

-1-

Date: <u>9/7/00</u>	Express Mail Label No. <u>EL552573612VS</u>
---------------------	---

Inventors: Eric S. Lander, Michele Cargill, James S. Ireland, Stacey
Bolk, George Q. Daley and Jeanette J. McCarthy
Attorney's Docket No.: 2825.1027-001

SINGLE NUCLEOTIDE POLYMORPHISMS IN GENES

RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Serial No. 60/153,357, filed September 10, 2000, U.S. Provisional Application Serial No. 60/220,947, filed July 26, 2000, and U.S. Provisional Application Serial No. 60/225,724, filed August 16, 2000, the entire teachings of all of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

The genomes of all organisms undergo spontaneous mutation in the course of their continuing evolution, generating variant forms of progenitor nucleic acid sequences (Gusella, *Ann. Rev. Biochem.* 55, 831-854 (1986)). The variant form may confer an evolutionary advantage or disadvantage relative to a progenitor form, or may be neutral. In some instances, a variant form confers a lethal disadvantage and is not transmitted to subsequent generations of the organism. In other instances, a variant form confers an evolutionary advantage to the species and is eventually incorporated into the DNA of many or most members of the species and effectively becomes the progenitor form. In many instances, both progenitor and variant form(s) survive and co-

exist in a species population. The coexistence of multiple forms of a sequence gives rise to polymorphisms.

Several different types of polymorphism have been reported. A restriction fragment length polymorphism (RFLP) is a variation in DNA sequence that alters the length of a restriction fragment (Botstein *et al.*, *Am. J. Hum. Genet.* 32, 314-331 (1980)). The restriction fragment length polymorphism may create or delete a restriction site, thus changing the length of the restriction fragment. RFLPs have been widely used in human and animal genetic analyses (see WO 90/13668; W090/11369; Donis-Keller, *Cell* 51, 319-337 (1987); Lander *et al.*, *Genetics* 121, 85-99 (1989)).

When a heritable trait can be linked to a particular RFLP, the presence of the RFLP in an individual can be used to predict the likelihood that the animal will also exhibit the trait.

Other polymorphisms take the form of short tandem repeats (STRs) that include tandem di-, tri- and tetra-nucleotide repeated motifs. These tandem repeats are also referred to as variable number tandem repeat (VNTR) polymorphisms. VNTRs have been used in identity and paternity analysis (US 5,075,217; Armour *et al.*, *FEBS Lett.* 307, 113-115 (1992); Horn *et al.*, W0 91/14003; Jeffreys, EP 370,719), and in a large number of genetic mapping studies.

Other polymorphisms take the form of single nucleotide variations between individuals of the same species. Such polymorphisms are far more frequent than RFLPs, STRs and VNTRs. Some single nucleotide polymorphisms (SNP) occur in protein-coding nucleic acid sequences (coding sequence SNP (cSNP)), in which case, one of the polymorphic forms may give rise to the expression of a defective or otherwise variant protein and, potentially, a genetic disease. Examples of genes in which polymorphisms within coding sequences give rise to genetic disease include β -globin (sickle cell anemia), apoE4 (Alzheimer's Disease), Factor V Leiden (thrombosis), and CFTR (cystic fibrosis). cSNPs can alter the codon sequence of the gene and therefore specify an alternative amino acid. Such changes are called "missense" when another amino acid is substituted, and "nonsense" when the alternative codon specifies a

002000 2242990

stop signal in protein translation. When the cSNP does not alter the amino acid specified the cSNP is called "silent".

Other single nucleotide polymorphisms occur in noncoding regions. Some of these polymorphisms may also result in defective protein expression (e.g., as a result of defective splicing). Other single nucleotide polymorphisms have no phenotypic effects.

Single nucleotide polymorphisms can be used in the same manner as RFLPs and VNTRs, but offer several advantages. Single nucleotide polymorphisms occur with greater frequency and are spaced more uniformly throughout the genome than other forms of polymorphism. The greater frequency and uniformity of single nucleotide polymorphisms means that there is a greater probability that such a polymorphism will be found in close proximity to a genetic locus of interest than would be the case for other polymorphisms. The different forms of characterized single nucleotide polymorphisms are often easier to distinguish than other types of polymorphism (e.g., by use of assays employing allele-specific hybridization probes or primers).

Only a small percentage of the total repository of polymorphisms in humans and other organisms has been identified. The limited number of polymorphisms identified to date is due to the large amount of work required for their detection by conventional methods. For example, a conventional approach to identifying polymorphisms might be to sequence the same stretch of DNA in a population of individuals by dideoxy sequencing. In this type of approach, the amount of work increases in proportion to both the length of sequence and the number of individuals in a population and becomes impractical for large stretches of DNA or large numbers of persons.

SUMMARY OF THE INVENTION

Work described herein pertains to the identification of polymorphisms which can predispose individuals to disease, by resequencing large numbers of genes in a large number of individuals. Various genes from a number of individuals have been resequenced as described herein, and SNPs in these genes have been discovered (see the Table and Fig. 3). Some of these SNPs are cSNPs which specify a different amino acid

The invention relates to a gene which comprises a single nucleotide polymorphism at a specific location. In a particular embodiment the invention relates to the variant allele of a gene having a single nucleotide polymorphism, which variant allele differs from a reference allele by one nucleotide at the site(s) identified in the Table and Fig. 3. Complements of these nucleic acid sequences are also included. The nucleic acid molecules can be DNA or RNA, and can be double- or single-stranded. Nucleic acid molecules can be, for example, 5-10, 5-15, 10-20, 5-25, 10-30, 10-50 or 10-100 bases long.

The invention further provides allele-specific oligonucleotides that hybridize to the reference or variant allele of a gene comprising a single nucleotide polymorphism or to the complement thereof. These oligonucleotides can be probes or primers.

15 The invention further provides a method of analyzing a nucleic acid from an individual. The method determines which base is present at any one of the polymorphic sites shown in the Table and/or Fig. 3. Optionally, a set of bases occupying a set of the polymorphic sites shown in the Table and /or Fig. 3 is determined. This type of analysis can be performed on a number of individuals, who are tested for the presence of a

20 disease phenotype. The presence or absence of disease phenotype is then correlated with a base or set of bases present at the polymorphic site or sites in the individuals tested.

Thus, the invention further relates to a method of predicting the presence, absence, likelihood of the presence or absence, or severity of a particular phenotype or disorder associated with a particular genotype. The method comprises obtaining a nucleic acid sample from an individual and determining the identity of one or more bases (nucleotides) at polymorphic sites of genes described herein, wherein the presence of a particular base is correlated with a specified phenotype or disorder, thereby

predicting the presence, absence, likelihood of the presence or absence, or severity of the phenotype or disorder in the individual.

The thrombospondins are a family of extracellular matrix (ECM) glycoproteins that modulate many cell behaviors including adhesion, migration, and proliferation.

5 Thrombospondins (also known as thrombin sensitive proteins or TSPs) are large molecular weight glycoproteins composed of three identical disulfide-linked polypeptide chains. The results described herein also reveal an important association between alterations, particularly SNPs, in TSP genes, particularly TSP-1 and TSP-4, and vascular disease. In particular, SNPs in these genes which are associated with
10 premature coronary artery disease (CAD)(or coronary heart disease) and myocardial infarction (MI) have been identified and represent a potentially vital marker of upstream biology influencing the complex process of atherosclerotic plaque generation and vulnerability.

Thus, the invention relates to the TSP gene SNPs identified as described herein,
15 both singly and in combination, as well as to the use of these SNPs, and others in TSP genes, particularly those nearby in linkage disequilibrium with these SNPs, for diagnosis, prediction of clinical course and treatment response for vascular disease, development of new treatments for vascular disease based upon comparison of the variant and normal versions of the gene or gene product, and development of cell-
20 culture based and animal models for research and treatment of vascular disease. The invention further relates to novel compounds and pharmaceutical compositions for use in the diagnosis and treatment of such disorders. In preferred embodiments, the vascular disease is CAD or MI.

The invention relates to isolated nucleic acid molecules comprising all or a
25 portion of the variant allele of TSP-1 (e.g., as exemplified by SEQ ID NO: 1), and to isolated nucleic acid molecules comprising all or a portion of the variant allele of TSP-4 (e.g., as exemplified by SEQ ID NO: 3). Preferred portions are at least 10 contiguous nucleotides and comprise the polymorphic site, e.g., a portion of SEQ ID NO: 1 which is at least 10 contiguous nucleotides and comprises the "G" at position 2210, or a

0065472-000400

portion of SEQ ID NO: 3 which is at least 10 contiguous nucleotides and comprises the "C" at position 1186. The invention further relates to isolated gene products, e.g., polypeptides or proteins, which are encoded by a nucleic acid molecule comprising all or a portion of the variant allele of TSP-1 or TSP-4 (e.g., SEQ ID NO: 1 or SEQ ID NO: 3, respectively). The invention also relates to nucleic acid molecules which hybridize to and/or share identity with the variant alleles identified herein (or their complements) and which also comprise the variant nucleotide at the SNP site.

The invention further relates to isolated proteins or polypeptides comprising all or a portion of the variant amino acid sequence of TSP-1 (e.g., as exemplified by SEQ ID NO: 2), and to isolated proteins or polypeptides comprising all or a portion of the variant amino acid sequence of TSP-4 (e.g., as exemplified by SEQ ID NO: 4). Preferred polypeptides are at least 10 contiguous amino acids and comprise the polymorphic amino acid, e.g., a portion of SEQ ID NO: 2 which is at least 10 contiguous amino acids and comprises the serine at residue 700, or a portion of SEQ ID NO: 4 which is at least 10 contiguous amino acids and comprises the proline at residue 387. The invention further relates to isolated nucleic acid molecules encoding such proteins and polypeptides, as well as to antibodies which bind, e.g., specifically, to such proteins and polypeptides.

The invention further relates to a method of diagnosing or aiding in the diagnosis of a disorder associated with the presence of one or more of (a) a G at nucleotide position 2210 of SEQ ID NO: 1; or (b) a C at nucleotide position 1186 of SEQ ID NO: 3 in an individual. The method comprises obtaining a nucleic acid sample from the individual and determining the nucleotide present at one or more of the indicated nucleotide positions, wherein presence of one or more of (a) a G at nucleotide position 2210 of SEQ ID NO: 1; or (b) a C at nucleotide position 1186 of SEQ ID NO: 3 is indicative of increased likelihood of said disorder in the individual as compared with an appropriate control, e.g., an individual having the reference nucleotide at one or more of said positions. In a particular embodiment the disorder is a vascular disease selected from the group consisting of atherosclerosis, coronary heart or artery disease, MI,

stroke, peripheral vascular diseases, venous thromboembolism and pulmonary embolism. In a preferred embodiment, the vascular disease is selected from the group consisting of CAD and MI.

The invention further relates to a method of diagnosing or aiding in the diagnosis
5 of a disorder associated with one or more of (a) a G at nucleotide position 2210 of SEQ ID NO: 1; or (b) a C at nucleotide position 1186 of SEQ ID NO: 3 in an individual. The method comprises obtaining a nucleic acid sample from the individual and determining the nucleotide present at one or more of the indicated nucleotide positions, wherein presence of one or more of (a) an A at nucleotide position 2210 of SEQ ID NO: 1; or (b)
10 a G at nucleotide position 1186 of SEQ ID NO: 3 is indicative of decreased likelihood of said disorder in the individual as compared with an appropriate control, e.g., an individual having the variant nucleotide at said position. In a particular embodiment the disorder is a vascular disease selected from the group consisting of atherosclerosis, coronary heart or artery disease, MI, stroke, peripheral vascular diseases, venous
15 thromboembolism and pulmonary embolism. In a preferred embodiment, the vascular disease is selected from the group consisting of CAD and MI.

In one embodiment, the invention relates to a method for predicting the likelihood that an individual will have a vascular disease (or aiding in the diagnosis of a vascular disease), comprising the steps of obtaining a DNA sample from an individual to be
20 assessed and determining the nucleotide present at one or more of nucleotide positions 2210 of SEQ ID NO: 1 or 1186 of SEQ ID NO: 3. The presence of the reference nucleotide at one or more of these positions indicates that the individual has a lower likelihood of having a vascular disease than an individual having the variant nucleotide at one or more of these positions, or a lower likelihood of having severe symptomology.
25 In a particular embodiment, the individual is an individual at risk for development of a vascular disease.

The invention further relates to a method of diagnosing or aiding in the diagnosis of a disorder associated with the presence of one or more of (a) a serine at amino acid position 700 of SEQ ID NO: 2; or (b) a proline at amino acid position 387 of SEQ ID

0065472.00000

NO: 4 in an individual. The method comprises obtaining a biological sample containing the TSP-1 and/or TSP-4 protein or relevant portion thereof from the individual and determining the amino acid present at one or more of the indicated amino acid positions, wherein presence of one or more of (a) a serine at amino acid position 700 of SEQ ID
5 NO: 2; or (b) a proline at amino acid position 387 of SEQ ID NO: 4 is indicative of increased likelihood of said disorder in the individual as compared with an appropriate control, *e.g.*, an individual having the reference amino acid at one or more of said positions.

The invention further relates to a method of diagnosing or aiding in the diagnosis
10 of a disorder associated with one or more of (a) a serine at amino acid position 700 of SEQ ID NO: 2; or (b) a proline at amino acid position 387 of SEQ ID NO: 4 in an individual. The method comprises obtaining a biological sample containing the TSP-1 and/or TSP-4 protein or relevant portion thereof from the individual and determining the amino acid present at one or more of the indicated amino acid positions, wherein
15 presence of one or more of (a) an asparagine at amino acid position 700 of SEQ ID NO: 2; or (b) an alanine at amino acid position 387 of SEQ ID NO: 4 is indicative of decreased likelihood of said disorder in the individual as compared with an appropriate control, *e.g.*, an individual having the variant amino acid at one or more of said positions.

20 In one embodiment, the invention relates to a method for predicting the likelihood that an individual will have a vascular disease (or aiding in the diagnosis of a vascular disease), comprising the steps of obtaining a biological sample comprising the TSP-1 and/or TSP-4 protein or relevant portion thereof from an individual to be assessed and determining the amino acid present at one or more of amino acid positions 700 of SEQ
25 ID NO: 2 or 387 of SEQ ID NO: 4. The presence of the reference amino acid at one or more of these positions indicates that the individual has a lower likelihood of having a vascular disease than an individual having the variant amino acid at one or more of these positions, or a lower likelihood of having severe symptomology. In a particular

004000" 2442550

embodiment, the individual is an individual at risk for development of a vascular disease.

In another embodiment, the invention relates to pharmaceutical compositions comprising a reference TSP-1 and/or TSP-4 gene or gene product, or active portion thereof, for use in the treatment of vascular diseases. The invention further relates to the use of agonists and antagonists of TSP-1 and TSP-4 activity for use in the treatment of vascular diseases. In a particular embodiment the vascular disease is selected from the group consisting of atherosclerosis, coronary heart or artery disease, MI, stroke, peripheral vascular diseases, venous thromboembolism and pulmonary embolism. In a preferred embodiment, the vascular disease is selected from the group consisting of CAD and MI.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A-1D show the reference nucleotide (SEQ ID NO: 1) and amino acid (SEQ ID NO: 2) sequences for TSP-1.

Figs. 2A-2C show the reference nucleotide (SEQ ID NO: 3) and amino acid (SEQ ID NO: 4) sequences for TSP-4.

Fig. 3 shows a table providing detailed information about the SNPs identified herein. Column one shows the internal polymorphism identifier. Column two shows the accession number for the reference sequence in the TIGR database (http://www.tigr.org/tdb/hgi/searching/hgi_reports.html). Column three shows the nucleotide position for the SNP site. Column four shows the gene in which the polymorphism was identified. Column five shows the polymorphic site and additional flanking sequence on each side of the polymorphism. Column six shows the type of mutation produced by the polymorphism. Columns seven and eight show the reference and alternate (variant) nucleotides, respectively, for the SNP. Columns nine and ten show the reference and alternate (variant) amino acids, respectively, encoded by the alleles of the gene.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a gene which comprises a single nucleotide polymorphism (SNP) at a specific location. The gene which includes the SNP has at least two alleles, referred to herein as the reference allele and the variant allele. The reference allele (prototypical or wild type allele) has been designated arbitrarily and typically corresponds to the nucleotide sequence of the gene which has been deposited with GenBank or TIGR under a given Accession number. The variant allele differs from the reference allele by one nucleotide at the site(s) identified in the Table. The present invention also relates to variant alleles of the described genes and to complements of the variant alleles. The invention also relates to nucleic acid molecules which hybridize to and/or share identity with the variant alleles identified herein (or their complements) and which also comprise the variant nucleotide at the SNP site.

The invention further relates to portions of the variant alleles and portions of complements of the variant alleles which comprise (encompass) the site of the SNP and are at least 5 nucleotides in length. Portions can be, for example, 5-10, 5-15, 10-20, 5-25, 10-30, 10-50 or 10-100 bases long. For example, a portion of a variant allele which is 21 nucleotides in length includes the single nucleotide polymorphism (the nucleotide which differs from the reference allele at that site) and twenty additional nucleotides which flank the site in the variant allele. These nucleotides can be on one or both sides of the polymorphism. Polymorphisms which are the subject of this invention are defined in the Table with respect to the reference sequence deposited in GenBank or TIGR under the Accession number indicated. For example, the invention relates to a portion of a gene (e.g., AT3) having a nucleotide sequence as deposited in GenBank (e.g., U11270) comprising a single nucleotide polymorphism at a specific position (e.g., nucleotide 11918). The reference nucleotide for AT3 is shown in column 8, and the variant nucleotide is shown in column 9 of the Table. The nucleotide sequences of the invention can be double- or single-stranded.

The invention further provides allele-specific oligonucleotides that hybridize to the reference or variant allele of a gene comprising a single nucleotide polymorphism or to the complement thereof. These oligonucleotides can be probes or primers.

The invention further provides a method of analyzing a nucleic acid from an
5 individual. The method determines which base is present at any one of the polymorphic sites shown in the Table and/or Fig. 3. Optionally, a set of bases occupying a set of the polymorphic sites shown in the Table and/or Fig. 3 is determined. This type of analysis can be performed on a number of individuals, who are tested for the presence of a disease phenotype. The presence or absence of disease phenotype is then correlated
10 with a base or set of bases present at the polymorphic site or sites in the individuals tested.

Thus, the invention further relates to a method of predicting the presence, absence, likelihood of the presence or absence, or severity of a particular phenotype or disorder associated with a particular genotype. The method comprises obtaining a
15 nucleic acid sample from an individual and determining the identity of one or more bases (nucleotides) at polymorphic sites of genes described herein, wherein the presence of a particular base is correlated with a specified phenotype or disorder, thereby predicting the presence, absence, likelihood of the presence or absence, or severity of the phenotype or disorder in the individual.

20 DEFINITIONS

A nucleic acid molecule or oligonucleotide can be DNA or RNA, and single- or double-stranded. Nucleic acid molecules and oligonucleotides can be naturally occurring or synthetic, but are typically prepared by synthetic means. Preferred nucleic acid molecules and oligonucleotides of the invention include segments of DNA, or their
25 complements, which include any one of the polymorphic sites shown in the Table. The segments can be between 5 and 250 bases, and, in specific embodiments, are between 5-10, 5-20, 10-20, 10-50, 20-50 or 10-100 bases. For example, the segment can be 21

bases. The polymorphic site can occur within any position of the segment. The segments can be from any of the allelic forms of DNA shown in the Table.

As used herein, the terms "nucleotide", "base" and "nucleic acid" are intended to be equivalent. The terms "nucleotide sequence", "nucleic acid sequence", "nucleic acid molecule" and "segment" are intended to be equivalent.

Hybridization probes are oligonucleotides which bind in a base-specific manner to a complementary strand of nucleic acid. Such probes include peptide nucleic acids, as described in Nielsen *et al.*, *Science* 254, 1497-1500 (1991). Probes can be any length suitable for specific hybridization to the target nucleic acid sequence. The most appropriate length of the probe may vary depending upon the hybridization method in which it is being used; for example, particular lengths may be more appropriate for use in microfabricated arrays, while other lengths may be more suitable for use in classical hybridization methods. Such optimizations are known to the skilled artisan. Suitable probes and primers can range from about 5 nucleotides to about 30 nucleotides in length. For example, probes and primers can be 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 25, 26, 28 or 30 nucleotides in length. The probe or primer preferably overlaps at least one polymorphic site occupied by any of the possible variant nucleotides. The nucleotide sequence can correspond to the coding sequence of the allele or to the complement of the coding sequence of the allele.

As used herein, the term "primer" refers to a single-stranded oligonucleotide which acts as a point of initiation of template-directed DNA synthesis under appropriate conditions (*e.g.*, in the presence of four different nucleoside triphosphates and an agent for polymerization, such as DNA or RNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer, but typically ranges from 15 to 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template, but must be sufficiently complementary to hybridize with a template. The term primer site refers to the area of the target DNA to which a

primer hybridizes. The term primer pair refers to a set of primers including a 5' (upstream) primer that hybridizes with the 5' end of the DNA sequence to be amplified and a 3' (downstream) primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

- 5 As used herein, linkage describes the tendency of genes, alleles, loci or genetic markers to be inherited together as a result of their location on the same chromosome. It can be measured by percent recombination between the two genes, alleles, loci or genetic markers.

- 10 As used herein, polymorphism refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic marker or site is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency of greater than 1%, and more preferably greater than 10% or 20% of a selected population. A polymorphic locus may be as small as one base pair. Polymorphic markers include restriction fragment length polymorphisms, variable
15 number of tandem repeats (VNTR's), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide repeats, simple sequence repeats, and insertion elements such as Alu. The first identified allelic form is arbitrarily designated as the reference form and other allelic forms are designated as alternative or variant alleles. The allelic form occurring most frequently in a selected population is
20 sometimes referred to as the wildtype form. Diploid organisms may be homozygous or heterozygous for allelic forms. A diallelic or biallelic polymorphism has two forms. A triallelic polymorphism has three forms.

- 25 Work described herein pertains to the resequencing of large numbers of genes in a large number of individuals to identify polymorphisms which can predispose individuals to disease. For example, polymorphisms in genes which are expressed in liver may predispose individuals to disorders of the liver. By altering amino acid sequence, SNPs may alter the function of the encoded proteins. The discovery of the SNP facilitates biochemical analysis of the variants and the development of assays to characterize the variants and to screen for pharmaceutical that would interact directly

002000 244590 0953472 090700

with on or another form of the protein. SNPs (including silent SNPs) also enable the development of specific DNA, RNA, or protein-based diagnostics that detect the presence or absence of the polymorphism in particular conditions.

5 A single nucleotide polymorphism occurs at a polymorphic site occupied by a single nucleotide, which is the site of variation between allelic sequences. The site is usually preceded by and followed by highly conserved sequences of the allele (e.g., sequences that vary in less than 1/100 or 1/1000 members of the populations).

10 A single nucleotide polymorphism usually arises due to substitution of one nucleotide for another at the polymorphic site. A transition is the replacement of one purine by another purine or one pyrimidine by another pyrimidine. A transversion is the replacement of a purine by a pyrimidine or vice versa. Single nucleotide polymorphisms can also arise from a deletion of a nucleotide or an insertion of a nucleotide relative to a reference allele. Typically the polymorphic site is occupied by a base other than the reference base. For example, where the reference allele contains the
15 base "T" at the polymorphic site, the altered allele can contain a "C", "G" or "A" at the polymorphic site.

20 The invention also relates to nucleic acid molecules which hybridize to the variant alleles identified herein (or their complements) and which also comprise the variant nucleotide at the SNP site. Hybridizations are usually performed under stringent conditions, for example, at a salt concentration of no more than 1 M and a temperature of at least 25°C. For example, conditions of 5X SSPE (750 mM NaCl, 50 mM NaPhosphate, 5 mM EDTA, pH 7.4) and a temperature of 25-30°C, or equivalent conditions, are suitable for allele-specific probe hybridizations. Equivalent conditions can be determined by varying one or more of the parameters given as an example, as
25 known in the art, while maintaining a similar degree of identity or similarity between the target nucleotide sequence and the primer or probe used.

The invention also relates to nucleic acid molecules which share substantial sequence identity to the variant alleles identified herein (or their complements) and which also comprise the variant nucleotide at the SNP site. Particularly preferred are

004000 2445960

nucleic acid molecules and fragments which have at least about 60%, preferably at least about 70, 80 or 85%, more preferably at least about 90%, even more preferably at least about 95%, and most preferably at least about 98% identity with nucleic acid molecules described herein. The percent identity of two nucleotide or amino acid sequences can
5 be determined by aligning the sequences for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first sequence). The nucleotides or amino acids at corresponding positions are then compared, and the percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical positions/total # of positions x 100). In certain
10 embodiments, the length of a sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 60%, and even more preferably at least 70%, 80% or 90% of the length of the reference sequence. The actual comparison of the two sequences can be accomplished by well-known methods, for example, using a mathematical algorithm. A preferred, non-limiting example of such a mathematical
15 algorithm is described in Karlin *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) as described in Altschul *et al.*, *Nucleic Acids Res.*, 25:389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. In
20 one embodiment, parameters for sequence comparison can be set at score=100, wordlength=12, or can be varied (*e.g.*, W=5 or W=20).

The term "isolated" is used herein to indicate that the material in question exists in a physical milieu distinct from that in which it occurs in nature. For example, an isolated nucleic acid of the invention may be substantially isolated with respect to the
25 complex cellular milieu in which it naturally occurs. In some instances, the isolated material will form part of a composition (for example, a crude extract containing other substances), buffer system or reagent mix. In other circumstance, the material may be purified to essential homogeneity, for example as determined by PAGE or column

00657473-000700

chromatography such as HPLC. Preferably, an isolated nucleic acid comprises at least about 50, 80 or 90 percent (on a molar basis) of all macromolecular species present.

I. Novel Polymorphisms of the Invention

Some of the novel polymorphisms of the invention are shown in the Table.

- 5 Columns one and two show designations for the indicated polymorphism. Column three shows the Genbank or TIGR Accession number for the wild type (or reference) allele. Column four shows the location of the polymorphic site in the nucleic acid sequence with reference to the Genbank or TIGR sequence shown in column three. Column five shows common names for the gene in which the polymorphism is located.
- 10 Column six shows the polymorphism and a portion of the 3' and 5' flanking sequence of the gene. Column seven shows the type of mutation; N, non-sense, S, silent, M, missense. Columns eight and nine show the reference and alternate nucleotides, respectively, at the polymorphic site. Columns ten and eleven show the reference and alternate amino acids, respectively, encoded by the reference and variant, respectively,
- 15 alleles. Other novel polymorphisms of the invention are shown in Fig. 3.

II. Analysis of Polymorphisms

A. Preparation of Samples

- Polymorphisms are detected in a target nucleic acid from an individual being analyzed. For assay of genomic DNA, virtually any biological sample (other than pure
- 20 red blood cells) is suitable. For example, convenient tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, buccal, skin and hair. For assay of cDNA or mRNA, the tissue sample must be obtained from an organ in which the target nucleic acid is expressed. For example, if the target nucleic acid is a cytochrome P450, the liver is a suitable source.

- 25 Many of the methods described below require amplification of DNA from target samples. This can be accomplished by e.g., PCR. *See generally PCR Technology: Principles and Applications for DNA Amplification* (ed. H.A. Erlich, Freeman Press,

NY, NY, 1992); *PCR Protocols: A Guide to Methods and Applications* (eds. Innis, *et al.*, Academic Press, San Diego, CA, 1990); Mattila *et al.*, *Nucleic Acids Res.* 19, 4967 (1991); Eckert *et al.*, *PCR Methods and Applications* 1, 17 (1991); *PCR* (eds. McPherson *et al.*, IRL Press, Oxford); and U.S. Patent 4,683,202.

- 5 Other suitable amplification methods include the ligase chain reaction (LCR) (see Wu and Wallace, *Genomics* 4, 560 (1989), Landegren *et al.*, *Science* 241, 1077 (1988), transcription amplification (Kwoh *et al.*, *Proc. Natl. Acad. Sci. USA* 86, 1173 (1989)), and self-sustained sequence replication (Guatelli *et al.*, *Proc. Nat. Acad. Sci. USA*, 87, 1874 (1990)) and nucleic acid based sequence amplification (NASBA). The latter two
- 10 amplification methods involve isothermal reactions based on isothermal transcription, which produce both single stranded RNA (ssRNA) and double stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

B. Detection of Polymorphisms in Target DNA

- There are two distinct types of analysis of target DNA for detecting
- 15 polymorphisms. The first type of analysis, sometimes referred to as *de novo* characterization, is carried out to identify polymorphic sites not previously characterized (i.e., to identify new polymorphisms). This analysis compares target sequences in different individuals to identify points of variation, i.e., polymorphic sites. By analyzing groups of individuals representing the greatest ethnic diversity among humans
- 20 and greatest breed and species variety in plants and animals, patterns characteristic of the most common alleles/haplotypes of the locus can be identified, and the frequencies of such alleles/haplotypes in the population can be determined. Additional allelic frequencies can be determined for subpopulations characterized by criteria such as geography, race, or gender. The *de novo* identification of polymorphisms of the
- 25 invention is described in the Examples section. The second type of analysis determines which form(s) of a characterized (known) polymorphism are present in individuals under test. There are a variety of suitable procedures, which are discussed in turn.

1. Allele-Specific Probes

The design and use of allele-specific probes for analyzing polymorphisms is described by e.g., Saiki *et al.*, *Nature* 324, 163-166 (1986); Dattagupta, EP 235,726, Saiki, WO 89/11548. Allele-specific probes can be designed that hybridize to a
5 segment of target DNA from one individual but do not hybridize to the corresponding segment from another individual due to the presence of different polymorphic forms in the respective segments from the two individuals. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe
10 hybridizes to only one of the alleles. Some probes are designed to hybridize to a segment of target DNA such that the polymorphic site aligns with a central position (e.g., in a 15-mer at the 7 position; in a 16-mer, at either the 8 or 9 position) of the probe. This design of probe achieves good discrimination in hybridization between different allelic forms.
15 Allele-specific probes are often used in pairs, one member of a pair showing a perfect match to a reference form of a target sequence and the other member showing a perfect match to a variant form. Several pairs of probes can then be immobilized on the same support for simultaneous analysis of multiple polymorphisms within the same target sequence.

2. Tiling Arrays

20 The polymorphisms can also be identified by hybridization to nucleic acid arrays, some examples of which are described in WO 95/11995. One form of such arrays is described in the Examples section in connection with de novo identification of polymorphisms. The same array or a different array can be used for analysis of
25 characterized polymorphisms. WO 95/11995 also describes subarrays that are optimized for detection of a variant form of a precharacterized polymorphism. Such a subarray contains probes designed to be complementary to a second reference sequence, which is an allelic variant of the first reference sequence. The second group of probes is

designed by the same principles as described in the Examples, except that the probes exhibit complementarity to the second reference sequence. The inclusion of a second group (or further groups) can be particularly useful for analyzing short subsequences of the primary reference sequence in which multiple mutations are expected to occur

5 within a short distance commensurate with the length of the probes (e.g., two or more mutations within 9 to 21 bases).

3. Allele-Specific Primers

An allele-specific primer hybridizes to a site on target DNA overlapping a polymorphism and only primes amplification of an allelic form to which the primer

10 exhibits perfect complementarity. See Gibbs, *Nucleic Acid Res.* 17, 2427-2448 (1989). This primer is used in conjunction with a second primer which hybridizes at a distal site. Amplification proceeds from the two primers, resulting in a detectable product which indicates the particular allelic form is present. A control is usually performed with a second pair of primers, one of which shows a single base mismatch at the

15 polymorphic site and the other of which exhibits perfect complementarity to a distal site. The single-base mismatch prevents amplification and no detectable product is formed. The method works best when the mismatch is included in the 3'-most position of the oligonucleotide aligned with the polymorphism because this position is most destabilizing to elongation from the primer (see, e.g., WO 93/22456).

20 4. Direct-Sequencing

The direct analysis of the sequence of polymorphisms of the present invention can be accomplished using either the dideoxy chain termination method or the Maxam - Gilbert method (see Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2nd Ed., CSHP, New York 1989); Zyskind *et al.*, *Recombinant DNA Laboratory Manual*,

25 (Acad. Press, 1988)).

5. Denaturing Gradient Gel Electrophoresis

Amplification products generated using the polymerase chain reaction can be analyzed by the use of denaturing gradient gel electrophoresis. Different alleles can be identified based on the different sequence-dependent melting properties and

- 5 electrophoretic migration of DNA in solution. Erlich, ed., *PCR Technology, Principles and Applications for DNA Amplification*, (W.H. Freeman and Co, New York, 1992), Chapter 7.

6. Single-Strand Conformation Polymorphism Analysis

- Alleles of target sequences can be differentiated using single-strand conformation polymorphism analysis, which identifies base differences by alteration in
10 electrophoretic migration of single stranded PCR products, as described in Orita *et al.*, *Proc. Nat. Acad. Sci.* 86, 2766-2770 (1989). Amplified PCR products can be generated as described above, and heated or otherwise denatured, to form single stranded amplification products. Single-stranded nucleic acids may refold or form secondary
15 structures which are partially dependent on the base sequence. The different electrophoretic mobilities of single-stranded amplification products can be related to base-sequence differences between alleles of target sequences.

7. Single-Base Extension

- An alternative method for identifying and analyzing polymorphisms is based on
20 single-base extension (SBE) of a fluorescently-labeled primer coupled with fluorescence resonance energy transfer (FRET) between the label of the added base and the label of the primer. Typically, the method, such as that described by Chen *et al.*, (*PNAS* 94:10756-61 (1997), incorporated herein by reference) uses a locus-specific oligonucleotide primer labeled on the 5' terminus with 5-carboxyfluorescein (FAM).
25 This labeled primer is designed so that the 3' end is immediately adjacent to the polymorphic site of interest. The labeled primer is hybridized to the locus, and single base extension of the labeled primer is performed with fluorescently labeled

dideoxyribonucleotides (ddNTPs) in dye-terminator sequencing fashion, except that no deoxyribonucleotides are present. An increase in fluorescence of the added ddNTP in response to excitation at the wavelength of the labeled primer is used to infer the identity of the added nucleotide.

5 III. Methods of Use

After determining polymorphic form(s) present in an individual at one or more polymorphic sites, this information can be used in a number of methods.

A. Forensics

Determination of which polymorphic forms occupy a set of polymorphic sites in
10 an individual identifies a set of polymorphic forms that distinguishes the individual.
See generally National Research Council, *The Evaluation of Forensic DNA Evidence*
(Eds. Pollard *et al.*, National Academy Press, DC, 1996). The more sites that are
analyzed, the lower the probability that the set of polymorphic forms in one individual
is the same as that in an unrelated individual. Preferably, if multiple sites are analyzed,
15 the sites are unlinked. Thus, polymorphisms of the invention are often used in
conjunction with polymorphisms in distal genes. Preferred polymorphisms for use in
forensics are biallelic because the population frequencies of two polymorphic forms can
usually be determined with greater accuracy than those of multiple polymorphic forms
at multi-allelic loci.

20 The capacity to identify a distinguishing or unique set of forensic markers in an
individual is useful for forensic analysis. For example, one can determine whether a
blood sample from a suspect matches a blood or other tissue sample from a crime scene
by determining whether the set of polymorphic forms occupying selected polymorphic
sites is the same in the suspect and the sample. If the set of polymorphic markers does
25 not match between a suspect and a sample, it can be concluded (barring experimental
error) that the suspect was not the source of the sample. If the set of markers does
match, one can conclude that the DNA from the suspect is consistent with that found at

00657472-00000

10

Both Heterozygotes: $p(\text{AB}+\text{BA})=2xy = 2x(1-x)$

$$p(\text{ID}) = (x^2)^2 + (2xy)^2 + (y^2)^2.$$
$$p(\mathbb{ID}) = x^4 + (2xy)^2 + (2yz)^2 + (2xz)^2 + z^4 + y^4$$

25 The cumulative probability of identity (cum p(ID)) for each of multiple unlinked loci is determined by multiplying the probabilities provided by each locus.

$$\text{cum } p(\text{ID}) = p(\text{ID1})p(\text{ID2})p(\text{ID3}).... p(\text{IDn})$$

cum p(nonID) = 1-cum p(ID).

B. Paternity Testing

15 If the set of polymorphisms in the child attributable to the father does not match
the set of polymorphisms of the putative father, it can be concluded, barring
experimental error, that the putative father is not the real father. If the set of
polymorphisms in the child attributable to the father does match the set of
polymorphisms of the putative father, a statistical calculation can be performed to
20 determine the probability of coincidental match.

$$p(\text{exc}) = xy(1-xy)$$

(At a triallelic site $p(\text{exc}) = xy(1-xy) + yz(1-yz) + xz(1-xz) + 3xyz(1-xyz)$), where x, y and z are the respective population frequencies of alleles A, B and C).

The probability of non-exclusion is

$$p(\text{non-exc}) = 1 - p(\text{exc})$$

The cumulative probability of non-exclusion (representing the value obtained when n loci are used) is thus:

$$5 \quad \text{cum } p(\text{non-exc}) = p(\text{non-exc1})p(\text{non-exc2})p(\text{non-exc3})\dots p(\text{non-excn})$$

The cumulative probability of exclusion for n loci (representing the probability that a random male will be excluded)

$$\text{cum } p(\text{exc}) = 1 - \text{cum } p(\text{non-exc}).$$

- If several polymorphic loci are included in the analysis, the cumulative
- 10 probability of exclusion of a random male is very high. This probability can be taken into account in assessing the liability of a putative father whose polymorphic marker set matches the child's polymorphic marker set attributable to his/her father.

C. Correlation of Polymorphisms with Phenotypic Traits

- The polymorphisms of the invention may contribute to the phenotype of an
- 15 organism in different ways. Some polymorphisms occur within a protein coding sequence and contribute to phenotype by affecting protein structure. The effect may be neutral, beneficial or detrimental, or both beneficial and detrimental, depending on the circumstances. For example, a heterozygous sickle cell mutation confers resistance to malaria, but a homozygous sickle cell mutation is usually lethal. Other polymorphisms
- 20 occur in noncoding regions but may exert phenotypic effects indirectly via influence on replication, transcription, and translation. A single polymorphism may affect more than one phenotypic trait. Likewise, a single phenotypic trait may be affected by polymorphisms in different genes. Further, some polymorphisms predispose an individual to a distinct mutation that is causally related to a certain phenotype.

- 25 Phenotypic traits include diseases that have known but hitherto unmapped genetic components (e.g., agammaglobulinemia, diabetes insipidus, Lesch-Nyhan syndrome, muscular dystrophy, Wiskott-Aldrich syndrome, Fabry's disease, familial hypercholesterolemia, polycystic kidney disease, hereditary spherocytosis, von

Willebrand's disease, tuberous sclerosis, hereditary hemorrhagic telangiectasia, familial colonic polyposis, Ehlers-Danlos syndrome, osteogenesis imperfecta, and acute intermittent porphyria). Phenotypic traits also include symptoms of, or susceptibility to, multifactorial diseases of which a component is or may be genetic, such as autoimmune diseases, inflammation, cancer, diseases of the nervous system, and infection by pathogenic microorganisms. Some examples of autoimmune diseases include rheumatoid arthritis, multiple sclerosis, diabetes (insulin-dependent and non-independent), systemic lupus erythematosus and Graves disease. Some examples of cancers include cancers of the bladder, brain, breast, colon, esophagus, kidney, leukemia, liver, lung, oral cavity, ovary, pancreas, prostate, skin, stomach and uterus. Phenotypic traits also include characteristics such as longevity, appearance (e.g., baldness, obesity), strength, speed, endurance, fertility, and susceptibility or receptivity to particular drugs or therapeutic treatments.

The correlation of one or more polymorphisms with phenotypic traits can be facilitated by knowledge of the gene product of the wild type (reference) gene. The genes in which cSNPs of the present invention have been identified are genes which have been previously sequenced and characterized in one of their allelic forms.

Correlation is performed for a population of individuals who have been tested for the presence or absence of a phenotypic trait of interest and for polymorphic markers sets. To perform such analysis, the presence or absence of a set of polymorphisms (i.e. a polymorphic set) is determined for a set of the individuals, some of whom exhibit a particular trait, and some of which exhibit lack of the trait. The alleles of each polymorphism of the set are then reviewed to determine whether the presence or absence of a particular allele is associated with the trait of interest. Correlation can be performed by standard statistical methods such as a K-squared test and statistically significant correlations between polymorphic form(s) and phenotypic characteristics are noted. For example, it might be found that the presence of allele A1 at polymorphism A correlates with heart disease. As a further example, it might be found that the combined

presence of allele A1 at polymorphism A and allele B1 at polymorphism B correlates with increased milk production of a farm animal.

Such correlations can be exploited in several ways. In the case of a strong correlation between a set of one or more polymorphic forms and a disease for which
 5 treatment is available, detection of the polymorphic form set in a human or animal patient may justify immediate administration of treatment, or at least the institution of regular monitoring of the patient. Detection of a polymorphic form correlated with serious disease in a couple contemplating a family may also be valuable to the couple in their reproductive decisions. For example, the female partner might elect to undergo *in*
 10 *vitro* fertilization to avoid the possibility of transmitting such a polymorphism from her husband to her offspring. In the case of a weaker, but still statistically significant correlation between a polymorphic set and human disease, immediate therapeutic intervention or monitoring may not be justified. Nevertheless, the patient can be motivated to begin simple life-style changes (e.g., diet, exercise) that can be
 15 accomplished at little cost to the patient but confer potential benefits in reducing the risk of conditions to which the patient may have increased susceptibility by virtue of variant alleles. Identification of a polymorphic set in a patient correlated with enhanced receptiveness to one of several treatment regimes for a disease indicates that this treatment regime should be followed.

20 For animals and plants, correlations between characteristics and phenotype are useful for breeding for desired characteristics. For example, Beitz *et al.*, US 5,292,639 discuss use of bovine mitochondrial polymorphisms in a breeding program to improve milk production in cows. To evaluate the effect of mtDNA D-loop sequence polymorphism on milk production, each cow was assigned a value of 1 if variant or 0 if
 25 wildtype with respect to a prototypical mitochondrial DNA sequence at each of 17 locations considered. Each production trait was analyzed individually with the following animal model:

$$Y_{ijkpn} = \mu + YS_i + P_j + X_k + \beta_1 + \dots \beta_{17} + PE_n + a_n + e_p$$

where Y_{ijknp} is the milk, fat, fat percentage, SNF, SNF percentage, energy concentration, or lactation energy record; μ is an overall mean; YS_i is the effect common to all cows calving in year-season; X_k is the effect common to cows in either the high or average selection line; β_1 to β_{17} are the binomial regressions of production record on mtDNA D-loop sequence polymorphisms; PE_n is permanent environmental effect common to all records of cow n ; a_n is effect of animal n and is composed of the additive genetic contribution of sire and dam breeding values and a Mendelian sampling effect; and e_p is a random residual. It was found that eleven of seventeen polymorphisms tested influenced at least one production trait. Bovines having the best polymorphic forms for milk production at these eleven loci are used as parents for breeding the next generation of the herd.

D. Genetic Mapping of Phenotypic Traits

The previous section concerns identifying correlations between phenotypic traits and polymorphisms that directly or indirectly contribute to those traits. The present section describes identification of a physical linkage between a genetic locus associated with a trait of interest and polymorphic markers that are not associated with the trait, but are in physical proximity with the genetic locus responsible for the trait and co-segregate with it. Such analysis is useful for mapping a genetic locus associated with a phenotypic trait to a chromosomal position, and thereby cloning gene(s) responsible for the trait. See Lander *et al.*, *Proc. Natl. Acad. Sci. (USA)* 83, 7353-7357 (1986); Lander *et al.*, *Proc. Natl. Acad. Sci. (USA)* 84, 2363-2367 (1987); Donis-Keller *et al.*, *Cell* 51, 319-337 (1987); Lander *et al.*, *Genetics* 121, 185-199 (1989)). Genes localized by linkage can be cloned by a process known as directional cloning. See Wainwright, *Med. J. Australia* 159, 170-174 (1993); Collins, *Nature Genetics* 1, 3-6 (1992).

Linkage studies are typically performed on members of a family. Available members of the family are characterized for the presence or absence of a phenotypic trait and for a set of polymorphic markers. The distribution of polymorphic markers in an informative meiosis is then analyzed to determine which polymorphic markers co-

002060 244560

segregate with a phenotypic trait. See, e.g., Kerem *et al.*, *Science* 245, 1073-1080 (1989); Monaco *et al.*, *Nature* 316, 842 (1985); Yamoka *et al.*, *Neurology* 40, 222-226 (1990); Rossiter *et al.*, *FASEB Journal* 5, 21-27 (1991).

Linkage is analyzed by calculation of LOD (log of the odds) values. A lod value
5 is the relative likelihood of obtaining observed segregation data for a marker and a
genetic locus when the two are located at a recombination fraction θ , versus the
situation in which the two are not linked, and thus segregating independently
(Thompson & Thompson, *Genetics in Medicine* (5th ed, W.B. Saunders Company,
Philadelphia, 1991); Strachan, "Mapping the human genome" in *The Human Genome*
10 (BIOS Scientific Publishers Ltd, Oxford), Chapter 4). A series of likelihood ratios are
calculated at various recombination fractions (θ), ranging from $\theta = 0.0$ (coincident loci)
to $\theta = 0.50$ (unlinked). Thus, the likelihood at a given value of θ is: probability of data
if loci linked at θ to probability of data if loci unlinked. The computed likelihoods are
usually expressed as the \log_{10} of this ratio (i.e., a lod score). For example, a lod score
15 of 3 indicates 1000:1 odds against an apparent observed linkage being a coincidence.
The use of logarithms allows data collected from different families to be combined by
simple addition. Computer programs are available for the calculation of lod scores for
differing values of θ (e.g., LIPED, MLINK (Lathrop, *Proc. Nat. Acad. Sci. (USA)* 81,
3443-3446 (1984)). For any particular lod score, a recombination fraction may be
20 determined from mathematical tables. See Smith *et al.*, *Mathematical tables for*
research workers in human genetics (Churchill, London, 1961); Smith, *Ann. Hum.*
Genet. 32, 127-150 (1968). The value of θ at which the lod score is the highest is
considered to be the best estimate of the recombination fraction.

Positive lod score values suggest that the two loci are linked, whereas negative
25 values suggest that linkage is less likely (at that value of θ) than the possibility that the
two loci are unlinked. By convention, a combined lod score of +3 or greater (equivalent
to greater than 1000:1 odds in favor of linkage) is considered definitive evidence that
two loci are linked. Similarly, by convention, a negative lod score of -2 or less is taken
as definitive evidence against linkage of the two loci being compared. Negative linkage

data are useful in excluding a chromosome or a segment thereof from consideration. The search focuses on the remaining non-excluded chromosomal locations.

IV. Modified Polypeptides and Gene Sequences

The invention further provides variant forms of nucleic acids and corresponding
5 proteins. The nucleic acids comprise one of the sequences described in the Table, column 5, in which the polymorphic position is occupied by one of the alternative bases for that position. Some nucleic acids encode full-length variant forms of proteins. Similarly, variant proteins have the prototypical amino acid sequences encoded by nucleic acid sequences shown in the Table, column 5, (read so as to be in-frame with the
10 full-length coding sequence of which it is a component) except at an amino acid encoded by a codon including one of the polymorphic positions shown in the Table. That position is occupied by the amino acid coded by the corresponding codon in any of the alternative forms shown in the Table.

Variant genes can be expressed in an expression vector in which a variant gene is
15 operably linked to a native or other promoter. Usually, the promoter is a eukaryotic promoter for expression in a mammalian cell. The transcription regulation sequences typically include a heterologous promoter and optionally an enhancer which is recognized by the host. The selection of an appropriate promoter, for example trp, lac, phage promoters, glycolytic enzyme promoters and tRNA promoters, depends on the
20 host selected. Commercially available expression vectors can be used. Vectors can include host-recognized replication systems, amplifiable genes, selectable markers, host sequences useful for insertion into the host genome, and the like.

The means of introducing the expression construct into a host cell varies depending upon the particular construction and the target host. Suitable means include
25 fusion, conjugation, transfection, transduction, electroporation or injection, as described in Sambrook, *supra*. A wide variety of host cells can be employed for expression of the variant gene, both prokaryotic and eukaryotic. Suitable host cells include bacteria such as *E. coli*, yeast, filamentous fungi, insect cells, mammalian cells, typically

00567473-000700

The protein may be isolated by conventional means of protein biochemistry and purification to obtain a substantially pure product, i.e., 80, 95 or 99% free of cell component contaminants, as described in Jacoby, *Methods in Enzymology* Volume 104, Academic Press, New York (1984); Scopes, *Protein Purification, Principles and Practice*, 2nd Edition, Springer-Verlag, New York (1987); and Deutscher (ed), *Guide to Protein Purification, Methods in Enzymology*, Vol. 182 (1990). If the protein is secreted, it can be isolated from the supernatant in which the host cell is grown. If not secreted, the protein can be isolated from a lysate of the host cells.

The invention further provides transgenic nonhuman animals capable of expressing an exogenous variant gene and/or having one or both alleles of an endogenous variant gene inactivated. Expression of an exogenous variant gene is usually achieved by operably linking the gene to a promoter and optionally an enhancer, and microinjecting the construct into a zygote. *See Hogan et al.*, "Manipulating the Mouse Embryo, A Laboratory Manual," Cold Spring Harbor Laboratory. Inactivation of endogenous variant genes can be achieved by forming a transgene in which a cloned variant gene is inactivated by insertion of a positive selection marker. *See Capecchi, Science* 244, 1288-1292 (1989). The transgene is then introduced into an embryonic stem cell, where it undergoes homologous recombination with an endogenous variant gene. Mice and other rodents are preferred animals. Such animals provide useful drug screening systems.

In addition to substantially full-length polypeptides expressed by variant genes, the present invention includes biologically active fragments of the polypeptides, or analogs thereof, including organic molecules which simulate the interactions of the peptides. Biologically active fragments include any portion of the full-length

polypeptide which confers a biological function on the variant gene product, including ligand binding, and antibody binding. Ligand binding includes binding by nucleic acids, proteins or polypeptides, small biologically active molecules, or large cellular structures.

- 5 Polyclonal and/or monoclonal antibodies that specifically bind to variant gene products but not to corresponding prototypical gene products are also provided. Antibodies can be made by injecting mice or other animals with the variant gene product or synthetic peptide fragments thereof. Monoclonal antibodies are screened as are described, for example, in Harlow & Lane, *Antibodies, A Laboratory Manual*, Cold
10 Spring Harbor Press, New York (1988); Goding, *Monoclonal antibodies, Principles and Practice* (2d ed.) Academic Press, New York (1986). Monoclonal antibodies are tested for specific immunoreactivity with a variant gene product and lack of immunoreactivity to the corresponding prototypical gene product. These antibodies are useful in diagnostic assays for detection of the variant form, or as an active ingredient in a
15 pharmaceutical composition.

V. Kits

- The invention further provides kits comprising at least one allele-specific oligonucleotide as described herein. Often, the kits contain one or more pairs of allele-specific oligonucleotides hybridizing to different forms of a polymorphism. In some
20 kits, the allele-specific oligonucleotides are provided immobilized to a substrate. For example, the same substrate can comprise allele-specific oligonucleotide probes for detecting at least 10, 100 or all of the polymorphisms shown in the Table. Optional additional components of the kit include, for example, restriction enzymes, reverse-transcriptase or polymerase, the substrate nucleoside triphosphates, means used to label
25 (for example, an avidin-enzyme conjugate and enzyme substrate and chromogen if the label is biotin), and the appropriate buffers for reverse transcription, PCR, or hybridization reactions. Usually, the kit also contains instructions for carrying out the methods.

000000-242560

The thrombospondins are a family of extracellular matrix (ECM) glycoproteins that modulate many cell behaviors including adhesion, migration, and proliferation. Thrombospondins (also known as thrombin sensitive proteins or TSPs) are large molecular weight glycoproteins composed of three identical disulfide-linked

5 polypeptide chains. TSPs are stored in the alpha-granules of platelets and secreted by a variety of mesenchymal and epithelial cells (Majack *et al.*, *Cell Membrane* 3:57-77 (1987)). Platelets secrete TSPs when activated in the blood by such physiological agonists such as thrombin. TSPs have lectin properties and a broad function in the regulation of fibrinolysis and as a component of the ECM, and are one of a group of

10 ECM proteins which have adhesive properties. TSPs bind to fibronectin and fibrinogen (Lahav *et al.*, *Eur J Biochem* 145:151-6 (1984)), and these proteins are known to be involved in platelet adhesion to substratum and platelet aggregation (Leung, *J Clin Invest* 74:1764-1772 (1986)).

Recent work has implicated TSPs in response of cells to growth factors.

15 Submitogenic doses of PDGF induce a rapid but transitory, increase in TSP synthesis and secretion by rat aortic smooth muscle cells (Majack *et al.*, *J Biol Chem* 101:1059-70 (1985)). PDGF responsiveness to TSP synthesis in glial cells has also been shown (Asch *et al.*, *Proc Natl Acad Sci* 83:2904-8 (1986)). TSP mRNA levels rise rapidly in response to PDGF (Majack *et al.*, *J Biol Chem* 262:8821-5 (1987)). TSPs act

20 synergistically with epidermal growth factor to increase DNA synthesis in smooth muscle cells (Majack *et al.*, *Proc Natl Acad Sci* 83:9050-4 (1986)), and monoclonal antibodies to TSPs inhibit smooth muscle cell proliferation (Majack *et al.*, *J Biol Chem* 106:415-22 (1988)). TSPs modulate local adhesions in endothelial cells, and TSPs, particularly TSP-1 primarily derived from platelet granules, are known to be an

25 important activator of transforming growth factor beta-1 (TGFB-1) (Crawford *et al.*, *Cell* 93:1159 (1998)) and appear to be a potential link between platelet-thrombosis and development of atherosclerosis.

To determine pivotal genes associated with premature coronary artery disease, we analyzed DNA from 347 patients with MI or coronary revascularization before age 40

(men) or 45 (women) and 422 general population controls. Cases were drawn (one per family) from a retrospective collection of sibling pairs with premature CAD. Controls were ascertained through random-digit dialing. Both cases and controls were Caucasian. A complete database of phenotypic and laboratory variables for the affected
5 patients afforded logistic regression to control for age, diabetes, body mass index, gender.

Thrombospondin (TSP) 4 and 1 emerged as important SNPs associated with premature CAD and MI. For CAD, 148 of 347 patients carried at least one copy of the TSP-4 variant compared with 142 of 422 control subjects; adjusted odds ratio 1.47,
10 $p=0.01$. For premature MI, the association was even stronger: 91 of 187 cases vs. 142 of 422 controls had the variant; adjusted odds ratio 2.08, $p=0.0003$. The TSP-1 SNP was rare. Nonetheless, homozygosity for the variant allele gave an adjusted odds ratio of 9.5, $p=.04$.

Specific reference nucleotide (SEQ ID NO: 1) and amino acid (SEQ ID NO: 2)
15 sequences for TSP-1 are shown in Figs. 1A-1D. Specific reference nucleotide (SEQ ID NO: 3) and amino acid (SEQ ID NO: 4) sequences for TSP-4 are shown in Figs. 2A-2C. It is understood that the invention is not limited by these exemplified reference sequences, as variants of these sequences which differ at locations other than the SNP sites identified herein can also be utilized. The skilled artisan can readily determine the
20 SNP sites in these other reference sequences which correspond to the SNP sites identified herein by aligning the sequence of interest with the reference sequences specifically disclosed herein, and programs for performing such alignments are commercially available. For example, the ALIGN program in the GCG software package can be used, utilizing a PAM120 weight residue table, a gap length penalty of
25 12 and a gap penalty of 4, for example.

Two SNPs have been specifically studied as described herein. The first (G334u4) is a change from A (reference nucleotide) to G (alternate or variant nucleotide) at nucleotide position 2210 of the nucleic acid sequence of TSP-1 (Figs. 1A-1D), resulting in a missense amino acid mutation from asparagine (reference) to serine (alternate) at

amino acid 700. The second SNP (G355u2) is a change from G (reference) to C (alternate) at nucleotide position 1186 of the nucleic acid sequence of TSP-4 (Figs. 2A-2C), resulting in a missense amino acid alteration from alanine (reference) to proline (alternate) at amino acid 387. With respect to the G355u2 SNP, individuals with CAD
5 carried at least one copy of the variant "C" allele more frequently than control individuals (43% as compared with 34%). With respect to the G355u2 SNP, individuals with MI carried at least one copy of the variant "C" allele more frequently than control individuals (49% as compared with 34%). With respect to the G334u4 SNP, individuals with CAD carried two copies of the variant "G" allele more frequently than
10 control individuals (1.7% as compared with 0.2%). With respect to the G334u4 SNP, individuals with MI carried two copies of the variant "G" allele more frequently than control individuals (2% as compared with 0.2%).

As used herein, the term "polymorphism" refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic
15 marker or site is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency of greater than 1%, and more preferably greater than 10% or 20% of a selected population. A polymorphic locus may be as small as one base pair, in which case it is referred to as a single nucleotide polymorphism (SNP).

Thus, the invention relates to a method for predicting the likelihood that an
20 individual will have a vascular disease, or for aiding in the diagnosis of a vascular disease, or predicting the likelihood of having altered symptomology associated with a vascular disease, comprising the steps of obtaining a DNA sample from an individual to be assessed and determining the nucleotide present at one or more of nucleotide positions 2210 of the TSP-1 gene or 1186 of the TSP-4 gene. In a preferred
25 embodiment, the nucleotides present at both of these nucleotide positions are determined. In one embodiment the TSP-1 gene has the nucleotide sequence of SEQ ID NO: 1 and the TSP-4 gene has the nucleotide sequence of SEQ ID NO: 3. The presence of one or more of a G (the variant nucleotide) at position 2210 of SEQ ID NO: 1 or a C (the variant nucleotide) at position 1186 of SEQ ID NO: 1186 indicates that the

individual has a greater likelihood of having a vascular disease, or a greater likelihood of having severe symptomology associated with a vascular disease, than if that individual had the reference nucleotide at one or more of these positions. Conversely, the presence of one or more of an A (the reference nucleotide) at position 2210 of SEQ ID NO: 1 or a G (the reference nucleotide) at position 1186 of SEQ ID NO: 3 indicates that the individual has a reduced likelihood of having a vascular disease or a likelihood of having reduced symptomology associated with a vascular disease than if that individual had the variant nucleotide at one or more of these positions.

In a particular embodiment, the individual is an individual at risk for development of a vascular disease. In another embodiment the individual exhibits clinical symptomology associated with a vascular disease. In one embodiment, the individual has been clinically diagnosed as having a vascular disease. Vascular diseases include, but are not limited to, atherosclerosis, coronary heart disease, myocardial infarction (MI), stroke, peripheral vascular diseases, venous thromboembolism and pulmonary embolism. In preferred embodiments, the vascular disease is CAD or MI.

The genetic material to be assessed can be obtained from any nucleated cell from the individual. For assay of genomic DNA, virtually any biological sample (other than pure red blood cells) is suitable. For example, convenient tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, skin and hair. For assay of cDNA or mRNA, the tissue sample must be obtained from a tissue or organ in which the target nucleic acid is expressed.

Many of the methods described herein require amplification of DNA from target samples. This can be accomplished by e.g., PCR. See generally *PCR Technology: Principles and Applications for DNA Amplification* (ed. H.A. Erlich, Freeman Press, NY, NY, 1992); *PCR Protocols: A Guide to Methods and Applications* (eds. Innis, et al., Academic Press, San Diego, CA, 1990); Mattila et al., *Nucleic Acids Res.* 19, 4967 (1991); Eckert et al., *PCR Methods and Applications* 1, 17 (1991); PCR (eds. McPherson et al., IRL Press, Oxford); and U.S. Patent 4,683,202.

Other suitable amplification methods include the ligase chain reaction (LCR) (see Wu and Wallace, *Genomics* 4, 560 (1989), Landegren *et al.*, *Science* 241, 1077 (1988), transcription amplification (Kwoh *et al.*, *Proc. Natl. Acad. Sci. USA* 86, 1173 (1989)), and self-sustained sequence replication (Guatelli *et al.*, *Proc. Nat. Acad. Sci. USA*, 87, 1874 (1990)) and nucleic acid based sequence amplification (NASBA). The latter two
5 amplification methods involve isothermal reactions based on isothermal transcription, which produce both single stranded RNA (ssRNA) and double stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

The nucleotide which occupies the polymorphic site of interest (e.g., nucleotide
10 position 2210 in TSP-1 and/or nucleotide position 1186 in TSP-4) can be identified by a variety of methods, such as Southern analysis of genomic DNA; direct mutation analysis by restriction enzyme digestion; Northern analysis of RNA; denaturing high pressure liquid chromatography (DHPLC); gene isolation and sequencing; hybridization of an allele-specific oligonucleotide with amplified gene products; single base extension
15 (SBE). In a preferred embodiment, determination of the allelic form of TSP is carried out using SBE-FRET methods as described herein, or using chip-based oligonucleotide arrays as described herein.

The invention also relates to a method for predicting the likelihood that an individual will have a vascular disease, or for aiding in the diagnosis of a vascular
20 disease, or predicting the likelihood of having altered symptomology associated with a vascular disease, comprising the steps of obtaining a biological sample comprising TSP-1 and/or TSP-4 protein or relevant portion thereof from an individual to be assessed and determining the amino acid present at one or more of amino acid positions 700 of the TSP-1 gene product (e.g., as exemplified by SEQ ID NO: 2) or 387 of the TSP-4 gene
25 product (e.g., as exemplified by SEQ ID NO: 4). In a preferred embodiment, the amino acids present at both of these amino acid positions are determined. As used herein, the term "relevant portion" of the TSP-1 and TSP-4 proteins is intended to encompass any portion of the protein which comprises the polymorphic amino acid positions. The presence of one or more of a serine (the variant amino acid) at position 700 of SEQ ID

00667422-000000

NO: 2, or a proline (the variant amino acid) at position 387 of SEQ ID NO: 4 indicates that the individual has a greater likelihood of having a vascular disease, or a greater likelihood of having severe symptomology associated with a vascular disease, than if that individual had the reference amino acid at one or more of these positions.

- 5 Conversely, the presence of one or more of an asparagine (the reference amino acid) at position 700 of SEQ ID NO: 2, or an alanine (the reference amino acid) at position 387 of SEQ ID NO: 4 indicates that the individual has a reduced likelihood of having a vascular disease or a likelihood of having reduced symptomology associated with a vascular disease, than if that individual had the variant amino acid at one or more of
10 these positions.

In a particular embodiment, the individual is an individual at risk for development of a vascular disease. In another embodiment the individual exhibits clinical symptomology associated with a vascular disease. In one embodiment, the individual has been clinically diagnosed as having a vascular disease.

- 15 In this embodiment of the invention, the biological sample contains protein molecules from the test subject. *In vitro* techniques for detection of protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. Furthermore, *in vivo* techniques for detection of protein include introducing into a subject a labeled anti-protein antibody. For example, the
20 antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Polyclonal and/or monoclonal antibodies that specifically bind to variant gene products but not to corresponding reference gene products, and vice versa, are also provided. Antibodies can be made by injecting mice or other animals with the variant gene product or synthetic peptide
25 fragments thereof comprising the variant portion. Monoclonal antibodies are screened as are described, for example, in Harlow & Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press, New York (1988); Goding, *Monoclonal antibodies, Principles and Practice* (2d ed.) Academic Press, New York (1986). Monoclonal antibodies are tested for specific immunoreactivity with a variant gene product and lack

of immunoreactivity to the corresponding prototypical gene product. These antibodies are useful in diagnostic assays for detection of the variant form, or as an active ingredient in a pharmaceutical composition.

The polymorphisms of the invention may be associated with vascular disease in
5 different ways. The polymorphisms may exert phenotypic effects indirectly via influence on replication, transcription, and translation. Additionally, the described polymorphisms may predispose an individual to a distinct mutation that is causally related to a certain phenotype, such as susceptibility or resistance to vascular disease and related disorders. The discovery of the polymorphisms and their correlation with
10 CAD and MI facilitates biochemical analysis of the variant and reference forms and the development of assays to characterize the variant and reference forms and to screen for pharmaceutical agents that interact directly with one or another form of the protein.

Alternatively, these particular polymorphisms may belong to a group of two or more polymorphisms in the TSP gene(s) which contributes to the presence, absence or
15 severity of vascular disease. An assessment of other polymorphisms within the TSP gene(s) can be undertaken, and the separate and combined effects of these polymorphisms, as well as alternations in other, distinct genes, on the vascular disease phenotype can be assessed.

Correlation between a particular phenotype, e.g., the CAD or MI phenotype, and
20 the presence or absence of a particular allele is performed for a population of individuals who have been tested for the presence or absence of the phenotype. Correlation can be performed by standard statistical methods such as a Chi-squared test and statistically significant correlations between polymorphic form(s) and phenotypic characteristics are noted. This correlation can be exploited in several ways. In the case
25 of a strong correlation between a particular polymorphic form, e.g., the variant allele for TSP-1 and/or TSP-4, and a disease for which treatment is available, detection of the polymorphic form in an individual may justify immediate administration of treatment, or at least the institution of regular monitoring of the individual. Detection of a polymorphic form correlated with a disorder in a couple contemplating a family may

006442-00000

also be valuable to the couple in their reproductive decisions. For example, the female partner might elect to undergo *in vitro* fertilization to avoid the possibility of transmitting such a polymorphism from her husband to her offspring. In the case of a weaker, but still statistically significant correlation between a polymorphic form and a particular disorder, immediate therapeutic intervention or monitoring may not be justified. Nevertheless, the individual can be motivated to begin simple life-style changes (e.g., diet modification, therapy or counseling) that can be accomplished at little cost to the individual but confer potential benefits in reducing the risk of conditions to which the individual may have increased susceptibility by virtue of the particular allele. Furthermore, identification of a polymorphic form correlated with enhanced receptiveness to one of several treatment regimes for a disorder indicates that this treatment regimen should be followed for the individual in question.

Furthermore, it may be possible to identify a physical linkage between a genetic locus associated with a trait of interest (e.g., CAD or MI) and polymorphic markers that are or are not associated with the trait, but are in physical proximity with the genetic locus responsible for the trait and co-segregate with it. Such analysis is useful for mapping a genetic locus associated with a phenotypic trait to a chromosomal position, and thereby cloning gene(s) responsible for the trait. See Lander *et al.*, *Proc. Natl. Acad. Sci. (USA)* 83, 7353-7357 (1986); Lander *et al.*, *Proc. Natl. Acad. Sci. (USA)* 84, 2363-2367 (1987); Donis-Keller *et al.*, *Cell* 51, 319-337 (1987); Lander *et al.*, *Genetics* 121, 185-199 (1989)). Genes localized by linkage can be cloned by a process known as directional cloning. See Wainwright, *Med. J. Australia* 159, 170-174 (1993); Collins, *Nature Genetics* 1, 3-6 (1992). Linkage studies are discussed in more detail above.

In another embodiment, the invention relates to pharmaceutical compositions comprising a reference TSP-1 and/or TSP-4 gene or gene product for use in the treatment of vascular disease, e.g., CAD and MI. As used herein, a reference TSP gene product is intended to mean gene products which are encoded by the reference allele of the TSP gene. In addition to substantially full-length polypeptides expressed by the genes, the present invention includes biologically active fragments of the polypeptides,

or analogs thereof, including organic molecules which simulate the interactions of the peptides. Biologically active fragments include any portion of the full-length polypeptide which confers a biological function on the variant gene product, including ligand binding, and antibody binding. Ligand binding includes binding by nucleic acids, proteins or polypeptides, small biologically active molecules, or large cellular structures.

For instance, the polypeptide or protein, or fragment thereof, of the present invention can be formulated with a physiologically acceptable medium to prepare a pharmaceutical composition. The particular physiological medium may include, but is not limited to, water, buffered saline, polyols (e.g., glycerol, propylene glycol, liquid polyethylene glycol) and dextrose solutions. The optimum concentration of the active ingredient(s) in the chosen medium can be determined empirically, according to procedures well known to medicinal chemists, and will depend on the ultimate pharmaceutical formulation desired. Methods of introduction of exogenous peptides at the site of treatment include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, oral and intranasal. Other suitable methods of introduction can also include rechargeable or biodegradable devices and slow release polymeric devices. The pharmaceutical compositions of this invention can also be administered as part of a combinatorial therapy with other agents and treatment regimens.

The invention further pertains to compositions, e.g., vectors, comprising a nucleotide sequence encoding reference or variant TSP-1 and/or TSP-4 gene products. For example, reference genes can be expressed in an expression vector in which a reference gene is operably linked to a native or other promoter. Usually, the promoter is a eukaryotic promoter for expression in a mammalian cell. The transcription regulation sequences typically include a heterologous promoter and optionally an enhancer which is recognized by the host. The selection of an appropriate promoter, for example trp, lac, phage promoters, glycolytic enzyme promoters and tRNA promoters, depends on the host selected. Commercially available expression vectors can be used. Vectors can

include host-recognized replication systems, amplifiable genes, selectable markers, host sequences useful for insertion into the host genome, and the like.

The means of introducing the expression construct into a host cell varies depending upon the particular construction and the target host. Suitable means include
5 fusion, conjugation, transfection, transduction, electroporation or injection, as described in Sambrook, *supra*. A wide variety of host cells can be employed for expression of the variant gene, both prokaryotic and eukaryotic. Suitable host cells include bacteria such as *E. coli*, yeast, filamentous fungi, insect cells, mammalian cells, typically immortalized, e.g., mouse, CHO, human and monkey cell lines and derivatives thereof.
10 Preferred host cells are able to process the variant gene product to produce an appropriate mature polypeptide. Processing includes glycosylation, ubiquitination, disulfide bond formation, general post-translational modification, and the like.

It is also contemplated that cells can be engineered to express the reference allele of the invention by gene therapy methods. For example, DNA encoding the reference
15 TSP gene product, or an active fragment or derivative thereof, can be introduced into an expression vector, such as a viral vector, and the vector can be introduced into appropriate cells in an animal. In such a method, the cell population can be engineered to inducibly or constitutively express active reference TSP gene product. In a preferred embodiment, the vector is delivered to the bone marrow, for example as described in
20 Corey *et al.* (*Science* 244:1275-1281 (1989)).

The invention further relates to the use of compositions (i.e., agonists) which enhance or increase the activity of the reference (or variant) TSP (e.g., TSP-1 or TSP-4) gene product, or a functional portion thereof, for use in the treatment of vascular disease. The invention also relates to the use of compositions (i.e., antagonists) which
25 reduce or decrease the activity of the variant (or reference) TSP (e.g., TSP-1 or TSP-4) gene product, or a functional portion thereof, for use in the treatment of vascular disease.

The invention also relates to constructs which comprise a vector into which a sequence of the invention has been inserted in a sense or antisense orientation. For

005472 000700
002000 22425960

example, a vector comprising a nucleotide sequence which is antisense to the variant TSP-1 or TSP-4 allele may be used as an antagonist of the activity of the TSP-1 or TSP-4 variant allele. Alternatively, a vector comprising a nucleotide sequence of the TSP-1 or TSP-4 reference allele may be used therapeutically to treat vascular diseases. As
5 used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of
10 autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors, expression vectors, are capable of directing the
15 expression of genes to which they are operably linked. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids (vectors). However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses) that serve equivalent functions.

20 Preferred recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell. This means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Within a recombinant
25 expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control

elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and
5 those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc.

The expression vectors of the invention can be introduced into host cells to
10 thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein. The recombinant expression vectors of the invention can be designed for expression of a polypeptide of the invention in prokaryotic or eukaryotic cells, *e.g.*, bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors), yeast cells or mammalian cells. Suitable host cells are discussed
15 further in Goeddel, *supra*. Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and
20 "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term
25 as used herein. A host cell can be any prokaryotic or eukaryotic cell. For example, a nucleic acid of the invention can be expressed in bacterial cells (*e.g.*, *E. coli*), insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including
5 calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*supra*), and other laboratory manuals.

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in
10 culture, can be used to produce (*i.e.*, express) a polypeptide of the invention. Accordingly, the invention further provides methods for producing a polypeptide using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of the invention (into which a recombinant expression vector encoding a polypeptide of the invention has been introduced) in a suitable medium such that the
15 polypeptide is produced. In another embodiment, the method further comprises isolating the polypeptide from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which a nucleic acid of the invention has been
20 introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous nucleotide sequences have been introduced into their genome or homologous recombinant animals in which endogenous nucleotide sequences have been altered. Such animals are useful for studying the function and/or activity of the nucleotide sequence and polypeptide encoded by the sequence and for identifying
25 and/or evaluating modulators of their activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the

00657472-000000

genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, an "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing a nucleic acid of the invention into the male pronuclei of a fertilized oocyte, *e.g.*, by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The sequence can be introduced as a transgene into the genome of a non-human animal. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of a polypeptide in particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, U.S. Patent No. 4,873,191 and in Hogan, *Manipulating the Mouse Embryo* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding the transgene can further be bred to other transgenic animals carrying other transgenes.

The invention also relates to the use of the variant and reference gene products to guide efforts to identify the causative mutation for vascular diseases or to identify or synthesize agents useful in the treatment of vascular diseases, *e.g.*, CAD and MI.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science*, 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then
5 tested for biological activity *in vitro*, or *in vitro* activity. Sites that are critical for polypeptide activity can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.*, 224:899-904 (1992); de Vos *et al.* *Science*, 255:306-312 (1992)).

Another aspect of the invention pertains to monitoring the influence of agents
10 (*e.g.*, drugs, compounds) on the expression or activity of proteins of the invention in clinical trials. An exemplary method for detecting the presence or absence of proteins or nucleic acids of the invention in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting the protein, or nucleic acid (*e.g.*, mRNA, genomic DNA) that
15 encodes the protein, such that the presence of the protein or nucleic acid is detected in the biological sample. A preferred agent for detecting mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to mRNA or genomic DNA sequences described herein, preferably in an allele-specific manner. The nucleic acid probe can be, for example, a full-length nucleic acid, or a portion thereof, such as an oligonucleotide
20 of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to appropriate mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

The invention also encompasses kits for detecting the presence of proteins or nucleic acid molecules of the invention in a biological sample. For example, the kit can
25 comprise a labeled compound or agent (*e.g.*, nucleic acid probe) capable of detecting protein or mRNA in a biological sample; means for determining the amount of protein or mRNA in the sample; and means for comparing the amount of protein or mRNA in the sample with a standard. The compound or agent can be packaged in a suitable

container. The kit can further comprise instructions for using the kit to detect protein or nucleic acid.

The following Examples are offered for the purpose of illustrating the present invention and are not to be construed to limit the scope of this invention. The teachings
5 of all references cited herein are hereby incorporated herein by reference.

EXAMPLES

Identification of Single Nucleotide Polymorphisms

The polymorphisms shown in the Table were identified by resequencing of target sequences from individuals of diverse ethnic and geographic backgrounds by
10 hybridization to probes immobilized to microfabricated arrays. The strategy and principles for design and use of such arrays are generally described in WO 95/11995.

A typical probe array used in this analysis has two groups of four sets of probes that respectively tile both strands of a reference sequence. A first probe set comprises a plurality of probes exhibiting perfect complementarity with one of the reference
15 sequences. Each probe in the first probe set has an interrogation position that corresponds to a nucleotide in the reference sequence. That is, the interrogation position is aligned with the corresponding nucleotide in the reference sequence, when the probe and reference sequence are aligned to maximize complementarity between the two. For each probe in the first set, there are three corresponding probes from three additional
20 probe sets. Thus, there are four probes corresponding to each nucleotide in the reference sequence. The probes from the three additional probe sets are identical to the corresponding probe from the first probe set except at the interrogation position, which occurs in the same position in each of the four corresponding probes from the four probe sets, and is occupied by a different nucleotide in the four probe sets. In the present
25 analysis, probes were 25 nucleotides long. Arrays tiled for multiple different reference sequences were included on the same substrate.

Publicly available sequences for a given gene were assembled into Gap4 (<http://www.biozentrum.unibas.ch/~biocomp/staden/Overview.html>). PCR primers covering each exon were designed using Primer 3 (<http://www-genome.wi.mit.edu/cgi-bin/primer/primer3.cgi>). Primers were not designed in regions where there were

5 sequence discrepancies between reads. Genomic DNA was amplified in at least 50 individuals using 2.5 pmol each primer, 1.5 mM MgCl₂, 100 μM dNTPs, 0.75 μM AmpliTaq GOLD polymerase, and 19 ng DNA in a 15 μl reaction. Reactions were assembled using a PACKARD MultiPROBE robotic pipetting station and then put in MJ 96-well tetrad thermocyclers (96°C for 10 minutes, followed by 35 cycles of 96°C

10 for 30 seconds, 59°C for 2 minutes, and 72°C for 2 minutes). A subset of the PCR assays for each individual were run on 3% NuSieve gels in 0.5X TBE to confirm that the reaction worked.

For a given DNA, 5 μl (about 50 ng) of each PCR or RT-PCR product were pooled (Final volume = 150-200 μl). The products were purified using QiaQuick PCR

15 purification from Qiagen. The samples were eluted once in 35 μl sterile water and 4 μl 10X One-Phor-All buffer (Pharmacia). The pooled samples were digested with 0.2 μ DNaseI (Promega) for 10 minutes at 37°C and then labeled with 0.5 nmols biotin-N6-ddATP and 15 μ Terminal Transferase (GibcoBRL Life Technology) for 60 minutes at 37°C. Both fragmentation and labeling reactions were terminated by incubating the

20 pooled sample for 15 minutes at 100°C.

Low-density DNA chips (Affymetrix, CA) were hybridized following the manufacturer's instructions. Briefly, the hybridization cocktail consisted of 3M TMACl, 10 mM Tris pH 7.8, 0.01% Triton X-100, 100 mg/ml herring sperm DNA (Gibco BRL), 200 pM control biotin-labeled oligo. The processed PCR products were

25 denatured for 7 minutes at 100°C and then added to prewarmed (37°C) hybridization solution. The chips were hybridized overnight at 44°C. Chips were washed in 1X SSPET and 6X SSPET followed by staining with 2 μg/ml SARPE and 0.5 mg/ml acetylated BSA in 200 μl of 6X SSPET for 8 minutes at room temperature. Chips were scanned using a Molecular Dynamics scanner.

Chip image files were analyzed using Ulysses (Affymetrix, CA) which uses four algorithms to identify potential polymorphisms. Candidate polymorphisms were visually inspected and assigned a confidence value: high confidence candidates displayed all three genotypes, while likely candidates showed only two genotypes
5 (homozygous for reference sequence and heterozygous for reference and variant). Some of the candidate polymorphisms were confirmed by ABI sequencing. Identified polymorphisms were compared to several databases to determine if they were novel. Results are shown in the Table.

Association of Thrombospondin Gene Polymorphisms with Vascular Disease

10 To determine pivotal genes associated with premature coronary artery disease, we analyzed DNA from 347 patients with MI or coronary revascularization before age 40 (men) or 45 (women) and 422 general population controls. Cases were drawn (one per family) from a retrospective collection of sibling pairs with premature CAD. Controls were ascertained through random-digit dialing. Both cases and controls were
15 Caucasian. A complete database of phenotypic and laboratory variables for the affected patients afforded logistic regression to control for age, diabetes, body mass index, gender.

Thrombospondin (TSP) 4 and 1 emerged as important SNPs associated with premature CAD and MI. For CAD, 148 of 347 patients carried at least one copy of the
20 TSP-4 variant compared with 142 of 422 control subjects; adjusted odds ratio 1.47, $p=0.01$. For premature MI, the association was even stronger: 91 of 187 cases vs. 142 of 422 controls had the variant; adjusted odds ratio 2.08, $p=0.0003$. The TSP-1 SNP was rare. Nonetheless, homozygosity for the variant allele gave an adjusted odds ratio of 9.5, $p=.04$.

Poly ID	WIAF ID	Genbank or TIGR Accession Number	Position in Sequence	Gene Description	Flanking Seq	Mutation Type	Ref NT	Alt NT	Ref AA	Alt AA
AT3a7	WIAF-13246	U11270	11918	AT3, antithrombin III	CTGCAGGAGT [G/A] GCTGGATGAA	N	G	A	W	*
DRD5u22	WIAF-12913	M67439	310	DRD1, dopamine receptor D1	CATCTGGACC [C/T] TGTCTGGCAA	S	C	T	L	L
DRD5u23	WIAF-12914	M67439	332	DRD1, dopamine receptor D1	GTGCTGTGT [G/C] CGCAGCCATC	M	G	C	C	S
DRD5u24	WIAF-12915	M67439	369	DRD1, dopamine receptor D1	TGCGCGCCAA [C/G] ATGACCAACG	M	C	G	N	K
DRD5u25	WIAF-12916	M67439	522	DRD1, dopamine receptor D1	TGTCTCCAC [T/C] GCCTCCATCC	S	T	C	T	T
DRD5u26	WIAF-12917	M67439	953	DRD1, dopamine receptor D1	GCAGAGCACG [C/T] GCAGAGCTGC	M	C	T	A	V
DRD5u27	WIAF-12918	M67439	635	DRD1, dopamine receptor D1	ATGCTCGGCC [T/C] GGCATGGACC	M	T	C	L	P
DRD5u28	WIAF-12919	M67439	606	DRD1, dopamine receptor D1	GCAAGATGAC [T/C] CAGCGCATGG	S	T	C	T	T
DRD5u29	WIAF-12920	M67439	845	DRD1, dopamine receptor D1	TCGCTCATCA [G/A] CTTCTACATC	M	G	A	S	N
DRD5u30	WIAF-12921	M67439	720	DRD1, dopamine receptor D1	GGGGGGGGCT [G/T] GACCTGCCAA	S	G	T	L	L
DRD5u31	WIAF-12922	M67439	1044	DRD1, dopamine receptor D1	AGACCTGTC [G/A] GTGATCATGG	S	G	A	S	S
DRD5u32	WIAF-12923	M67439	766	DRD1, dopamine receptor D1	GGAGGAGGAC [T/G] TTTGGAGCC	M	T	G	F	V
DRD5u33	WIAF-12924	M67439	777	DRD1, dopamine receptor D1	TTTGGAGCC [C/T] GACGTGAATG	S	C	T	P	P
DRD5u34	WIAF-12925	M67439	786	DRD1, dopamine receptor D1	CCGACGTGAA [T/G] GCAGAGAACT	M	T	G	N	K

DRD5u35	WIAF-12926	M67439	887	DRD1, dopamine receptor D1	ACCTACAGC [G/A] CATCTACCGC	M	G	A	R	H
DRD5u36	WIAF-12927	M67439	1279	DRD1, dopamine receptor D1	GTGAGCCAC [T/G] TCTGCTCCCG	M	T	G	F	V
DRD5u37	WIAF-12928	M67439	1370	DRD1, dopamine receptor D1	GAAATCGCAG [C/T] TGCCTACATC	M	C	T	A	V
DRD5u38	WIAF-12929	M67439	1500	DRD1, dopamine receptor D1	ACCTGTGTGC [T/A] GAGTCTGTCT	S	T	A	A	A
DRD5u39	WIAF-12930	M67439	1338	DRD1, dopamine receptor D1	TCTCTACAA [C/T] CAAGACATCG	S	C	T	N	N
DRD5u40	WIAF-12931	M67439	1215	DRD1, dopamine receptor D1	CACTCAACCC [C/A] GTCATCTATG	S	C	A	P	P
DRD5u41	WIAF-12932	M67439	1242	DRD1, dopamine receptor D1	ACGCCGACTT [T/C] CAGAAAGGTGT	S	T	C	F	F
DRD5u42	WIAF-12933	M67439	1441	DRD1, dopamine receptor D1	CGAGGAGGAG [G/A] GTCCTTTTGA	M	G	A	G	S
DRD5u43	WIAF-12934	M67439	1460	DRD1, dopamine receptor D1	GATCGCATGT [T/C] CCAGATCTAT	M	T	C	F	S
DRD5u44	WIAF-12960	M67439	399	DRD1, dopamine receptor D1	TGTCTCTGGC [C/T] GTGCTGACC	S	C	T	A	A
DRD5u45	WIAF-12961	M67439	162	DRD1, dopamine receptor D1	TGCGGCCAGG [C/G] AGCAACGGCA	S	C	G	G	G
DRD5u46	WIAF-12962	M67439	195	DRD1, dopamine receptor D1	GGCAGTTGCG [T/G] CTATACCAGC	S	T	G	A	A
DRD5u47	WIAF-12963	M67439	264	DRD1, dopamine receptor D1	TGGGGCCCTC [A/G] CAGGTGGTCA	S	A	G	S	S
DRD5u48	WIAF-12964	M67439	465	DRD1, dopamine receptor D1	TGGCCGGTTA [C/T] TGGCCCTTTG	S	C	T	Y	Y
DRD5u49	WIAF-12965	M67439	511	DRD1, dopamine receptor D1	CTTCGACATC [A/T] TGTGCTCCAC	M	A	T	M	L
DRD5u50	WIAF-12966	M67439	557	DRD1, dopamine receptor D1	ATCAGCGTGG [A/G] CCGCTACTGG	M	A	G	D	G
DRD5u51	WIAF-12967	M67439	476	DRD1, dopamine receptor D1	TGGCCCTTTG [G/A] AGCGTTCTGC	M	G	A	G	E
DRD5u52	WIAF-12968	M67439	1004	DRD1, dopamine receptor D1	AGCTGCGCG [C/T] TTCCATCAAG	M	C	T	A	V
DRD5u53	WIAF-12969	M67439	1036	DRD1, dopamine receptor D1	GGTTCTCAAG [A/C] CCCTGTGGGT	M	A	C	T	P
DRD5u54	WIAF-12970	M67439	859	DRD1, dopamine receptor D1	CTACATCCCC [G/A] TTGCCATCAT	M	G	A	V	I
DRD5u55	WIAF-12971	M67439	931	DRD1, dopamine receptor D1	GATTCCTCC [C/T] TGGAGAGGGC	S	C	T	L	L

G10u1	WIAF-10234	J04111	1308	JUN, v-jun avian sarcoma virus 17 oncogene homolog	CCCTCAACGC[C/T]TCGTTCTCTCC	S	C	T	A	A
G10u2	WIAF-10235	J04111	1471	JUN, v-jun avian sarcoma virus 17 oncogene homolog	GCTGTCTCAAG[C/T]TGGCGTCGCC	S	C	T	L	L
G10u3	WIAF-10253	J04111	2010	JUN, v-jun avian sarcoma virus 17 oncogene homolog	TGGAGTCCCA[G/A]GAGCGGATCA	S	G	A	Q	Q
G1001u1	WIAF-13746	D26135	993	DGKG, diacylglycerol kinase, gamma (90kD)	CCCCAGTGGT[G/A]TACCTGAAGG	S	G	A	V	V
G1001u2	WIAF-13764	D26135	2313	DGKG, diacylglycerol kinase, gamma (90kD)	ATGTGATGAG[A/T]GAGAAACATC	M	A	T	R	S
G1002u1	WIAF-13918	X57206	334	ITPKB, inositol 1,4,5-trisphosphate 3-kinase B	CCCCAAGATC[A/C]GGACAAGCCT	M	A	C	Q	P
G1002u2	WIAF-13925	X57206	575	ITPKB, inositol 1,4,5-trisphosphate 3-kinase B	CCAACCTCAGC[T/C]TTCTGTCATA	S	T	C	A	A
G1004u1	WIAF-13567	L36151	1854	PIK4CA, phosphatidylinositol 4-kinase, catalytic, alpha polypeptide	GCCGCTCAGA[C/T]TCCGAGGATG	S	C	T	D	D
G1006u1	WIAF-12375	HT2690	858	PRKCA, protein kinase C, alpha	GGTACAAGTT[G/A]CTTAACCAAG	S	G	A	L	L
G1008u1	WIAF-12397	HT2136	300	PRKCZ, protein kinase C, zeta	CTGGCCTGCC[A/G]TGTCGGGAG	S	A	G	P	P
G1008u2	WIAF-12398	HT2136	246	PRKCZ, protein kinase C, zeta	AGTGCAGGGA[T/C]GAAGGCCTCA	S	T	C	D	D
G1008u3	WIAF-12399	HT2136	504	PRKCZ, protein kinase C, zeta	GCTCCACGG[C/T]CTCGTCCCGC	S	C	T	G	G
G1008u4	WIAF-12403	HT2136	807	PRKCZ, protein kinase C, zeta	AGAAGAATGA[C/T]CAAAATTACG	S	C	T	D	D
G1008u5	WIAF-12404	HT2136	1514	PRKCZ, protein kinase C, zeta	GGATTTCTG[A/T]CATCAAGTCC	M	A	T	D	V
G1008u6	WIAF-12412	HT2136	166	PRKCZ, protein kinase C, zeta	CAAGTGGGTG[G/A]ACAGCGAAGG	M	G	A	D	N
G1008u7	WIAF-12418	HT2136	560	PRKCZ, protein kinase C, zeta	TCCCAAGAGC[C/T]TCCAGTAGAC	M	C	T	P	L
G1009u1	WIAF-12396	L05186	2495	PTK2, PTK2 protein tyrosine kinase 2	TCATCAACAA[G/A]ATGAAACTGG	S	G	A	K	K
G1011u1	WIAF-11988	X07876	1250	WNT2, wingless-type MMTV integration site family member 2	TCCCATGTCA[C/A]CCGATGACC	M	C	A	T	N
G1011u2	WIAF-11997	X07876	788	WNT2, wingless-type MMTV integration site family member 2	GACTATGGGA[T/C]CAAATTGGC	M	T	C	I	T

G1011u3	WIAF-12014	X07876	WNT2, wingless-type MMTV 1338 integration site family member 2	TGCACACATG [C/A] AAGGCCCCCA	N	C	A	C	*
G1011u4	WIAF-13475	X07876	WNT2, wingless-type MMTV 856 integration site family member 2	CCTGATGAT [C/T] TTCACAACA	M	C	T	L	F
G1011u5	WIAF-13476	X07876	WNT2, wingless-type MMTV 958 integration site family member 2	GACATGCTGG [C/T] TGGCCATGGC	S	C	T	L	L
G1011u6	WIAF-13477	X07876	WNT2, wingless-type MMTV 789 integration site family member 2	ACTATGGGAT [C/T] AAATTGCCC	S	C	T	I	I
G1011u7	WIAF-13478	X07876	WNT2, wingless-type MMTV 823 integration site family member 2	TGCAAGGAA [A/G] GGAAGGAAA	M	A	G	R	G
G1012u1	WIAF-12408	HT48910	WNT2B, wingless-type MMTV 1574 integration site family, member 2B	ATACTTGCAA [A/G] GCCCCCAAGA	S	A	G	K	K
G1016a1	WIAF-12125	Z22534	793 ACVR1, activin A receptor, type I	GGCAAGGGA [A/G] AATGTCGG	S	A	G	E	E
G1016u2	WIAF-12392	Z22534	373 ACVR1, activin A receptor, type I	CTGCCAAGC [T/C] GTGGAGTGCT	S	T	C	A	A
G1018u1	WIAF-12413	X74210	ADCY2, adenylate cyclase 2 1150 (brain)	CAAAATTGCGA [G/T] TGGGTATTAA	M	G	T	V	L
G1019u1	WIAF-12394	U83867	SPTAN1, spectrin, alpha, non-erythrocytic 1 (alpha-fodrin) 5475	GGGACCTAAC [T/C] GSGGTGAGA	S	T	C	T	T
G1019u2	WIAF-12406	U83867	SPTAN1, spectrin, alpha, non-erythrocytic 1 (alpha-fodrin) 1223	GCCCTCATCA [A/G] TGCAGATGAG	M	A	G	N	S
G1019u3	WIAF-12409	U83867	SPTAN1, spectrin, alpha, non-erythrocytic 1 (alpha-fodrin) 3555	CTGAAGGTCT [T/C] ATGGCAGAGG	S	T	C	L	L
G1019u4	WIAF-12415	U83867	SPTAN1, spectrin, alpha, non-erythrocytic 1 (alpha-fodrin) 3369	TCCGTGAAGC [G/A] AATGAATAC	S	G	A	A	A
G1019u5	WIAF-12417	U83867	SPTAN1, spectrin, alpha, non-erythrocytic 1 (alpha-fodrin) 5839	TCAGACAGAC [T/A] TCACCGTCCA	M	T	A	F	I
G1022u1	WIAF-12393	U45945	ATP1B2, ATPase, Na+/K+ transporting, beta 2 polypeptide 631	CATGAATGTT [A/G] CCTGTGCTGG	M	A	G	T	A

G1022u2	WIAF-12400	U45945	432	ATP1B2, ATPase, Na+/K+ transporting, beta 2 polypeptide	GCCGCCCTGG[G/A]CGCTATTACG	S	G	A	G	G
G1023u1	WIAF-12401	D89722	395	ARNTL, aryl hydrocarbon receptor nuclear translocator-like	AACATTAAAG[G/C]GTGCCACCAA	M	G	C	G	R
G1023u2	WIAF-12407	D89722	681	ARNTL, aryl hydrocarbon receptor nuclear translocator-like	CTCATAGATG[C/T]AAAACTGGA	M	C	T	A	V
G1024u1	WIAF-12410	U85946	731	Homo sapiens brain secretory protein hSec10p (HSEC10) mRNA, complete cds.	GATAGATTTT[C/T]AGAAAGTTAAA	M	C	T	S	L
G1027u1	WIAF-12402	L47647	1135	CKB, creatine kinase, brain	TCGAGATGGA[A/G]CAGCGGCTGG	S	A	G	E	E
G1027u2	WIAF-12405	L47647	499	CKB, creatine kinase, brain	GGGAGCGCCG[A/C]GCCATCGAGA	S	A	C	R	R
G103u1	WIAF-10427	HT2269	335	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	GGGATCGCCA[T/C]GGGAACTCAA	S	T	C	H	H
G103u2	WIAF-10429	HT2269	1221	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	CCCTCCTTCT[C/T]CAAGAACTTT	M	C	T	P	S
G103u3	WIAF-10431	HT2269	1783	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	TCTCCAACTT[G/C]TACAAATTCT	M	G	C	C	S

G103u4	WIAF-10432	HT2269	2077	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	ACTGAATCTG [C/A]AGGCAGGAT	M	C	A	A	E
G103u5	WIAF-10446	HT2269	3338	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	AATTGAGCT [A/T]CTTGATAAGG	S	A	T	L	L
G103u6	WIAF-10447	HT2269	3487	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	TCAGAAATCAT [C/T]TGATGGATCT	M	C	T	S	F
G103u7	WIAF-10448	HT2269	3507	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	TTCAAGTGAA [C/G]ATGCTGAAAG	M	C	G	H	D
G103u8	WIAF-10457	HT2269	1388	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	CTCTTGACGA [T/G]GACGAAGATG	M	T	G	D	E
G103u9	WIAF-10458	HT2269	1362	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	CCGACTCTT [T/C]CAGCATTA	M	T	C	S	P

004000" 2442550

G103u10	WIAF-10459	HT2269	2357	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	CTGAGAAAGA [T/C] GCGGAAGATT	S	T	C	D	D
G103u11	WIAF-10462	HT2269	3109	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	TGGAACAGAA [C/T] GAAGACAGAT	M	C	T	T	M
G103u12	WIAF-10463	HT2269	3138	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	GTTTCCTGTA [T/C] TAAAGCAACT	S	T	C	L	L
G103u14	WIAF-10484	HT2269	3553	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	AGAACAGCTG [C/T] GAAAGAGCCA	M	C	T	A	V
G103u15	WIAF-10485	HT2269	1429	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	GATGTGCAGA [C/T] GGGAGGGCCA	M	C	T	T	M
G103u16	WIAF-12097	HT2269	3335	ERCC5, excision repair cross-complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome))	AAGAATTGGA [G/T] CTACTTGATA	M	G	T	E	D

G1030u1	WIAF-12411	U07358	ZPK, zipper (leucine) protein 203 kinase	ACACTTCTGA [C/T] TGCACCTCCG	S	C	T	D	D
G1030u2	WIAF-12416	U07358	ZPK, zipper (leucine) protein 1806 kinase	GCCACCCAT [G/T] AACTGGAGG	N	G	T	E	*
G1031a1	WIAF-12124	U87460	GPR37, G protein-coupled receptor 37 (endothelin receptor type B- 2825 like)	GAGTCACCAC [C/T] TTCACCTTAT	S	C	T	T	T
G1032u1	WIAF-12381	U57911	CL10RF8, chromosome 11 open 926 reading frame 8	ACGTACATCA [A/C] TGCCTCGACG	M	A	C	N	T
G1033u1	WIAF-12437	M65188	GJA1, gap junction protein, alpha 431 L, 43kD (connexin 43)	TCCTGTACCCA [C/T] ACTCTGTGAC	M	C	T	T	I
G1033u2	WIAF-12438	M65188	GJA1, gap junction protein, alpha 169 L, 43kD (connexin 43)	AGCAACATG [G/C] GTGACTGGAG	M	G	C	G	R
G1033u3	WIAF-12439	M65188	GJA1, gap junction protein, alpha 467 L, 43kD (connexin 43)	TATGTGATGC [G/A] AAAGGAAGAG	M	G	A	R	Q
G1033u4	WIAF-12440	M65188	GJA1, gap junction protein, alpha 263 L, 43kD (connexin 43)	TTCATTTTCC [G/A] AATCCTGCTG	M	G	A	R	Q
G1033u5	WIAF-12441	M65188	GJA1, gap junction protein, alpha 218 L, 43kD (connexin 43)	CAAGCCTACT [C/T] AACTGCTGGA	M	C	T	S	L
G1033u6	WIAF-12442	M65188	GJA1, gap junction protein, alpha 498 L, 43kD (connexin 43)	AGAAAGAGGA [A/G] GAACTCAAGG	S	A	G	E	E
G1033u7	WIAF-12465	M65188	GJA1, gap junction protein, alpha 550 L, 43kD (connexin 43)	GCACTTGAAG [C/A] AGATTGAGAT	M	C	A	Q	K
G1033u8	WIAF-12466	M65188	GJA1, gap junction protein, alpha 548 L, 43kD (connexin 43)	ATGCACCTTGA [A/G] GCAGATTGAG	M	A	G	K	R
G1033u9	WIAF-12486	M65188	GJA1, gap junction protein, alpha 933 L, 43kD (connexin 43)	CGCTGAGCCC [T/C] GCCAAAGACT	S	T	C	P	P
G1033u10	WIAF-12487	M65188	GJA1, gap junction protein, alpha 990 L, 43kD (connexin 43)	CCTCACCAC [C/T] GCTCCCTCT	S	C	T	T	T
G1033u11	WIAF-12488	M65188	GJA1, gap junction protein, alpha 1034 L, 43kD (connexin 43)	AAGCTGGTTA [C/A] TGGCGACAGA	M	C	A	T	N

G1033u12 G1033u13 G1033u14 G1033u15 G1033u16 G1033u17 G1033u18 G1033u19 G1033u20 G1033u21 G1033u22 G1034u1 G1034u2 G1034u3 G1034u4

G1033u12	WIAF-12489	M65188	GJA1, gap junction protein, alpha 1158 1, 43kD (connexin 43)	CTAACTCCCA [T/C] GCACAGCCTT	S	T	C	H	H
G1033u13	WIAF-12490	M65188	GJA1, gap junction protein, alpha 1222 1, 43kD (connexin 43)	TGCACATGAA [T/C] TACAGCCACT	S	T	C	L	L
G1033u14	WIAF-12491	M65188	GJA1, gap junction protein, alpha 1069 1, 43kD (connexin 43)	CCGCAATATAC [A/G] ACAAGCAAGC	M	A	G	N	D
G1033u15	WIAF-12492	M65188	GJA1, gap junction protein, alpha 1250 1, 43kD (connexin 43)	GTGGACCAGC [G/A] ACCTTCAAGC	M	G	A	R	Q
G1033u16	WIAF-12496	M65188	GJA1, gap junction protein, alpha 423 1, 43kD (connexin 43)	TATTGTGTC [T/C] GTACCCACAC	S	T	C	S	S
G1033u17	WIAF-12503	M65188	GJA1, gap junction protein, alpha 880 1, 43kD (connexin 43)	CGTTAAGGAT [C/T] GGGTTAAGGG	M	C	T	R	W
G1033u18	WIAF-12504	M65188	GJA1, gap junction protein, alpha 855 1, 43kD (connexin 43)	AACTCTCTTA [T/C] GTTTTCTTCA	S	T	C	Y	Y
G1033u19	WIAF-12505	M65188	GJA1, gap junction protein, alpha 576 1, 43kD (connexin 43)	AGTTCAAGTA [C/T] GGTATTGAAG	S	C	T	Y	Y
G1033u20	WIAF-12512	M65188	GJA1, gap junction protein, alpha 1255 1, 43kD (connexin 43)	CCAGGACCT [T/G] CAAGCAGAGC	M	T	G	S	A
G1033u21	WIAF-12513	M65188	GJA1, gap junction protein, alpha 1078 1, 43kD (connexin 43)	CAACAAGCAA [G/A] CAAGTGAGCA	M	G	A	A	T
G1033u22	WIAF-12514	M65188	GJA1, gap junction protein, alpha 1097 1, 43kD (connexin 43)	CAAAACTGGG [C/G] TAATTACAGT	M	C	G	A	G
G1034u1	WIAF-12443	J03544	PYGB, phosphorylase, glycogen; brain	AGACCTGTGC [A/G] TACACCAACC	S	A	G	A	A
G1034u2	WIAF-12469	J03544	PYGB, phosphorylase, glycogen; brain	GACACCCAG [T/C] GCCCGGTAC	M	T	C	V	A
G1034u3	WIAF-12470	J03544	PYGB, phosphorylase, glycogen; brain	TCCACTCGGA [G/C] ATCGTGAAC	M	G	C	E	D
G1034u4	WIAF-12471	J03544	PYGB, phosphorylase, glycogen; brain	GGGGCTGGCC [G/A] ATACCATCGT	M	G	A	D	N

G1034u5	WIAF-12472	J03544	1774	pyGB, phosphorylase, glycogen; brain	CCATGTTTGA [T/C] GTGCAATGTA	S	T	C	D	D
G1034u6	WIAF-12474	J03544	2449	pyGB, phosphorylase, glycogen; brain	AGGTGGACCA [G/A] CTGTACCGGA	S	G	A	Q	Q
G1034u7	WIAF-12508	J03544	718	pyGB, phosphorylase, glycogen; brain	CCCCCGACGG [C/T] GTGAAGTGGC	S	C	T	G	G
G1035u1	WIAF-12484	U97105	1962	DPYSL2, dihydropyrimidinase-like 2	GCAGAGGAGC [A/G] GCAGAGGATC	M	A	G	Q	R
G1035u2	WIAF-12485	U97105	2842	DPYSL2, dihydropyrimidinase-like 2	ATGACGGACC [T/C] GTGTGTGAAG	S	T	C	P	P
G1035u3	WIAF-12511	U97105	2062	DPYSL2, dihydropyrimidinase-like 2	CCATCACCAT [C/T] GCCAACCCAGA	S	C	T	I	I
G1036u1	WIAF-12444	D88460	311	WASL, Wiskott-Aldrich syndrome- like	ACGTGGGGTC [C/T] CTGTTGTCTCA	S	C	T	S	S
G1038u1	WIAF-12445	HT2746	994	PCTK2, PCTAIRE protein kinase 2	TAGAAGRAAG [G/A] TATTGCATCG	M	G	A	V	I
G1039u1	WIAF-12429	HT2747	955	serine/threonine kinase, PCTAIRE-3	ATCCAGAGT [C/T] GCATGTCTAGC	M	C	T	R	C
G1039u2	WIAF-12458	HT2747	808	serine/threonine kinase, PCTAIRE-3	CACAGAAGAG [A/T] CGTGGCCCGG	M	A	T	T	S
G1041u1	WIAF-12459	X72886	544	H.sapiens TYRO3 mRNA.	CAAGTGGCTG [G/C] CCTTGAGAG	M	G	C	A	P
G1041u2	WIAF-12460	X72886	693	H.sapiens TYRO3 mRNA.	TTGGCGGGAA [C/T] CGCCTGAAAC	S	C	T	N	N
G1041u3	WIAF-12502	X72886	561	H.sapiens TYRO3 mRNA.	AGAGCCTGGC [C/T] GACAACTGT	S	C	T	A	A
G1043u1	WIAF-12448	M94055	5481	Human voltage-gated sodium channel mRNA, complete cds.	CTCTGAGTGA [G/A] GATGACTTTG	S	G	A	E	E
G1043u2	WIAF-12449	M94055	5205	Human voltage-gated sodium channel mRNA, complete cds.	TTGAGACCTT [T/C] GGCAACAGCA	S	T	C	F	F
G1043u3	WIAF-12450	M94055	5224	Human voltage-gated sodium channel mRNA, complete cds.	CATGATCTGC [C/T] TGTTCCTCAAT	S	C	T	L	L
G1043u4	WIAF-12451	M94055	5514	Human voltage-gated sodium channel mRNA, complete cds.	AGGTTTGGGA [G/A] AAGTTTGATC	S	G	A	E	E
G1043u5	WIAF-12452	M94055	5217	Human voltage-gated sodium channel mRNA, complete cds.	GCAACAGCAT [G/C] ATCTGCCTGT	M	G	C	M	I
G1043u6	WIAF-12453	M94055	5334	Human voltage-gated sodium channel mRNA, complete cds.	GCTCACTTAA [A/G] GGAGACTGTG	S	A	G	K	K

G1043u7	WIAF-12454	M94055	Human voltage-gated sodium channel mRNA, complete cds.	5424	TGTACATCGC [G/C] GTCATCTCTGG	S	G	C	A	A
G1043u8	WIAF-12455	M94055	Human voltage-gated sodium channel mRNA, complete cds.	5322	ATCACCTGG [A/C] AGCTCAGTTA	S	A	C	G	G
G1043u9	WIAF-12456	M94055	Human voltage-gated sodium channel mRNA, complete cds.	1200	ATGGGTACAC [G/A] AGCTTTGACA	S	G	A	T	T
G1043u10	WIAF-12499	M94055	Human voltage-gated sodium channel mRNA, complete cds.	1170	TCTGTGTGAA [G/T] GCTGGTAGAA	M	G	T	K	N
G1046a1	WIAF-13187	U50352	ACCN1, amiloride-sensitive cation channel 1, neuronal (degenerin)	267	TCCAGCTGT [G/A] ACCCTCTGTA	S	G	A	V	V
G1046a2	WIAF-13188	U50352	ACCN1, amiloride-sensitive cation channel 1, neuronal (degenerin)	282	TCTGTAACCT [C/G] AATGGCTTCC	S	C	G	L	L
G1046a3	WIAF-13189	U50352	ACCN1, amiloride-sensitive cation channel 1, neuronal (degenerin)	315	TCACCACCAA [C/T] GACCTGTACC	S	C	t	N	N
G1046a4	WIAF-13190	U50352	ACCN1, amiloride-sensitive cation channel 1, neuronal (degenerin)	386	CCCCATCTGG [C/a] TGACCCCTCC	M	C	a	A	D
G1046a5	WIAF-13191	U50352	ACCN1, amiloride-sensitive cation channel 1, neuronal (degenerin)	417	CCCTCGGCA [G/A] AAGGCCAACT	S	G	A	Q	Q
G1048u1	WIAF-12641	HT5174S	REST, RE1-silencing transcription factor	3214	CAGTCAAAGC [G/A] GCTAAGGAG	S	G	A	A	A
G1048u2	WIAF-12642	HT5174S	REST, RE1-silencing transcription factor	3199	CAAGGAAGC [C/G] TTGGCAGTCA	S	C	G	A	A
G1048u3	WIAF-12657	HT5174S	REST, RE1-silencing transcription factor	2125	CTCCCATGGA [G/T] ACTGCTCAGA	M	G	T	E	D
G1048u4	WIAF-12660	HT5174S	REST, RE1-silencing transcription factor	2333	GGAACTGTT [A/C] AGATAGAGCT	M	A	C	K	Q
G1051u1	WIAF-12431	HT28321	SCNN1G, sodium channel, nonvoltage-gated 1, gamma	658	ATGACACCTC [C/T] GACTGTGCA	S	C	T	S	S
G1051u2	WIAF-12434	HT28321	SCNN1G, sodium channel, nonvoltage-gated 1, gamma	1735	AAGCAAAGGA [G/A] TGGTGGGCT	S	G	A	E	E
G1051u3	WIAF-12473	HT28321	SCNN1G, sodium channel, nonvoltage-gated 1, gamma	409	AGTCCCTGTA [T/C] GGCTTTCCAG	S	T	C	Y	Y

[illegible]

G1054u1	WIAF-12419	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	2252	TTGCAAGAG [C/T]TACAGGAGT	S	C	T	S	S
G1054u2	WIAF-12423	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4559	TGGTCATGTT [C/T]ATCTACTCCA	S	C	T	F	F
G1054u3	WIAF-12424	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4856	TCAACATGTA [C/G]ATCGCCATCA	N	C	G	Y	*
G1054u4	WIAF-12425	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4777	GTCAAGGGTG [A/G]CTCGGGCAAC	M	A	G	D	G
G1054u5	WIAF-12426	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4863	GTACATCGCC [A/G]TCATCCTGGA	M	A	G	I	V
G1054u6	WIAF-12427	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4566	GTTTCATCTAC [T/G]CCATCTTCGG	M	T	G	S	A
G1054u7	WIAF-12428	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4923	TGCTGAAGAT [G/T]ACTTTGAGAT	M	G	T	D	Y
G1054u8	WIAF-12446	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	3595	TTCTGGCTGA [T/C]CTTCAGCAFC	M	T	C	I	T
G1054u9	WIAF-12447	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4203	GGAGACAGAC [G/A]ACCAGAGCCA	M	G	A	D	N
G1054u10	WIAF-12495	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	4811	TCTGTTCTTT [C/A]TGCAGCTATA	M	C	A	F	L
G1054u11	WIAF-12497	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	5555	CAGGCAGAC [T/G]GTGCGCCAG	S	T	G	T	T
G1054u12	WIAF-12498	HT2202		SCN4A, sodium channel, voltage-gated, type IV, alpha polypeptide	5480	CAGGGACGC [C/T]GGACCCACTA	S	C	T	A	A
G1059u1	WIAF-12432	HT33704		APLP1, amyloid beta (A4) precursor-like protein 1	112	CGCTGTGCT [G/A]CCACTATTGC	S	G	A	L	L
G1059u2	WIAF-12433	HT33704		APLP1, amyloid beta (A4) precursor-like protein 1	140	TCTGCGCGC [C/T]AGCCCGCCAT	N	C	T	Q	*
G1059u3	WIAF-12435	HT33704		APLP1, amyloid beta (A4) precursor-like protein 1	1344	CAGCATGTGG [C/T]CGCCGTGGAT	M	C	T	A	V

G1059u4	WIAF-12457	HT33704	APLP1, amyloid beta (A4) precursor-like protein 1	1687	ATGAGCGAAA[G/A]GTGAATCGT	S	G	A	K	K
G1059u5	WIAF-12500	HT33704	APLP1, amyloid beta (A4) precursor-like protein 1	976	GGTTCTGTAG[A/G]GCCAAGATGG	S	A	G	R	R
G1059u6	WIAF-12501	HT33704	APLP1, amyloid beta (A4) precursor-like protein 1	1786	GTGAGGCTGT[A/G]TCGGGTCTGC	S	A	G	V	V
G1060u1	WIAF-12436	HT1418	APLP2, amyloid beta (A4) precursor-like protein 2	1744	CCAAGAAATT[C/G]AAGAGGAAAT	M	C	G	Q	E
G1060u2	WIAF-12467	HT1418	APLP2, amyloid beta (A4) precursor-like protein 2	2213	ATCAGGCTGG[T/G]GATGCTGAGG	M	T	G	V	G
G1060u3	WIAF-12468	HT1418	APLP2, amyloid beta (A4) precursor-like protein 2	2256	GCCACGGGAT[C/T]GTGGAGGTTG	S	C	T	I	I
G1066a1	WIAF-13195	HT3538	566 CCKBR, cholecystokinin B receptor	566	CTTTGGCACC[G/A]TCATCTGCAA	M	G	A	V	I
G1066a2	WIAF-13196	HT3538	607 CCKBR, cholecystokinin B receptor	607	GGGTGCTGT[G/A]AGTGTGTCCA	S	G	A	V	V
G1066a3	WIAF-13206	HT3538	864 CCKBR, cholecystokinin B receptor	864	CTGCTGCTTC[T/A]GCTCTGTTC	M	T	A	L	Q
G1067u1	WIAF-12478	HT0830	KCNA1, potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	684	AAACGCTGTG[C/T]ATCATCTGCT	S	C	T	C	C
G1067u2	WIAF-12479	HT0830	KCNA1, potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	722	GTGCGCTTCT[T/C]CGCCTGCCCC	M	T	C	F	S
G1067u3	WIAF-12480	HT0830	KCNA1, potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	804	ATTTTCATCAC[C/G]CTGGGCACCG	S	C	G	T	T
G1067u4	WIAF-12509	HT0830	KCNA1, potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)	690	TGTGCATCAT[C/T]TGCTTCTCCT	S	C	T	I	I
G1068u1	WIAF-12493	HT0831	KCNA2, potassium voltage-gated channel, shaker-related subfamily, member 2	774	TGAACATCAT[T/A]GACATTGTGG	S	T	A	I	I

G1070a1	WIAF-13197	HT27728		KCNJ6, potassium inwardly-rectifying channel, subfamily J, member 6	522		CACAGTACC [T/C] GGCTCTTTT	M	T	C	W	R
G1070a2	WIAF-13201	HT27728		KCNJ6, potassium inwardly-rectifying channel, subfamily J, member 6	1244		CCCTGGAGG [T/C] GGGTTCTACG	S	T	C	D	D
G1070a3	WIAF-13207	HT27728		KCNJ6, potassium inwardly-rectifying channel, subfamily J, member 6	707		ATAAATGCC [G/A] GAGGGAATTA	S	G	A	P	P
G1071u1	WIAF-12422	HT48672		KCNJ3, potassium inwardly-rectifying channel, subfamily J, member 3	1534		TTCCGGGCAA [C/T] TCAGAAGAAA	S	C	T	N	N
G1073u1	WIAF-12461	HT4556		KCNJ1, potassium inwardly-rectifying channel, subfamily J, member 1	1127		CACGTGTCCA [T/C] GTGCCCTTAT	M	T	C	M	T
G1074u1	WIAF-12462	HT27804		KCNAB2, potassium voltage-gated channel, shaker-related subfamily, beta member 2	289		ACCTCTTCGA [T/C] ACACAGAGAAG	S	T	C	D	D
G1079u1	WIAF-12463	HT27383		potassium channel, inwardly rectifying (GB:D50582)	1130		ACCTGGCCGA [T/A] GAGATCCTGT	M	T	A	D	E
G1079u2	WIAF-12464	HT27383		potassium channel, inwardly rectifying (GB:D50582)	1192		CGTTACTCTG [T/G] GGACTACTCC	M	T	G	V	G
G1079u3	WIAF-12481	HT27383		potassium channel, inwardly rectifying (GB:D50582)	708		GCTTGGCTGC [A/G] TCTTTCATGAA	M	A	G	I	V
G1079u4	WIAF-12482	HT27383		potassium channel, inwardly rectifying (GB:D50582)	779		CGGTGATGCG [T/C] CTGCGCCACG	S	T	C	A	A
G1079u5	WIAF-12483	HT27383		potassium channel, inwardly rectifying (GB:D50582)	276		GGACCCCTGCC [G/A] AGCCGAGGTA	M	G	A	E	K
G1079u6	WIAF-12510	HT27383		potassium channel, inwardly rectifying (GB:D50582)	489		GTGGCTCATC [G/A] CCTTCGCCCA	M	G	A	A	T
G1080u1	WIAF-12536	HT4412		KCNJ4, potassium inwardly-rectifying channel, subfamily J, member 4	1099		TGGACTACTC [A/G] CGTTTTCACA	S	A	G	S	S

G1080u2	WIAF-12537	HT4412	1050	KCNJ4, potassium inwardly-rectifying channel, subfamily J, member 4	GGCCACCGCT [T/A] TGAGCCTGTG	M	T	A	F	Y
G1081u1	WIAF-12538	HT27724	1090	KCNJ2, potassium inwardly-rectifying channel, subfamily J, member 2	GGCCACCGCT [A/T] TGAGCCTGTG	M	A	T	Y	F
G1082u1	WIAF-12662	HT28319	768	potassium channel, inwardly rectifying, high conductance, alpha subunit	CGCGGGTAC [C/T] GAGAGGGCG	S	C	T	T	T
G1082u2	WIAF-12663	HT28319	854	potassium channel, inwardly rectifying, high conductance, alpha subunit	CTGGTGTGC [C/T] CATCACCATC	M	C	T	P	L
G1082u3	WIAF-12679	HT28319	471	potassium channel, inwardly rectifying, high conductance, alpha subunit	TCTCCATCGA [G/C] ACAGAGACCA	M	G	C	E	D
G1084a1	WIAF-13198	HT0383	2028	KCNB1, potassium voltage-gated channel, Shab-related subfamily, member 1	CACGCCCCAG [C/A] AAGACTGGG	M	C	A	S	R
G1084a2	WIAF-13199	HT0383	2033	KCNB1, potassium voltage-gated channel, Shab-related subfamily, member 1	CCCAGCAAGA [C/G] TGGGGGACG	M	C	G	T	S
G1084a3	WIAF-13200	HT0383	2321	KCNB1, potassium voltage-gated channel, Shab-related subfamily, member 1	GAGTGTGCCA [C/A] GCTTTTGAC	M	C	A	T	K
G1084a4	WIAF-13208	HT0383	870	KCNB1, potassium voltage-gated channel, Shab-related subfamily, member 1	ACAACCCCA [G/A] CTGGCCACG	S	G	A	Q	Q
G1088u1	WIAF-12516	HT0522	1503	KCNA5, potassium voltage-gated channel, shaker-related subfamily, member 5	TCCTGGGCAA [G/A] ACCTTGACG	S	G	A	K	K
G1088u2	WIAF-12519	HT0522	1249	KCNA5, potassium voltage-gated channel, shaker-related subfamily, member 5	CGAGCTGCTC [G/A] TGCGCTTCTT	M	G	A	V	M

G1088u3	WIAF-12520	HT0522		KCN5, potassium voltage-gated channel, shaker-related subfamily, member 5	973	CTCTGGGTCC [G/A] CGCGGGCCAT	M	G	A	A	T
G1088u4	WIAF-12521	HT0522		KCN5, potassium voltage-gated channel, shaker-related subfamily, member 5	1013	GTATCTCTCA [T/C] CTCCATCATC	M	T	C	I	T
G1090u1	WIAF-12651	HT1497		KCN6, potassium voltage-gated channel, shaker-related subfamily, member 6	1836	CAACCAGCCA [G/A] TGGAGGAGGC	M	G	A	S	N
G1091u1	WIAF-12714	HT0222		KCN3, potassium voltage-gated channel, shaker-related subfamily, member 3	843	CATCATCTGG [T/C] TCTCCTTCCA	M	T	C	F	L
G1094a1	WIAF-13218	HT27381		KCNJ8, potassium inwardly-rectifying channel, subfamily J, member 8	1280	GTCTATTCTG [T/a] GGATTACTCC	M	T	a	V	E
G1095u1	WIAF-12532	HT2629		KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member 1	765	TTCTCTACTT [C/T] GGCTTGGGT	S	C	T	F	F
G1095u2	WIAF-12533	HT2629		KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member 1	2441	GTGCTCTGCA [T/C] CTTTGGCGAC	M	T	C	I	T
G1095u3	WIAF-12534	HT2629		KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member 1	2714	GATGATACTT [C/G] GCTGCAGGAC	M	C	G	S	W
G1095u4	WIAF-12535	HT2629		KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member 1	2439	TCGTGGTCTG [C/T] ATCTTTGGCG	S	C	T	C	C

002060-24425960

G1095u5	WIAF-12539	HT2629	3048	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	CACTCATGAG [C/T] GCGACGFACT	S	C	T	S	S
G1095u6	WIAF-12544	HT2629	2352	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	GGATGTTTCA [C/T] TGGTGTGCAC	S	C	T	H	H
G1095u7	WIAF-12545	HT2629	2392	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	CATCTTGACT [C/T] GAAGTGAAGC	N	C	T	R	*
G1095u8	WIAF-12546	HT2629	2295	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	CTGGCAATGA [T/C] CAGATTGACA	S	T	C	D	D
G1095u9	WIAF-12548	HT2629	2949	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	AGTTTTTTGGA [C/T] CAAGACGATG	S	C	T	D	D
G1095u10	WIAF-12549	HT2629	2865	KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member	TGCACGGGAT [G/A] TTAGGTCAAC	M	G	A	M	I
G1096u1	WIAF-12547	L26318	930	PRKM8, protein kinase mitogen-activated 8 (MAP kinase)	TGCTGGTAAAT [A/T] GATGCACTCTA	S	A	T	I	I
G1098u1	WIAF-12515	L19711	2650	DAG1, dystroglycan 1 (dystrophin-associated glycoprotein 1)	TCTACCTGCA [C/T] ACAGTCAATTC	S	C	T	H	H
G110u1	WIAF-10385	HT27392	230	meiosis-specific recA homolog, Hslim15	CAAAGGTATA [C/T] AGATGACAAC	N	C	T	Q	*
G110u2	WIAF-10397	HT27392	1050	meiosis-specific recA homolog, Hslim15	CCTGAAAATG [A/G] AGCCACCTTC	M	A	G	E	G
G110u3	WIAF-10399	HT27392	674	meiosis-specific recA homolog, Hslim15	TGAACATCAG [A/G] TGGAGCTACT	M	A	G	M	V

G1106u1	WIAF-12647	HT5073	5781	MAP1B, microtubule-associated protein 1B	ACTATGAGAA [G/A] ATAGAGAGAA	S	G	A	K	K
G1106u2	WIAF-12648	HT5073	5916	MAP1B, microtubule-associated protein 1B	CTGAAGAGGG [C/T] GGGTACTCAT	S	C	T	G	G
G1106u3	WIAF-12650	HT5073	1837	MAP1B, microtubule-associated protein 1B	AGACAAGCCA [G/A] TAAAAACAGA	M	G	A	V	I
G1106u4	WIAF-12653	HT5073	2476	MAP1B, microtubule-associated protein 1B	CACCACAGCA [G/A] CTGTCTATGGC	M	G	A	A	T
G1106u5	WIAF-12656	HT5073	3913	MAP1B, microtubule-associated protein 1B	GCCCAATGAG [A/G] TTAAGTCTC	M	A	G	I	V
G1106u6	WIAF-12667	HT5073	559	MAP1B, microtubule-associated protein 1B	GATTTTCACC [G/A] ATCAAGAGAT	M	G	A	D	N
G1106u7	WIAF-12668	HT5073	570	MAP1B, microtubule-associated protein 1B	ATCAAGAGAT [C/T] GGGGAGTTAC	S	C	T	I	I
G1106u8	WIAF-12669	HT5073	6175	MAP1B, microtubule-associated protein 1B	TACTTCCACA [T/C] ACTGTTACGA	M	T	C	Y	H
G1106u9	WIAF-12670	HT5073	1215	MAP1B, microtubule-associated protein 1B	TCACTCTCCA [G/C] TACCTAAACA	M	G	C	Q	H
G1106u10	WIAF-12672	HT5073	1821	MAP1B, microtubule-associated protein 1B	AGGTAATGGT [G/A] AAAAAAGACA	S	G	A	V	V
G1106u11	WIAF-12673	HT5073	2727	MAP1B, microtubule-associated protein 1B	GTCTGCGCA [G/T] TCCCTGTATG	M	G	T	E	D
G1106u12	WIAF-12674	HT5073	2739	MAP1B, microtubule-associated protein 1B	CCCCTGATGA [G/A] GGAATCACTA	S	G	A	E	E
G1106u13	WIAF-12676	HT5073	3643	MAP1B, microtubule-associated protein 1B	AGATGCCACT [G/A] ATGGCAAGGA	M	G	A	D	N
G1106u14	WIAF-12677	HT5073	3609	MAP1B, microtubule-associated protein 1B	CACCGCTCAA [C/T] GGATTTTCTG	S	C	T	N	N
G1106u15	WIAF-12682	HT5073	4752	MAP1B, microtubule-associated protein 1B	TTCCAGAGCC [A/T] ACACAGATG	S	A	T	P	P
G1110u1	WIAF-12517	HT1096	1527	myelin associated glycoprotein	GCGSCCTCGT [G/C] CTCACCAGCA	S	G	C	V	V
G1110u2	WIAF-12518	HT1096	1678	myelin associated glycoprotein	TGTGGGCGCC [G/T] TGGTGGCCTT	M	G	T	V	L
G1110u3	WIAF-12522	HT1096	1271	myelin associated glycoprotein	GCCGTGTAC [C/T] CGAGGATGAT	M	C	T	P	L
G1113u1	WIAF-12523	HT2242	353	myelin transcription factor 1	AATTCCGATC [G/T] GATCCTCAGG	M	G	T	R	L
G1116a1	WIAF-13217	HT28451	417	myelin oligodendrocyte glycoprotein (MOG)	CAAGTTATC [G/A] AGACCCCTCTC	S	G	A	S	S
G1116a2	WIAF-13219	HT28451	913	myelin oligodendrocyte glycoprotein (MOG)	GCATATCACT [C/G] TTGGCCTCGT	M	C	G	L	V

004060" 2444560

G1116a3	WIAP-13220	HT28451	myelin oligodendrocyte 922 glycoprotein (MOG)	TCTTGGCTC [G/A]TCTTCTCTG	M	G	A	V	I
G1120u1	WIAP-12525	HT3695	1200 neurofilament, subunit H	TAGAGATAGC [T/C]GCTTACAGAA	S	T	C	A	A
G1123u1	WIAP-12542	HT2569	OMG, oligodendrocyte myelin glycoprotein	CAGCTGCAAC [T/C]CTAACTATTC	S	T	C	T	T
G1126u1	WIAP-12526	HT28354	PSEN2, presenilin 2 (Alzheimer disease 4)	GAGCGAAGCA [T/C]GTGATCATGC	S	T	C	H	H
G1126u2	WIAP-12527	HT28354	PSEN2, presenilin 2 (Alzheimer disease 4)	ATGGAGAGAA [T/C]ACTGCCCACT	S	T	C	N	N
G1126u3	WIAP-12528	HT28354	PSEN2, presenilin 2 (Alzheimer disease 4)	TAATGTCGGC [C/T]GAGAGCCCCA	S	C	T	A	A
G1126u4	WIAP-12543	HT28354	PSEN2, presenilin 2 (Alzheimer disease 4)	GACCTGACC [G/A]CTATGTCTGT	M	G	A	R	H
G117u1	WIAP-10391	HT27765	GTPB, G/T mismatch-binding protein	ACTTCTCACC [A/G]GGAGATTGG	S	A	G	P	P
G117u2	WIAP-10392	HT27765	GTPB, G/T mismatch-binding protein	AACGTGCAGA [T/C]GAAGCCTTAA	S	T	C	D	D
G117u3	WIAP-10407	HT27765	GTPB, G/T mismatch-binding protein	CCCACGTTAG [T/C]GGAGGTGGTG	S	T	C	S	S
G117u4	WIAP-10411	HT27765	GTPB, G/T mismatch-binding protein	CATTGTTCGA [G/A]ATTAGGACT	M	G	A	R	K
G117u5	WIAP-10412	HT27765	GTPB, G/T mismatch-binding protein	GACAGCAGGG [C/T]TATAATGTAT	M	C	T	A	V
G117u6	WIAP-10413	HT27765	GTPB, G/T mismatch-binding protein	AAGAGTCAGA [A/T]CCACCCAGAC	M	A	T	N	I
G125u1	WIAP-10371	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	CAGTAATTTT [C/T]CTCATCTTGT	M	C	T	P	S
G125u2	WIAP-10372	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	TAATGATGA [C/A]ATTGCAGATA	M	C	A	D	E
G125u3	WIAP-10373	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	CAATGGAAGA [T/G]GTTCTTGAA	M	T	G	D	E
G125u5	WIAP-10375	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	CACTTATACC [C/T]CTTGTGTATG	S	C	T	P	P

G125u6	WIAF-10383	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	8713	ATCCTTGGAT [C/T] CAGCTATTGG	M	C	T	P	S
G125u7	WIAF-10396	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	1825	GACTTTGGCA [C/G] TGACCACCAG	M	C	G	L	V
G125u8	WIAF-10398	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	2924	ACTACTGCTC [A/G] GACCAATACT	M	A	G	Q	R
G125u9	WIAF-10405	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	8967	TTGAAGGTGT [C/T] TTCAGAAGAT	S	C	T	V	V
G125u10	WIAF-10408	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	6954	CCAAACACCT [T/C] GTAGAACTCT	S	T	C	L	L
G125u11	WIAF-10409	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	6855	TTCAGGAGCC [T/C] ATCATGGCTC	S	T	C	P	P
G125u12	WIAF-10410	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	6801	TATATATTAA [G/T] TGGCAGAAAC	M	G	T	K	N
G125u13	WIAF-10421	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	335	CATTCAGATT [C/G] CAAACAAGGA	M	C	G	S	C
G125u14	WIAF-11607	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	3966	TTCCACATCT [G/A] GTGATTAGAA	S	G	A	L	L
G125a15	WIAF-13130	HT28632	ATM, ataxia telangiectasia mutated (includes complementation groups A, C and D)	8642	GAGAAATATG [A/C] AGTCCTCATG	M	A	C	E	A

004060" 2442566

G136u1	WIAF-10388	HT3337	535 2)	MLH1, mutL (E. coli) homolog 1 (colon cancer, nonpolyposis type	AGGAGAAAAG [C/T] TTTAAAAAT	M C T A V
G136u2	WIAF-10389	HT3337	769 2)	MLH1, mutL (E. coli) homolog 1 (colon cancer, nonpolyposis type	TTCAAAATGA [A/G] TGGTTACATA	M A G N S
G144u1	WIAF-11638	HT3625	1129	FOS, v-fos FBJ murine osteosarcoma viral oncogene homolog	CCTGTGCACT [C/T] CGTGTGTAC	M C T P S
G1461u1	WIAF-12562	HT0329	684	pRB-binding protein	TTGCCAAGAA [G/A] TCCAAGAACC	S G A K K
G1466u1	WIAF-12571	HT27849	2128	API2, apoptosis inhibitor 2	ATGATCCATG [G/C] GTAGAACATG	M G C W C
G1468u1	WIAF-12563	HT4986	1928	apoptosis inhibitor, neuronal	CCACGAGACC [A/T] GACGAGGGGC	S A T P P
G1468u2	WIAF-12564	HT4986	3057	apoptosis inhibitor, neuronal	TTTGCAATTC [C/G] TTCAAGGGAG	M C G L V
G1472u1	WIAF-12565	HT28478	242	BAK1, BCL2-antagonist/killer 1	GGCAGGAGTG [C/T] GGAGAGCCTG	S C T C C
G1472u2	WIAF-12572	HT28478	509	BAK1, BCL2-antagonist/killer 1	TGCAGCCAC [G/A] GCAGAGAATG	S G A T T
G1473u1	WIAF-12568	HT28606	394	CASP6, caspase 6, apoptosis- related cysteine protease	GGTGTCAACT [G/C] TTAGCCACGC	M G C V L
G1473u2	WIAF-12576	HT28606	411	CASP6, caspase 6, apoptosis- related cysteine protease	ACGAGATGC [C/T] GATTGCTTTG	S C T A A
G1479u1	WIAF-12550	Y09077	711	ATR, ataxia telangiectasia and Rad3 related	ACTTTATTAA [T/C] GGTTCCTTACT	M T C M T
G1479u2	WIAF-12551	Y09077	4303	ATR, ataxia telangiectasia and Rad3 related	TTGGGTATGC [T/C] GATAATAGCC	S T C A A
G1479u3	WIAF-12552	Y09077	1894	ATR, ataxia telangiectasia and Rad3 related	ATTCTGATGA [T/C] GGCTGTTAA	S T C D D
G1479u4	WIAF-12553	Y09077	1855	ATR, ataxia telangiectasia and Rad3 related	ATTTATGTGG [T/A] ATGCTCTCAC	S T A G G
G1479u5	WIAF-12558	Y09077	5287	ATR, ataxia telangiectasia and Rad3 related	TCATTCATTA [T/C] CATGGTGTAG	S T C Y Y

000000242560

004000-2445960

G1479u6	WIAF-12559	Y09077		ATR, ataxia telangiectasia and 5539 Rad3 related	CAGCTTTTAA[T/C]GACTCACTGA	S	T	C	Y	Y
G1479u7	WIAF-12569	Y09077		ATR, ataxia telangiectasia and 1540 Rad3 related	ATCCTGTTAT[T/C]GAGATGTTAG	S	T	C	I	I
G1479u8	WIAF-12570	Y09077		ATR, ataxia telangiectasia and 2521 Rad3 related	ATTTAATGGA[A/G]GATCCAGACA	S	A	G	E	E
G1482u1	WIAF-12560	HT27870		3176 BLM, Bloom syndrome	AAAATATAAC[G/A]GAATGCAGGA	S	G	A	T	T
G1482u2	WIAF-12561	HT27870		3605 BLM, Bloom syndrome	GAAATAAGC[C/A]CAAACTGTAC	S	C	A	A	A
G1482u3	WIAF-12573	HT27870		2677 BLM, Bloom syndrome	TATGTATTAC[C/T]GAAAAAGCCT	M	C	T	P	L
G1483u1	WIAF-12597	HT1470		MYBL2, v-myb avian myeloblastosis 1910 viral oncogene homolog-like 2	GGATGAGGAT[G/A]TGAAGCTGAT	M	G	A	V	M
G1483u2	WIAF-12610	HT1470		MYBL2, v-myb avian myeloblastosis 244 viral oncogene homolog-like 2	ATGAGGAGGA[C/T]GAGCAGCTGA	S	C	T	D	D
G1483u3	WIAF-12611	HT1470		MYBL2, v-myb avian myeloblastosis 1406 viral oncogene homolog-like 2	CACTGAGAAT[A/G]GCACAGTCT	M	A	G	S	G
G1485u1	WIAF-12581	HT1432		1941 BCR, breakpoint cluster region	TGGAGATGAG[A/G]AAATGGGTCC	S	A	G	R	R
G1485u2	WIAF-12582	HT1432		3144 BCR, breakpoint cluster region	TGACCATCAA[T/C]AAGGAAGATG	S	T	C	N	N
G1485u3	WIAF-12583	HT1432		3777 BCR, breakpoint cluster region	ATAACAAGGA[T/C]GTGTGGTGA	S	T	C	D	D
G1485u4	WIAF-12603	HT1432		2831 BCR, breakpoint cluster region	CAGATCAAGA[G/A]TGACATCCAG	M	G	A	S	N
G1485u5	WIAF-12608	HT1432		4217 BCR, breakpoint cluster region	ATCCCTGCCC[C/T]GGACAGCAAG	M	C	T	P	L
G1486u1	WIAF-12578	HT33770		BRCA2, breast cancer 2, early 1909 onset	ATTGATAATG[G/A]AAGCTGGCCA	M	G	A	G	E
G1486u2	WIAF-12579	HT33770		BRCA2, breast cancer 2, early 3623 onset	AGTTTAGAAA[A/G]CCAAGCTACA	S	A	G	K	K
G1486u3	WIAF-12586	HT33770		BRCA2, breast cancer 2, early 1341 onset	AAATGTAGCA[A/C]ATCAGAAGCC	M	A	C	N	H
G1486u4	WIAF-12594	HT33770		BRCA2, breast cancer 2, early 446 onset	CTTATAATCA[G/A]CTGGCTTCAA	S	G	A	Q	Q
G1486u5	WIAF-12598	HT33770		BRCA2, breast cancer 2, early 3013 onset	ACCATGGTTT[T/C]ATATGGAGAC	M	T	C	L	S

G1486u6	WIAF-12599	HT33770	BRCA2, breast cancer 2, early onset	3187	GAAAAAATA[A/T]TGATTACATG	M	A	T	N	I
G1486u7	WIAF-12604	HT33770	BRCA2, breast cancer 2, early onset	4971	AGCATGTGAG[A/C]CCATTGAGAT	M	A	C	T	P
G1486u8	WIAF-12607	HT33770	BRCA2, breast cancer 2, early onset	4034	ATGATTCTGT[C/T]GTTCAATGT	S	C	T	V	V
G1487u1	WIAF-12584	HT27632	BRCA1, breast cancer 1, early onset	2536	AGTCAGTGTG[C/G]AGCATTGAA	M	C	G	A	G
G1487u2	WIAF-12587	HT27632	BRCA1, breast cancer 1, early onset	4697	CATCTCAAGA[G/C]GAGCTCATTA	M	G	C	E	D
G1487u3	WIAF-12595	HT27632	BRCA1, breast cancer 1, early onset	469	TCTCCTGAAC[A/G]TCTAAAAGAT	M	A	G	H	R
G1487u4	WIAF-12600	HT27632	BRCA1, breast cancer 1, early onset	3667	AGCGTCCAGA[A/G]AGGAGAGCTT	M	A	G	K	R
G1487u5	WIAF-12601	HT27632	BRCA1, breast cancer 1, early onset	3537	TATGGGAAGT[A/G]GTCAATGCATC	M	A	G	S	G
G1487u6	WIAF-12602	HT27632	BRCA1, breast cancer 1, early onset	4956	ATCTGCCAG[A/G]GTCCAGCTGC	M	A	G	S	G
G1487u7	WIAF-12605	HT27632	BRCA1, breast cancer 1, early onset	2090	AGTACAAACCA[A/G]ATGCCAGTCA	S	A	G	Q	Q
G1487u8	WIAF-12614	HT27632	BRCA1, breast cancer 1, early onset	233	TCTCCACAAA[G/A]TGTACCACAA	S	G	A	K	K
G1492u1	WIAF-12585	HT3506	3912 cell death-associated kinase		TCCAGGTCCG[T/C]GGCCTGGAGA	S	T	C	R	R
G1492u2	WIAF-12593	HT3506	4352 cell death-associated kinase		TACAACACCA[A/G]TAACGGGGCT	M	A	G	N	S
G1492u3	WIAF-12606	HT3506	2127 cell death-associated kinase		GCAATTTTGA[C/T]ATCTCCAACA	S	C	T	D	D
G1492u4	WIAF-12612	HT3506	1605 cell death-associated kinase		TGAAATTTCT[C/T]AGTGAGAAACA	S	C	T	L	L
G1494u1	WIAF-12589	HT28507	366 cell death-inducing protein Bik		TTTCTTCCCT[G/C]ATCCTGATCT	S	G	C	L	L
G1495u1	WIAF-12580	HT27803	CSE1L, chromosome segregation 1 (yeast homolog)-like	759	CAGCAATGAC[A/C]TTCCCATCGC	M	A	C	I	L
G1501u1	WIAF-13502	HT1949	MCC, mutated in colorectal cancers	1181	CAGCTGAGAA[C/T]GCTGCCAAGG	S	C	T	N	N
G1501u2	WIAF-13503	HT1949	MCC, mutated in colorectal cancers	1753	TGTCCCTAGC[T/C]GAACTCAGGA	S	T	C	A	A
G1501u3	WIAF-13504	HT1949	MCC, mutated in colorectal cancers	2344	AGCGAACGAC[G/A]CTTCGCTATG	S	G	A	T	T
G1501u4	WIAF-13521	HT1949	MCC, mutated in colorectal cancers	445						

G1501u5	WIAF-13522	HT1949	1504	MCC, mutated in colorectal cancers	AAAGCAATGC [T/C] GAGAGGATGA	S	T	C	A	A
G1501u6	WIAF-13527	HT1949	2511	MCC, mutated in colorectal cancers	TTCTGTAATG [A/G] TCTAAAGCGG	M	A	G	D	G
G1502u1	WIAF-12633	HT1547	870	CCND1, cyclin D1 (PRAD1: parathyroid adenomatosis 1)	AGTGTGACCC [A/G] GACTGCGCTCC	S	A	G	P	P
G1503u1	WIAF-13741	U37022	1151	CDK4, cyclin-dependent kinase 4	CATGCCAATT [G/A] CATCGTTTAC	M	G	A	C	Y
G1503u2	WIAF-13742	U37022	1410	CDK4, cyclin-dependent kinase 4	CTGAAGCCGA [C/T] CAGTTGGGCA	S	C	T	D	D
G1503u3	WIAF-13743	U37022	1328	CDK4, cyclin-dependent kinase 4	TATGCAACAC [C/T] TGTGGACATG	M	C	T	P	L
G1503u4	WIAF-13780	U37022	1194	CDK4, cyclin-dependent kinase 4	TTCTGGTGCAC [A/G] AGTGGTGGAA	S	A	G	T	T
G1503u5	WIAF-13781	U37022	1443	CDK4, cyclin-dependent kinase 4	TGATTGGGCT [G/A] CCTCCAGAGG	S	G	A	L	L
G1503u6	WIAF-13787	U37022	1633	CDK4, cyclin-dependent kinase 4	CTCTTATCTA [C/T] ATAAGGATGA	M	C	T	H	Y
G1517u1	WIAF-12618	HT1132	3894	ERBB3, v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 3	CAGACCTCAG [T/C] GCCTCTCTGG	S	T	C	S	S
G152u1	WIAF-11608	HT3854	1673	HSPALL, heat shock 70kD protein-like 1	GTGAGTGATG [A/C] AGGTTTGAAG	M	A	C	E	A
G152u2	WIAF-11629	HT3854	1683	HSPALL, heat shock 70kD protein-like 1	AAGGTTTGAA [G/A] GGCAAGATTA	S	G	A	K	K
G152u3	WIAF-11609	HT3854	1478	HSPALL, heat shock 70kD protein-like 1	GTCAACAGCCA [C/T] GGACAAGAGC	M	C	T	T	M
G152u4	WIAF-11610	HT3854	1443	HSPALL, heat shock 70kD protein-like 1	TGACGTTTGA [C/T] ATTGATGCCA	S	C	T	D	D
G1520u1	WIAF-12162	HT1175	2211	DNA excision repair protein ERCC2, 5' end	TGACCCGTGGA [C/T] GAGGGTGTCC	S	C	T	D	D
G1520u2	WIAF-12166	HT1175	546	DNA excision repair protein ERCC2, 5' end	CCCACTGCCG [A/C] TTTCTATGAGG	S	A	C	R	R
G1527u1	WIAF-12168	HT0086	577	GSTM2, glutathione S-transferase M2 (muscle)	TCATCTCCCG [A/C] TTTGAGGGCT	S	A	C	R	R
G1527u2	WIAF-12169	HT0086	644	GSTM2, glutathione S-transferase M2 (muscle)	ACCTGTGTTT [A/T] CAARGATGGC	M	A	T	T	S
G1527u3	WIAF-12171	HT0086	100	GSTM2, glutathione S-transferase M2 (muscle)	ACTCAAGCTA [C/T] GAGGAAAAGA	S	C	T	Y	Y
G1527u4	WIAF-12172	HT0086	41	GSTM2, glutathione S-transferase M2 (muscle)	GGGTACTTGG [A/G] ACATCCGCGG	M	A	G	N	D

G1527u5	WIAF-12173	HT0086	215 M2	glutathione S-transferase	GSTM2, glutathione S-transferase	215 M2 (muscle)	GATTGATGGG [A/G] CTCACAAGAT	M	A	G	T	A
G1527u6	WIAF-12194	HT0086	238 M2	glutathione S-transferase	GSTM2, glutathione S-transferase	238 M2 (muscle)	CCCAGAGCAA [T/C] GCCATCCTGC	S	T	C	N	N
G1528u1	WIAF-11950	HT1811	529 M3	glutathione S-transferase	GSTM3, glutathione S-transferase	529 M3 (brain)	GTATATTGA [C/G] CCCAAGTGCC	M	C	G	D	E
G1528u2	WIAF-11951	HT1811	674 M3	glutathione S-transferase	GSTM3, glutathione S-transferase	674 M3 (brain)	CAACAAGCCT [G/A] TATGCTGAGC	M	G	A	V	I
G1528u3	WIAF-11989	HT1811	572 M3	glutathione S-transferase	GSTM3, glutathione S-transferase	572 M3 (brain)	GGCTTTCAAG [T/G] GCCGTTTGA	M	T	G	C	G
G1528u4	WIAF-13470	HT1811	240 M3	glutathione S-transferase	GSTM3, glutathione S-transferase	240 M3 (brain)	CAGAGCAATG [C/A] CATCTTGGC	M	C	A	A	D
G1529u1	WIAF-14146	HT2006	797 M4	glutathione S-transferase	GSTM4, glutathione S-transferase	797 M4	TGGACGCCTT [C/T] CCAAATCTGA	S	C	T	F	F
G153u1	WIAF-12163	HT3856	1212	HSPA1B, heat shock 70kD protein 1	GSTM5, glutathione S-transferase	1212 HSPA1B, heat shock 70kD protein 1	TGGGGCTGGA [G/A] ACGCCGGAG	S	G	A	E	E
G153u2	WIAF-12182	HT3856	676	HSPA1B, heat shock 70kD protein 1	GSTM5, glutathione S-transferase	676 HSPA1B, heat shock 70kD protein 1	GGCCGGGGAC [A/G] CCCACCTGGG	M	A	G	T	A
G153u3	WIAF-12183	HT3856	1695	HSPA1B, heat shock 70kD protein 1	GSTM5, glutathione S-transferase	1695 HSPA1B, heat shock 70kD protein 1	TCAGCGAGGC [C/G] GACAAGAAGA	S	C	G	A	A
G153u4	WIAF-12189	HT3856	330	HSPA1B, heat shock 70kD protein 1	GSTM5, glutathione S-transferase	330 HSPA1B, heat shock 70kD protein 1	ACAAGGGGA [G/C] ACCAAGGCAT	M	G	C	E	D
G153u5	WIAF-12190	HT3856	1053	HSPA1B, heat shock 70kD protein 1	GSTM5, glutathione S-transferase	1053 HSPA1B, heat shock 70kD protein 1	AGCTGCTGCA [A/G] GACTTCTTCA	S	A	G	Q	Q
G1530u1	WIAF-11964	HT3010	673 M5	glutathione S-transferase	GSTM5, glutathione S-transferase	673 M5	ATTCTCCGA [G/A] GTCTTTTGT	M	G	A	G	S
G1530u2	WIAF-11995	HT3010	593 M5	glutathione S-transferase	GSTM5, glutathione S-transferase	593 M5	GACGCTTCC [T/C] AAACCTGAAG	M	T	C	L	P
G1530u3	WIAF-13473	HT3010	693 M5	glutathione S-transferase	GSTM5, glutathione S-transferase	693 M5	TTGGAAGTC [A/G] GCTACATGGA	S	A	G	S	S
G1533u1	WIAF-13458	HT27460	543	glutathione S-transferase	GSTM5, glutathione S-transferase	543	CTCTCGGCTA [C/T] GAACCTGTTG	S	C	T	Y	Y
G1533u2	WIAF-13460	HT27460	417	glutathione S-transferase	GSTM5, glutathione S-transferase	417	GGACTGCCAT [G/A] GACCAGGCC	M	G	A	M	I
G1533u3	WIAF-13461	HT27460	359	glutathione S-transferase	GSTM5, glutathione S-transferase	359	CAGGTGTGG [G/A] GCCACTCAT	M	G	A	G	E
G1533u4	WIAF-13462	HT27460	363	glutathione S-transferase	GSTM5, glutathione S-transferase	363	TGTTGGGGCC [A/C] CTCATTGGGG	S	A	C	P	P
G1533u5	WIAF-13463	HT27460	385	glutathione S-transferase	GSTM5, glutathione S-transferase	385	CCAGGTGCC [G/A] AGGAGAAGGT	M	G	A	E	K
G1535u1	WIAF-11952	HT0436	517	HCK, hemopoietic cell kinase	GSTM5, glutathione S-transferase	517	CCGCGTTGAC [T/C] CTCTGGAGAC	M	T	C	S	P

G1535u2	WIAF-12013	HT0436	793 HCK, hemopoietic cell kinase	TGGACCACTA [C/T] AAGAAGGGA	S	C	T	Y	Y
G1535u3	WIAF-13464	HT0436	357 HCK, hemopoietic cell kinase	TCATCGTGGT [T/C] GCCTGTATG	S	T	C	V	V
G1535u4	WIAF-13465	HT0436	387 HCK, hemopoietic cell kinase	CCATTACCA [C/T] GAAGACTCA	S	C	T	H	H
G1535u5	WIAF-13466	HT0436	471 HCK, hemopoietic cell kinase	CCCTGGCCAC [C/G] CGGAAGGAGG	S	C	G	T	T
G1535u6	WIAF-13467	HT0436	240 HCK, hemopoietic cell kinase	CCAGGCCAG [C/T] CCACACTGTC	S	C	T	S	S
G1535u7	WIAF-13468	HT0436	394 HCK, hemopoietic cell kinase	CCACGAAGAC [C/T] TCAGCTTCCA	M	C	T	L	F
G1537u1	WIAF-12020	U04045	MSH2, mutS (E. coli) homolog 2 (colon cancer, nonpolyposis type 1514 1)	GTGAATTAAAG [A/G] GAAATAATGA	S	A	G	R	R
G1537u2	WIAF-12044	U04045	MSH2, mutS (E. coli) homolog 2 (colon cancer, nonpolyposis type 599 1)	GACTGTGTGA [A/T] TTCCTGTATA	M	A	T	E	D
G1537u3	WIAF-12045	U04045	MSH2, mutS (E. coli) homolog 2 (colon cancer, nonpolyposis type 1452 1)	AGATATGGAT [C/T] AGTGGAAAA	N	C	T	Q	*
G1537u4	WIAF-12076	U04045	MSH2, mutS (E. coli) homolog 2 (colon cancer, nonpolyposis type 938 1)	GACAGTTTGA [A/T] CTGACTACTT	M	A	T	E	D
G1537u5	WIAF-12077	U04045	MSH2, mutS (E. coli) homolog 2 (colon cancer, nonpolyposis type 1878 1)	TCAGCTAGAT [G/A] CTGTTGTGAG	M	G	A	A	T
G1543u1	WIAF-13856	J00119	MOS, v-mos Moloney murine sarcoma viral oncogene homolog	GAGTTTCTGG [G/T] CTGAGCTCAA	M	G	T	A	S
G1543u2	WIAF-13857	J00119	MOS, v-mos Moloney murine sarcoma viral oncogene homolog	GCACGGCCAC [G/A] CCCGCAGGGT	S	G	A	T	T
G1544u1	WIAF-12018	U59464	PTCH, patched (Drosophila) homolog	CATCCCGAAT [C/T] CAGGCATCAC	M	C	T	S	F
G1544u2	WIAF-12019	U59464	PTCH, patched (Drosophila) homolog	GCGTGGTCCG [C/T] TTCGCCATGC	S	C	T	R	R
G1544u3	WIAF-12027	U59464	PTCH, patched (Drosophila) homolog	ATTTTGCCAT [G/T] GTTCGTGCTCA	M	G	T	M	I
G1544u4	WIAF-12029	U59464	PTCH, patched (Drosophila) homolog	CTGCCATGGG [C/T] AGTCCGTGTC	S	C	T	G	G

G1544u5	WIAF-12043	U59464	3845	PTCH, patched (Drosophila) homolog	CCTCGAAC[C/T]GAGACAGCAG	M	C	T	P	L
G1544u6	WIAF-12056	U59464	1433	PTCH, patched (Drosophila) homolog	CTGCTGTTG[C/T]ACTGTCAGTG	M	C	T	A	V
G1544u7	WIAF-12058	U59464	3298	PTCH, patched (Drosophila) homolog	CACCGTTAC[G/C]TTGCTTTGGC	M	G	C	V	L
G1544u8	WIAF-12062	U59464	3986	PTCH, patched (Drosophila) homolog	TCTACTGAAG[G/A]GCATTCTGGC	M	G	A	G	E
G1544u9	WIAF-13489	U59464	1665	PTCH, patched (Drosophila) homolog	CCATCAGCAA[T/C]GTCACAGCCT	S	T	C	N	N
G1544u10	WIAF-13490	U59464	2396	PTCH, patched (Drosophila) homolog	AAATACTTTT[C/T]TTTCTACAAC	M	C	T	S	F
G1544u11	WIAF-13491	U59464	2199	PTCH, patched (Drosophila) homolog	GGACACTCTC[A/G]TCTTTTGCTG	S	A	G	S	S
G1544u12	WIAF-13492	U59464	2222	PTCH, patched (Drosophila) homolog	AAGCACTATG[C/T]TCCTTTCTCTC	M	C	T	A	V
G1544u13	WIAF-13500	U59464	1686	PTCH, patched (Drosophila) homolog	TCTTCATGGC[C/T]CGGTTAATCC	S	C	T	A	A
G1545u1	WIAF-12032	HT0473	1835	RAG1, recombination activating gene 1	GGACATGGAA[G/A]AAGACATCTT	M	G	A	E	K
G1545u2	WIAF-12035	HT0473	2519	RAG1, recombination activating gene 1	TGACATTGGC[A/G]ATGCAGCTGA	M	A	G	N	D
G1545u3	WIAF-12046	HT0473	3045	RAG1, recombination activating gene 1	CGGAAAATGA[A/G]TGCACGGCAG	M	A	G	N	S
G1545u4	WIAF-12047	HT0473	3146	RAG1, recombination activating gene 1	TCATAATGCA[T/C]TAAAAACCTC	S	T	C	L	L
G1545u5	WIAF-12075	HT0473	2513	RAG1, recombination activating gene 1	CCACTGTGAC[A/T]TTGGCAATGC	M	A	T	I	F
G1545u6	WIAF-13484	HT0473	1322	RAG1, recombination activating gene 1	GTGCTGACT[C/T]GGAGAGCTCA	M	C	T	R	W
G1545u7	WIAF-13494	HT0473	2571	RAG1, recombination activating gene 1	GAAGTGATA[A/G]GAATCCCAAT	M	A	G	K	R
G1545u8	WIAF-13498	HT0473	1018	RAG1, recombination activating gene 1	TTCTGGCTGA[C/A]CCTGTGGAGA	M	C	A	D	E
G1545u9	WIAF-13499	HT0473	2782	RAG1, recombination activating gene 1	ATCTTTACCT[G/C]AAGATGAAAAC	S	G	C	L	L
G1548u1	WIAF-12015	HT4999	133	IFI27, interferon, alpha- inducible protein 27	CTCTGCCGTA[G/A]TTTTCGCCCT	M	G	A	V	I
G1548u2	WIAF-13482	HT4999	380	IFI27, interferon, alpha- inducible protein 27	ATCCTGGGCT[C/T]CATTTGGTCT	M	C	T	S	F
G1548u3	WIAF-13483	HT4999	135	IFI27, interferon, alpha- inducible protein 27	CTGCCGTAGT[T/C]TTGCCCTGG	S	T	C	V	V

002060-2445950

G155u1	WIAF-11634	HT3962	991	CHC1, chromosome condensation 1	AGCTGGATGT [G/A] CCTGTGTAA	S	G	A	V	V
G155u2	WIAF-11635	HT3962	1271	CHC1, chromosome condensation 1	CGGCTTCGGC [C/T] TCTCAAATA	M	C	T	L	F
G155u3	WIAF-11636	HT3962	1192	CHC1, chromosome condensation 1	GCCGGGGCCA [C/T] GTGAGATTCC	S	C	T	H	H
G155u4	WIAF-11637	HT3962	1267	CHC1, chromosome condensation 1	TGTACGGCTT [C/T] GGCCTCTCCA	S	C	T	F	F
G155u5	WIAF-11649	HT3962	1657	CHC1, chromosome condensation 1	TGATGGGCAA [A/G] CAGCTGGAGA	S	A	G	K	K
G1550u1	WIAF-12057	MT6038	611	LYN, v-yes-1 Yamaguchi sarcoma viral related oncogene homolog	GCAAAGTCCC [T/G] TTTAAACAAA	M	T	G	L	R
G1550u2	WIAF-12061	MT6038	1371	LYN, v-yes-1 Yamaguchi sarcoma viral related oncogene homolog	TGGCATAACAT [C/T] GAGCGGAAGA	S	C	T	I	I
G1550u3	WIAF-12080	MT6038	1059	LYN, v-yes-1 Yamaguchi sarcoma viral related oncogene homolog	AAAGGCTTGG [C/T] GCTGGGCAGT	S	C	T	G	G
G1550u4	WIAF-12081	MT6038	996	LYN, v-yes-1 Yamaguchi sarcoma viral related oncogene homolog	AGCCACAGAA [G/A] CCATGGGATA	S	G	A	K	K
G1552u1	WIAF-12030	HT4578	2355	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	CCTGCTATTT [A/T] AAAGACTTCT	N	A	T	K	*
G1552u2	WIAF-12031	HT4578	2231	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	ACAAAGTTGA [C/T] TTAGAAGAGA	S	C	T	D	D
G1552u3	WIAF-12040	HT4578	617	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TCATGAGCTT [T/C] GGTATCCTTA	S	T	C	F	F
G1552u4	WIAF-12063	HT4578	1723	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TCATGTAACA [A/G] AAAATCAAAT	M	A	G	K	R
G1552u5	WIAF-12064	HT4578	1732	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	AAAAATCAA [A/G] TGTAATAGAT	M	A	G	N	S
G1552u6	WIAF-12065	HT4578	1660	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TTACCATGTA [A/G] AGTAAGTAAT	M	A	G	K	R

G1552u7	WIAF-12066	HT4578	1975	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	GAACGATACA [A/G] TAGTCAAATG	M	A	G	N	S
G1552u8	WIAF-12067	HT4578	1881	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TTTAGAGGAT [G/T] CAACACTACA	M	G	T	A	S
G1552u9	WIAF-12068	HT4578	2454	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TTTAGACGTT [T/A] TATATAAAAT	M	T	A	L	I
G1552u10	WIAF-12069	HT4578	2457	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	AGACGTTTTA [T/C] ATAAATGAC	M	T	C	Y	H
G1552u11	WIAF-12082	HT4578	2557	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	ATACCAGGAG [T/C] TTCAATTACT	M	T	C	V	A
G1552u12	WIAF-12083	HT4578	971	PMS1, postmeiotic segregation increased (S. cerevisiae) 1	TTTCTTTTCT [G/T] AAAATCGATG	S	G	T	L	L
G1554u1	WIAF-12028	HT4161	1500	ELK3, ELK3, ETS-domain protein (SRF accessory protein 2) NOTE: Symbol and name provisional.	CTCAGAATC [C/T] TGAAGCGTC	S	C	T	S	S
G1554u2	WIAF-12059	HT4161	1380	ELK3, ELK3, ETS-domain protein (SRF accessory protein 2) NOTE: Symbol and name provisional.	CTGCCAGGCT [G/A] CAAGGGCCAA	S	G	A	L	L
G1554u3	WIAF-12060	HT4161	1436	ELK3, ELK3, ETS-domain protein (SRF accessory protein 2) NOTE: Symbol and name provisional.	CACATGCCAG [T/C] GCCAATCCCC	M	T	C	V	A
G1562u1	WIAF-12024	HT28220	804	PDCD1, programmed cell death 1	GGGGCTGAGC [T/C] GACGGCCCTC	S	T	C	A	A
G1562u2	WIAF-13488	HT28220	644	PDCD1, programmed cell death 1	GACCCCTCAG [C/T] CGTGCTGTG	M	C	T	A	V
G1563u1	WIAF-13493	HT1187	1748	EGFR, epidermal growth factor receptor (avian erythroblastic leukemia viral (v-erb-b) oncogene homolog)	CCGAGGCCCA [G/A] CGACTGCGTC	M	G	A	R	K

G1563u2	WIAF-13497	HT1187		EGFR, epidermal growth factor receptor (avian erythroblastic leukemia viral (v-erb-b) oncogene homolog)	2073	ACGGATGCAC [T/A] GGGCCAGGTC	S	T	A	T	T
G1566u1	WIAF-12016	HT27594		PCD2, programmed cell death 2	235	GCGCCGCTGC [C/G] TGGCCGCCCG	M	C	G	P	R
G1566u2	WIAF-12033	HT27594		PCD2, programmed cell death 2	904	TTGGAATTC [A/G] GGTCAATGCCT	M	A	G	Q	R
G1566u3	WIAF-12041	HT27594		PCD2, programmed cell death 2	331	AATCAACTAC [C/T] CAGGAAAAAC	M	C	T	P	L
G1566u4	WIAF-12071	HT27594		PCD2, programmed cell death 2	649	CCTGAGTTG [T/C] GGAAAAGGAA	M	T	C	V	A
G1566u5	WIAF-12072	HT27594		PCD2, programmed cell death 2	633	ACAAGATGAG [A/T] TTATGCCTGA	M	A	T	I	F
G1567u1	WIAF-12042	M95936		AKT2, v-akt murine thymoma viral oncogene homolog 2	293	GAGAGCCGC [G/A] ACCCAACACC	M	G	A	R	Q
G1572u1	WIAF-12212	HT3998		proto-oncogene c-abl, tyrosine protein kinase, alt. transcript 2	1894	TGTTCCAGGA [A/G] TCCAGTATCT	S	A	G	E	E
G1572u2	WIAF-12233	HT3998		proto-oncogene c-abl, tyrosine protein kinase, alt. transcript 2	3694	AGCTTCAGAT [C/T] TGCCCGGCGA	S	C	T	I	I
G1572u3	WIAF-12234	HT3998		proto-oncogene c-abl, tyrosine protein kinase, alt. transcript 2	3721	GCAGTGGTCC [G/A] GGGCCCACTC	S	G	A	P	P
G1573u1	WIAF-12021	HT0642		CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	343	TCATGGACAA [G/C] GTGTGCGGT	M	G	C	K	N
G1573u2	WIAF-12022	HT0642		CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	363	TTGTGTGAGA [A/T] CCCAAGCTG	M	A	T	N	I
G1573u3	WIAF-12034	HT0642		CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	2364	AATATTCACT [C/T] CCAGGCGCCA	M	C	T	S	F
G1573u4	WIAF-12049	HT0642		CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	387	CTAAGAATA [G/A] CCCACCTTAT	M	G	A	S	N
G1573u5	WIAF-12050	HT0642		CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	947	AACTCAATCCT [G/A] GCTACATGCG	M	G	A	G	S

G1573u6	WIAF-12070	HT0642	2740	CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	TCGAGAACCT[C/T]ATGAGTCAGG	S	C	T	L	L
G1573u7	WIAF-12073	HT0642	661	CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	TCTTTCCAAAG[T/C]GGACTCTTTC	S	T	C	S	S
G1573u8	WIAF-12074	HT0642	2569	CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	CTCTGGATGG[T/C]GATCCTACAA	S	T	C	G	G
G1573u9	WIAF-13486	HT0642	2006	CBL, Cas-Br-M (murine) ecotropic retroviral transforming sequence	CCGGCACTCA[C/T]TTCATTTC	M	C	T	L	F
G1574u1	WIAF-12037	HT1508	2493	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	AGCGGCCAG[C/T]TTCAGCACCA	S	C	T	S	S
G1574u2	WIAF-12051	HT1508	189	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	CCCAGCGGGT[C/T]AAGAGTGACA	S	C	T	V	V
G1574u3	WIAF-12052	HT1508	1441	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	GAAGCCCTG[C/T]ATGAGCAGCT	M	C	T	H	Y
G1574u4	WIAF-12053	HT1508	2202	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	GAGAGGAAGC[C/T]GATGGGGTCT	S	C	T	A	A
G1574u5	WIAF-12054	HT1508	2088	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	CTGCTGGCAT[G/T]GAGTACCTGG	M	G	T	M	I
G1574u6	WIAF-12078	HT1508	1577	FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-av) oncogene homolog	GATGCTCTGC[C/T]CCGGCACTTC	M	C	T	P	L

002060-2449960

G1574u7	WIAF-13495	HT1508		FES, feline sarcoma (Snyder-Theilen) viral (v-fes)/Fujinami avian sarcoma (PRCII) viral (v-fes) oncogene homolog	579	GTGACAAGGC [T/C] AAGGACAAGT	S	T	C	A	A
G1575u1	WIAF-12079	HT1052		FGF, Gardner-Rasheed feline sarcoma viral (v-fgr) oncogene homolog	963	TGGGCACCGG [C/T] TGCTTCGGGG	S	C	T	G	G
G1575u2	WIAF-13487	HT1052		FGF, Gardner-Rasheed feline sarcoma viral (v-fgr) oncogene homolog	232	CAGAAGCTAC [G/A] GGGCAGCAGA	M	G	A	G	R
G1585u1	WIAF-12017	HT1675		CRK, v-crck avian sarcoma virus Crt10 oncogene homolog	996	TGGATCAACA [G/A] AATCCGATG	S	G	A	Q	Q
G1585u2	WIAF-12036	HT1675		CRK, v-crck avian sarcoma virus Crt10 oncogene homolog	446	ACTACAACGT [T/C] GATAGAACCA	M	T	C	L	S
G1587u1	WIAF-12023	HT0590		1473 proto-oncogene db1	1473	GGCCAATCCA [A/G] TTTGTGGTAC	S	A	G	Q	Q
G1587u2	WIAF-12025	HT0590		2549 proto-oncogene db1	2549	GTCAGGCTT [C/T] TAACTGTAGAT	M	C	T	S	F
G1587u3	WIAF-12026	HT0590		2828 proto-oncogene db1	2828	GCATCACAAT [C/T] TGCAGAAATC	M	C	T	S	F
G1587u4	WIAF-12038	HT0590		982 proto-oncogene db1	982	AAATTCTCAG [G/C] AGCTATTATC	M	G	C	E	Q
G1587u5	WIAF-12039	HT0590		2343 proto-oncogene db1	2343	AACCAATGCA [G/T] CGACACCTTT	M	G	T	Q	H
G1587u6	WIAF-12048	HT0590		683 proto-oncogene db1	683	GACACTGAAG [G/A] AGCTGTCTAGT	M	G	A	G	E
G1587u7	WIAF-12055	HT0590		2686 proto-oncogene db1	2686	TTCTCTTCAG [C/T] AGAATGATGA	N	C	T	Q	*
G1587u8	WIAF-13485	HT0590		2136 proto-oncogene db1	2136	ACTGTGAAGG [T/A] TCTGCTCTGT	S	T	A	G	G
G1587u9	WIAF-13496	HT0590		1566 proto-oncogene db1	1566	AAAATCAGAG [C/T] AACTTAAAAA	S	C	T	S	S
G159u1	WIAF-11616	HT4209		RAD23B, RAD23 (S. cerevisiae) homolog B	1059	AGTACTGGGG [C/T] TCCTCAGTCT	M	C	T	A	V
G1590u1	WIAF-13897	HT2455		ETS2, v-ets avian erythroblastosis virus E26 oncogene homolog 2	1257	GCCAGTCTCT [C/G] TGCCTCAATA	S	C	G	L	L
G1590u2	WIAF-13913	HT2455		ETS2, v-ets avian erythroblastosis virus E26 oncogene homolog 2	1107	ATTCTGGGAC [T/G] CCCAAGACC	S	T	G	T	T
G1590u3	WIAF-13914	HT2455		ETS2, v-ets avian erythroblastosis virus E26 oncogene homolog 2	1314	GGAGTGACCC [A/G] GTGGAGCAAG	S	A	G	P	P
G1591u1	WIAF-13924	HT2333		HRAS, v-Ha-ras Harvey rat sarcoma viral oncogene homolog	417	TCCAGAACCA [T/C] TTTGTGGACG	S	T	C	H	H
G1595u1	WIAF-12262	HT33778		proto-oncogene l-myc, alt. transcript 1	1302	GCATACCTCA [G/C] TGGCTACTAA	M	G	C	S	T

G1597u1	WIAF-12243	HT0410	900	MAS1, MAS1 oncogene RAD23A, RAD23 (<i>S. cerevisiae</i>) 690 homolog A	CCATCTTGGT [C/T] GTGAAGATCC	S	C	T	V	V
G1601u1	WIAF-11630	HT4247			AGAGCCAGGT [A/G] TCGGAGCAGC	S	A	G	V	V
G1602u1	WIAF-14180	HT1903			GTGCGCGGG [C/A] CCAGCAATA	M	C	A	P	T
				REL, v-rel avian reticuloendotheliosis viral oncogene homolog						
G1604u1	WIAF-12319	HT2788	1182	receptor (TNFRSF) - interacting serine-threonine kinase 1	CCTCCAAAAG [T/C] GCTGGGATTA	S	T	C	S	S
G1609u1	WIAF-12358	HT33646	348		GAGCAGGGT [C/T] TCCCATGACC	S	C	T	V	V
				DNA repair and recombination homolog RAD52						
G1611u1	WIAF-11654	HT4251	1522		TATGATCCAT [C/T] TTAACCTAGG	M	C	T	S	F
G1610a1	WIAF-12101	HT27727	501	replication protein Rpa4, 30 kDa	TGCAACTCCT [G/A] CTATTAAAGAC	M	G	A	A	T
G1610a2	WIAF-12102	HT27727	554	replication protein Rpa4, 30 kDa	TACCGTGTA [C/T] GTGAACCAAGC	S	C	T	N	N
G1610u3	WIAF-12307	HT27727	450	replication protein Rpa4, 30 kDa	TTCTGTGCT [G/A] ATGAGCGGAG	M	G	A	D	N
G1610u4	WIAF-12320	HT27727	1037	replication protein Rpa4, 30 kDa	TGATTCATGA [G/C] TGTCCTCATC	M	G	C	E	D
G1610u5	WIAF-12321	HT27727	857	replication protein Rpa4, 30 kDa	TAGAGGACAT [G/A] AACGAGTTCA	M	G	A	M	I
G1610u6	WIAF-12343	HT27727	539	replication protein Rpa4, 30 kDa	GAATTCAGGA [C/T] GTTGTAACGT	S	C	T	D	D
G1630u1	WIAF-12302	HT3563	4312	DCC, deleted in colorectal carcinoma	ACTCATGAAG [C/T] AGCTTAATGC	N	C	T	Q	*
G1632u1	WIAF-13572	HT27355	742	tumor suppressor, PDGF receptor beta-like	TTTATGACAT [G/C] AAGCGGGGCT	M	G	C	M	I
G1632u2	WIAF-13584	HT27355	1102	tumor suppressor, PDGF receptor beta-like	TGGAAGACTT [C/T] GAGACGATG	S	C	T	F	F
G1632u3	WIAF-13601	HT27355	258	tumor suppressor, PDGF receptor beta-like	AAGACGAGT [C/T] TATCATGATG	M	C	T	S	F
				FER, fer (fps/fes related) tyrosine kinase (phosphoprotein NCP94)						
G1633u1	WIAF-13957	HT1778	1263		TTCAGGCAAA [T/C] GAGATCATGT	S	T	C	N	N
G1633u2	WIAF-13958	HT1778	2407	FER, fer (fps/fes related) tyrosine kinase (phosphoprotein NCP94)	TATGTTGTAT [C/T] TCGAGAGTAA	M	C	T	L	F
G1634u1	WIAF-13505	HT3216	1569	ELK1, ELK1, member of ETS oncoqene family	TCTCGACCCC [C/T] GTGTGTCTCT	S	C	T	P	P

G1634u2	WIAF-13858	HT3216	456	ELK1, ELK1, member of ETS oncogene family	GGCTGTGGGG [A/G]CTACGCAAGA	S	A	G	G	G
G1634u3	WIAF-13859	HT3216	745	ELK1, ELK1, member of ETS oncogene family	AGGCCAGGC [G/A]GTTTGGCAGG	M	G	A	G	S
G1638u1	WIAF-14172	HT1224	98	uracil-DNA glycosylase	GCTGGGACCT [G/C]TTCACAAAAT	-	G	C	-	-
G1643u1	WIAF-13517	HT3751	629	DXS648E, DNA segment on chromosome X (unique) 648 expressed sequence	TACATCCCA [G/A]TCGTGGCCCT	M	G	A	S	N
G1645u1	WIAF-14087	D21089	363	XPC, xeroderma pigmentosum, complementation group C	AAACCTCAA [G/A]GTTATAAAGG	S	G	A	K	K
G1645u2	WIAF-14088	D21089	2166	XPC, xeroderma pigmentosum, complementation group C	TGCATTCCAG [G/A]CACACGTGGC	S	G	A	R	R
G1645u3	WIAF-14089	D21089	1580	XPC, xeroderma pigmentosum, complementation group C	GGGAGCCATC [G/A]TAAGGACCCA	M	G	A	R	H
G1645u4	WIAF-14090	D21089	1601	XPC, xeroderma pigmentosum, complementation group C	AGCTTGCAG [T/C]GGCATCCTCA	M	T	C	V	A
G1645u5	WIAF-14091	D21089	2920	XPC, xeroderma pigmentosum, complementation group C	CCCATTTGAG [A/C]AGCTGTGAGC	M	A	C	K	Q
G1645u6	WIAF-14103	D21089	405	XPC, xeroderma pigmentosum, complementation group C	ATGACCTCAG [G/A]GACTTTCCAA	S	G	A	R	R
G1645u7	WIAF-14104	D21089	151	XPC, xeroderma pigmentosum, complementation group C	GGGACGCGAA [C/G]TGGCGAGCCA	M	C	G	L	V
G1645u8	WIAF-14105	D21089	2133	XPC, xeroderma pigmentosum, complementation group C	AAGCGTCTA [C/T]TCCAGGGATT	S	C	T	Y	Y
G167u1	WIAF-11632	HT4579	83	PMS2L8, postmeiotic segregation increased 2-like 8	CCTATTGATC [G/A]GAAGTCAGTC	M	G	A	R	Q
G167u2	WIAF-11633	HT4579	219	PMS2L8, postmeiotic segregation increased 2-like 8	GAGTGGATCT [T/C]ATTGAAGTTT	S	T	C	L	L
G167u3	WIAF-11644	HT4579	768	PMS2L8, postmeiotic segregation increased 2-like 8	TGCCCCCTAG [T/C]GACTCCGTGT	S	T	C	S	S

004060" 2446560

G167u4	WIAF-11622	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	1645	GAAAGCGCCT [G/A]AAACTGACGA	M	G	A	E	K
G167u5	WIAF-11645	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	1512	ACTCGGGGCA [C/T]GGCAGCACTT	S	C	T	H	H
G167u6	WIAF-11646	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	1619	TCGCAGGAAC [A/G]TGIGGACTCT	M	A	G	H	R
G167u7	WIAF-11647	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	1432	CGTCCTGAGA [C/T]CTCAGAAAGA	M	C	T	P	S
G167u8	WIAF-11625	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	2490	GGACTGCTCT [T/C]AACACAAGCG	S	T	C	L	L
G167u9	WIAF-11619	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	804	TGAGCTGTTC [G/C]GATGCTCTGC	S	G	C	S	S
G167u10	WIAF-11623	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	1555	CATCCCAGAC [A/G]CGGGCAGTCA	M	A	G	T	A
G167u11	WIAF-11624	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	2364	CCTTCGGACC [C/T]CAGGACGTGC	S	C	T	P	P
G167u12	WIAF-11626	HT4579		PMS2L8, postmeiotic segregation increased 2-like 8	2348	ACTAGTAAAA [A/G]CTGGACCTTC	M	A	G	N	S
G181u1	WIAF-11697	HT48793		ERCC4, excision repair cross- complementing rodent repair deficiency, complementation group 311 4	311 4	ATATTTCGGA [C/T]AAGTAGGATA	M	C	T	T	I
G181u2	WIAF-11698	HT48793		ERCC4, excision repair cross- complementing rodent repair deficiency, complementation group 295 4	295 4	CACACAAGGT [G/C]GTGTTATATT	M	G	C	G	R
G181u3	WIAF-11699	HT48793		ERCC4, excision repair cross- complementing rodent repair deficiency, complementation group 234 4	234 4	TTGAACACCT [C/T]CCTCGCGGTG	S	C	T	L	L

002000"2445960

G181u4	WIAF-11704	HT48793	808 4	ERCC4, excision repair cross-complementing rodent repair deficiency, complementation group	TTTGTGGCAC [C/T] AGCTTGGAGC	N	C	T	Q	*
G181u5	WIAF-11705	HT48793	640 4	ERCC4, excision repair cross-complementing rodent repair deficiency, complementation group	TTCTATGACA [C/T] CTACCATGCT	M	C	T	P	S
G181u6	WIAF-11670	HT48793	1117 4	ERCC4, excision repair cross-complementing rodent repair deficiency, complementation group	AGAAAGCAAC [C/T] CAAAGTGGGA	M	C	T	P	S
G185u1	WIAF-11668	HT5122	319 IIB	ACVR2B, activin A receptor, type IIB	TCTGCAACGA [G/A] CGCTTCACTC	S	G	A	E	E
G185u2	WIAF-11707	HT5122	70 IIB	ACVR2B, activin A receptor, type IIB	AGACACGGGA [G/C] TGCATCTACT	M	G	C	E	D
G185u3	WIAF-11672	HT5122	812 IIB	ACVR2B, activin A receptor, type IIB	CCTCACGGAT [T/C] ACCTCAAGGG	M	T	C	Y	H
G185u4	WIAF-13542	X77533	1109 IIB	ACVR2B, activin A receptor, type IIB	GGCTCCTGAG [G/A] TGCTCGAGGG	M	G	A	V	M
G185u5	WIAF-13558	X77533	997 IIB	ACVR2B, activin A receptor, type IIB	TGCTGAAGAG [C/T] GACCTCACAG	S	C	T	S	S
G187u1	WIAF-11669	HT97400	183 androgen	CXCR4, chemokine (C-X-C motif), receptor 4 (fusin)	CCAGAGACAG [C/T] GCGACCCGGA	M	C	T	R	C
G191u1	WIAF-10176	AF025375	414	CCR2, chemokine (C-C motif) receptor 2	ACCTGGCCAT [C/T] GTCCACGCCA	S	C	T	I	I
G193u1	WIAF-10178	D29984	231	CCR2, chemokine (C-C motif) receptor 2	AGTGCTTGAC [T/A] GACATTAC	S	T	A	T	T
G193u2	WIAF-10179	D29984	190	CCR2, chemokine (C-C motif) receptor 2	CATGCTGGTC [G/A] TCCTCATCTT	M	G	A	V	I
G194u1	WIAF-10211	D43767	121	SCYA17, small inducible cytokine subfamily A (Cys-Cys), member 17	ACATCCACGC [A/C] GCTCGAGGGA	S	A	C	A	A
G197u1	WIAF-10167	D50403	1515	NRAMP1, natural resistance-associated macrophage protein 1 (might include Leishmaniasis)	GGTGCTAGTC [T/C] GCGCCATCAA	M	T	C	C	R
G197u2	WIAF-10173	D50403	1629	NRAMP1, natural resistance-associated macrophage protein 1 (might include Leishmaniasis)	CACCTACCTG [G/C] TCTGGACCTG	M	G	C	V	L

G20u1	WIAF-10249	U14722	896 IB	ACVR1B, activin A receptor, type IB	CGGTACACAG [T/C] GACAATTGAG	M	T	C	V	A
G20u2	WIAF-10250	U14722	866 IB	ACVR1B, activin A receptor, type IB	GAGCAGGGT [C/T] CCTGTTGAT	M	C	T	S	F
G20u3	WIAF-10251	U14722	1391 IB	ACVR1B, activin A receptor, type IB	CAGAGTTATG [A/T] GGCACATGCGG	M	A	T	E	V
G20u4	WIAF-10252	U14722	1236 IB	ACVR1B, activin A receptor, type IB	TATATTGGGA [G/C] ATTGCTCGAA	M	G	C	E	D
G20u5	WIAF-10261	U14722	518 IB	ACVR1B, activin A receptor, type IB	GAGATGTGTC [T/C] CTCCAAAGAC	M	T	C	L	P
G207a1	WIAF-10516	L25259	866 2)	Human CTLA4 counter-receptor (B7-2) mRNA, complete cds.	AGCTGTACTT [C/T] CAACAGTTAT	M	C	T	P	S
G208u1	WIAF-10204	L31581	85	CCR7, chemokine (C-C motif) receptor 7	GGGGAACCA [A/G] TGAAGAGCGT	M	A	G	M	V
G211u1	WIAF-10213	M24545	174	SCYA2, small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse sig-je)	TCACCTGCTG [T/C] TATAACTTCA	S	T	C	C	C
G214u1	WIAF-10191	M27533	452	CD80, CD80 antigen (CD28 antigen ligand 1, B7-1 antigen)	TGAAGAAGT [G/A] GCAACGCTGT	S	G	A	V	V
G215u1	WIAF-11659	M28393	822	PRF1, perforin 1 (preforming protein)	GCATCTCTGC [C/T] GAAGCCAAGG	S	C	T	A	A
G215u2	WIAF-11723	M28393	159	PRF1, perforin 1 (preforming protein)	TGACCAGCCT [C/T] CGCCGCTCGG	S	C	T	L	L
G215u3	WIAF-11724	M28393	96	PRF1, perforin 1 (preforming protein)	CAGAGTGCAA [G/A] CGCAGCCACA	S	G	A	K	K
G215u4	WIAF-11725	M28393	1377	PRF1, perforin 1 (preforming protein)	ATAACAACCC [C/T] ATCTGGTCAG	S	C	T	P	P
G215u5	WIAF-11726	M28393	1326	PRF1, perforin 1 (preforming protein)	TGAAGCTCTT [C/T] TTGTGGCC	S	C	T	F	F
G215u6	WIAF-11727	M28393	1076	PRF1, perforin 1 (preforming protein)	CGGCGGGAGG [C/T] ACTGAGGAGG	M	C	T	A	V
G217u1	WIAF-11691	M31932	649	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	GCAGCTCTTC [A/G] CCAATGGGA	S	A	G	S	S
G217u2	WIAF-11692	M31932	625	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	TCAGTGTCCA [A/G] GTGCCAGCA	S	A	G	Q	Q

G217u3	WIAF-11712	M31932	332	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	GACTGGCCAG[A/C]CCAGCCTCAG	M	A	C	T	P
G217u4	WIAF-11713	M31932	101	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	GCCTTCTGCA[G/T]ACAGTCAAGC	M	G	T	D	Y
G218u1	WIAF-10184	M36712	677	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	TTTTACAAAT[A/G]AGCAGAGAAT	N	A	G	*	*
G218u2	WIAF-10188	M36712	326	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	GCTGTGTTTC[G/C]GGATGCAAGC	M	G	C	R	P
G218u3	WIAF-10189	M36712	196	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	CAGTAACATG[C/T]GCATCTACTG	M	C	T	R	C
G218u4	WIAF-10190	M36712	225	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	AGCGCCAGGC[A/C]CCGAGCAGTG	S	A	C	A	A
G218u5	WIAF-10194	M36712	583	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	GGTGGCTGGC[G/A]TCCTGGTTCT	M	G	A	V	I
G218u6	WIAF-10208	M36712	372	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	TGAAGCCGGA[A/G]GACAGTGGCA	S	A	G	E	E
G218u7	WIAF-10209	M36712	400	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	CTGCATGATC[G/T]TCGGGAGCCC	M	G	T	V	F
G218u8	WIAF-10210	M36712	270	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	TCTGGGATTC[C/T]GCAAAAGGGA	S	C	T	S	S
G218a9	WIAF-10518	M36712	618	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	GAGTGGCCAT[C/G]CACCTGTGCT	M	C	G	I	M
G218a10	WIAF-13223	M36712	556	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	TTGTAGCCCC[A/G]TCACCCTTGG	M	A	G	I	V
G218a11	WIAF-13224	M36712	836	CD8B1, CD8 antigen, beta polypeptide 1 (p37)	CTGTGTGTGA[T/C]CTGCATGGGA	-	T	C	-	-
G22u1	WIAF-10301	U86136	6719	Human telomerase-associated protein TP-1 mRNA, complete cds.	GGTGGTAACC[G/A]TCGGGCTAGA	M	G	A	V	I
G22u2	WIAF-10302	U86136	7537	Human telomerase-associated protein TP-1 mRNA, complete cds.	CTGATGGGAT[C/G]CTATGGAACC	M	C	G	I	M
G22u3	WIAF-10311	U86136	1798	Human telomerase-associated protein TP-1 mRNA, complete cds.	ATGATGCCAT[T/C]GATGCCCTCG	S	T	C	I	I
G22u4	WIAF-10312	U86136	2397	Human telomerase-associated protein TP-1 mRNA, complete cds.	CTGTCTCTGG[C/T]TGGCCAAAGG	M	C	T	A	V

002060-2446960

G22u5	WIAF-10313	U86136	3289	Human telomerase-associated protein TP-1 mRNA, complete cds.	AGAAAGGGAT [A/C] ACCTGCCGCA	S	A	C	I	I
G22u6	WIAF-10314	U86136	3242	Human telomerase-associated protein TP-1 mRNA, complete cds.	AGAGGCCGCA [T/C] GTCGGATCTC	M	T	C	C	R
G22u7	WIAF-10315	U86136	4482	Human telomerase-associated protein TP-1 mRNA, complete cds.	CCGTTTGCCT [G/A] CCTCGTCCAG	M	G	A	C	Y
G22u8	WIAF-10316	U86136	4363	Human telomerase-associated protein TP-1 mRNA, complete cds.	GTTTGACTGT [G/A] GACCAGCTGC	S	G	A	V	V
G22u9	WIAF-10317	U86136	4230	Human telomerase-associated protein TP-1 mRNA, complete cds.	GTGTCGTAGA [G/A] ACTCCGACCC	M	G	A	R	K
G22u10	WIAF-10318	U86136	4419	Human telomerase-associated protein TP-1 mRNA, complete cds.	GGGACTAAGA [G/C] CTGGGAAGAA	M	G	C	S	T
G22u11	WIAF-10319	U86136	5269	Human telomerase-associated protein TP-1 mRNA, complete cds.	TCTCCGATGA [T/C] ACACCTCTTC	S	T	C	D	D
G22u12	WIAF-10320	U86136	5015	Human telomerase-associated protein TP-1 mRNA, complete cds.	GCTGCTCTCC [C/T] GGAGATGGCA	M	C	T	R	W
G22u13	WIAF-10321	U86136	5133	Human telomerase-associated protein TP-1 mRNA, complete cds.	GTGGCCTTCT [C/T] CACCAATGGG	M	C	T	S	F
G22u14	WIAF-10322	U86136	7764	Human telomerase-associated protein TP-1 mRNA, complete cds.	ACAGCCCTCC [A/G] TGTGCTACCT	M	A	G	H	R
G22u15	WIAF-10323	U86136	7884	Human telomerase-associated protein TP-1 mRNA, complete cds.	TGCTTGGAAC [C/T] TTGGCTGGGC	M	C	T	P	L
G22u16	WIAF-10324	U86136	7744	Human telomerase-associated protein TP-1 mRNA, complete cds.	AGATTCACCTC [G/A] GGCCTGTCTCA	S	G	A	S	S
G22u17	WIAF-10337	U86136	1018	Human telomerase-associated protein TP-1 mRNA, complete cds.	CCATTGCTGC [T/C] TTCTTGCCGG	S	T	C	A	A
G22u18	WIAF-10338	U86136	1000	Human telomerase-associated protein TP-1 mRNA, complete cds.	TGGCCATAAA [C/A] ATCTTGGCAA	M	C	A	N	K

G22u19	WIAP-10339	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	1182	ATGACGGACA [A/G]ATTGCCCAG	M	A	G	K	R
G22u20	WIAP-10340	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	1939	AGCAGCTTCG [T/G]ATGCAATGA	S	T	G	R	R
G22u21	WIAP-10341	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	2227	TCACGAGGGC [G/A]GAGCAGGTGG	S	G	A	A	A
G22u22	WIAP-10342	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	2776	GGCGCAGCAT [C/T]CGGCTTTTCA	S	C	T	I	I
G22u23	WIAP-10343	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	2877	GCCCTCACC [G/A]TATCAGCCTT	M	G	A	R	H
G22u24	WIAP-10344	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	3087	TCAGGGCGCT [C/T]TGTGACAGAG	M	C	T	S	F
G22u25	WIAP-10345	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	3662	CAAGGTGGCA [C/T]CATTAGTCTT	M	C	T	P	S
G22u26	WIAP-10346	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	4762	TTTCGAAGTT [C/T]CTTACCAACC	S	C	T	F	F
G22u27	WIAP-10351	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	1737	CTCCAGCATG [G/C]GAAGTCGGTG	M	G	C	G	A
G22u28	WIAP-10352	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	3543	ACAGTGCAAC [A/G]GCTGATGCTG	M	A	G	Q	R
G22u29	WIAP-10353	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	4232	GTCTGAGAGA [C/T]TCCGACCCCT	M	C	T	L	F
G22u30	WIAP-10354	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	4523	GGAGGGCCCT [C/T]TGGAGCGCCC	S	C	T	L	L
G22u31	WIAP-10355	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	5333	TGGTTGTGGG [G/T]TGCTGAGAC	M	G	T	V	L
G22u32	WIAP-10356	U86136	Human telomerase-associated protein TP-1 mRNA, complete cds.	6208	AGCTGCTGAC [G/A]CGGCCACACA	S	G	A	T	T

002060" 2445960

G222u33	WIAF-10357	U86136		Human telomerase-associated protein TP-1 mRNA, complete cds.	7703		TAGTGAGCCA [A/G] CACCACATCT	M	A	G	T	A
G222u34	WIAF-10360	U86136		Human telomerase-associated protein TP-1 mRNA, complete cds.	3881		CATCGATGGG [G/A] CTGATAGGTT	M	G	A	A	T
G222u1	WIAF-11700	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	697		TGATGGGAT [G/C] GTGGAAGGGA	M	G	C	G	R
G222u2	WIAF-11701	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	708		GTGGAAGGGA [A/G] ACACACTTGG	S	A	G	E	E
G222u3	WIAF-11702	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	677		GAGGGAAGA [A/G] AATGAGGTGT	M	A	G	K	R
G222u4	WIAF-11706	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	1616		AAGAAATATA [T/C] ACTTGAGTGG	M	T	C	I	T
G222u5	WIAF-11667	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	1444		TGATCGCTAT [C/G] TAGCAACCTT	M	C	G	L	V
G222u6	WIAF-11708	M57230		IL6ST, interleukin 6 signal transducer (gp130, oncostatin M receptor)	981		TCTTAAAAATT [G/C] ACATGGACCA	M	G	C	L	F
G226u1	WIAF-11714	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	869		CACTGGGAGT [T/C] GCCATATCTG	S	T	C	V	V
G226u2	WIAF-11715	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	1749		AGATTATGAG [C/T] CTCCATTTGG	M	C	T	P	S
G226u3	WIAF-11716	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	1601		TGGGAAGTGC [A/G] AGATACATGG	S	A	G	A	A
G226u4	WIAF-11721	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	1256		TACTCCAGTT [C/G] CTGAGGGCTG	M	C	G	F	L
G226u5	WIAF-11722	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	1502		TGCTGAAGAA [C/T] GACCTAACCT	S	C	T	N	N
G226u6	WIAF-11671	M85079		TGFBR2, transforming growth factor, beta receptor II (70-80kD)	888		TGTCATCATC [A/C] TCTTCTACTG	M	A	C	I	L

G226u7	WIAF-11674	M85079	TGFR2, transforming growth factor, beta receptor II (70-80kd)	1425	CCTCCACAGT [G/A] ATCACACTCC	M	G	A	D	N
G227u1	WIAF-10197	M86511	CD14, CD14 antigen	685	CCTGTCTGAC [A/G] ATCTGGACT	M	A	G	N	D
G227u2	WIAF-10212	M86511	CD14, CD14 antigen	497	GAAGCCACAG [G/A] ACTTGCACTT	M	G	A	G	E
G2278u1	WIAF-14117	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	959	AGATAAATBA [T/C] GCGGCTGT	S	T	C	N	N
G2278u2	WIAF-14118	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	781	GGGTGGATGT [C/T] TTCACCCAAC	M	C	T	S	F
G2278u3	WIAF-14119	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	641	CTGAGACGTA [C/T] GGACCCCACT	S	C	T	Y	Y
G2278u4	WIAF-14121	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	1185	TGGTTATGGG [C/A] CAAATGGATG	M	C	A	P	T
G2278u5	WIAF-14133	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	1532	TCTGGTTAT [C/G] AAAACTGAAA	M	C	G	I	M
G2278u6	WIAF-14134	AF034611	CUBN, cubilin (intrinsic factor-cobalamin receptor)	2208	GCCTTTCAC [C/T] ACACCCAGCA	M	C	T	H	Y
G228u1	WIAF-10199	U00672	IL10RA, interleukin 10 receptor, alpha	586	GCTAAGTGCC [G/A] GGAACCTTCA	S	G	A	P	P
G228u2	WIAF-10200	U00672	IL10RA, interleukin 10 receptor, alpha	731	AGAGGAGTGC [A/G] TCTCCCTCAC	M	A	G	I	V
G2280u1	WIAF-13970	AJ001515	RXR3, ryanodine receptor 3	1747	CAGGTATCTT [G/A] GAAGTTTTCG	S	G	A	L	L
G2280u2	WIAF-13974	AJ001515	RXR3, ryanodine receptor 3	8593	TAGAAGCCAT [T/C] GTCACGAGTG	S	T	C	I	I
G2282u1	WIAF-12694	D00726	FECH, ferrochelatase (protoporphyrin)	263	ACATGGGAGG [C/T] CCTGAACTC	S	C	T	G	G
G2282u2	WIAF-12695	D00726	FECH, ferrochelatase (protoporphyrin)	514	TACTATATGT [G/A] ATTTCCGTAC	M	G	A	G	E
G2285u1	WIAF-12688	D16611	CPO, coproporphyrinogen oxidase (coproporphyrin, harderoporphyrin)	673	AGAAGACGCT [G/A] TCCATTTC	M	G	A	V	I
G2285u2	WIAF-12689	D16611	CPO, coproporphyrinogen oxidase (coproporphyrin, harderoporphyrin)	783	ATCTGGAGA [G/A] CGGGGGGCA	S	G	A	E	E
G2287u1	WIAF-12687	D28472	PTGER4, prostaglandin E receptor 4 (subtype EP4)	502	GGGCTCAGC [C/T] TCTTTGAGT	M	C	T	L	F
G2287u2	WIAF-12691	D28472	PTGER4, prostaglandin E receptor 4 (subtype EP4)	1309	TGMAAATGGC [C/T] TTGGAGGAG	M	C	T	L	F
G2287u3	WIAF-12707	D28472	PTGER4, prostaglandin E receptor 4 (subtype EP4)	243	AGGAGACGAC [C/T] TTCTACAGC	S	C	T	T	T

G2287u4	WIAF-12710	D28472	1343	PTGER4, prostaglandin E receptor 4 (subtype EP4)	GGTGTGCGCTG [G/A] CATGGGCGCTG	M	G	A	G	D
G229u1	WIAF-10185	U16752	202	SDF1, stromal cell-derived factor 1	CATGTTGCCA [G/A] AGCCAAGCTC	M	G	A	R	K
G2295u1	WIAF-12727	D89079	613	LTBR, leukotriene b4 receptor (chemokine receptor-like 1)	CTATGTCGTG [G/C] GAGTCAGCAT	M	G	C	G	R
G2295u2	WIAF-12728	D89079	1248	LTBR, leukotriene b4 receptor (chemokine receptor-like 1)	AGGGCACGGG [T/C] TCCGAGGCGT	S	T	C	G	G
G2295u3	WIAF-12753	D89079	1348	LTBR, leukotriene b4 receptor (chemokine receptor-like 1)	CCTCACTGCC [T/G] CCAGCCCTCT	M	T	G	S	A
G230u1	WIAF-10201	U31628	627	IL15RA, interleukin 15 receptor, alpha	ACAGCCAAGA [A/C] CTGGGAAGTC	M	A	C	N	T
G2300u1	WIAF-12735	J02959	102	LTA4H, leukotriene A4 hydrolase	ACCTGCACCT [G/T] CGCTGCAGCG	S	G	T	L	L
G2300u2	WIAF-12738	J02959	1380	LTA4H, leukotriene A4 hydrolase	CCTGGCTCTA [C/T] TCTCCTGGAC	S	C	T	Y	Y
G2302u1	WIAF-12741	J03037	627	CA2, carbonic anhydrase II	TCTGAAATCC [C/T] TGGATTACTG	S	C	T	L	L
G2302u2	WIAF-12742	J03037	819	CA2, carbonic anhydrase II	GCCACTGAAG [A/G] ACAGGCAAAAT	M	A	G	N	D
G2303u1	WIAF-12751	J03571	304	ALOX5, arachidonate 5-lipoxygenase	CGCTGAAGAC [G/A] CCCACGGGG	S	G	A	T	T
G2303u2	WIAF-12752	J03571	794	ALOX5, arachidonate 5-lipoxygenase	AGAGCTGCCC [G/A] AGAAGCTCCC	M	G	A	E	K
G2304u1	WIAF-12772	J03575	840	PDHA1, pyruvate dehydrogenase (lipoamide) alpha 1	TCCGAGAGGC [A/G] ACAAGGTTTG	S	A	G	A	A
G2304u2	WIAF-12779	J03575	1044	PDHA1, pyruvate dehydrogenase (lipoamide) alpha 1	CCAGTGTGGA [A/C] GAACATAAAGG	M	A	C	E	D
G2305u1	WIAF-12763	J03576	456	PDHB, pyruvate dehydrogenase (lipoamide) beta	TCTTCAGGGG [A/G] CCCAATGGTG	S	A	G	G	G
G2305u2	WIAF-12764	J03576	650	PDHB, pyruvate dehydrogenase (lipoamide) beta	GTTCTTTTGG [A/C] ATTTCCTCCG	M	A	C	E	A
G231u1	WIAF-10202	U32324	734	IL11RA, interleukin 11 receptor, alpha	CCAGGCGCTG [C/T] GGGTAGAGTC	M	C	T	R	W

G2312u1	WIAF-12762	J05096		ATP1A2, ATPase, Na+/K+ transporting, alpha 2 (+) polypeptide	3726	TCAGAAACCA [C/T] ACAGAGATCG	S	C	T	H	H
G2313u1	WIAF-12760	J05200		RYR1, ryanodine receptor 1 (skeletal)	6141	TGCAATTCAA [A/G] GATGGTACAG	S	A	G	K	K
G2313u2	WIAF-12767	J05200		RYR1, ryanodine receptor 1 (skeletal)	3048	CGGCGCAGAC [A/G] ACACTGTGTGG	S	A	G	T	T
G2313u3	WIAF-12768	J05200		RYR1, ryanodine receptor 1 (skeletal)	3084	ATGGGCACAA [C/T] GTGTGGGCCC	S	C	T	N	N
G2313u4	WIAF-12777	J05200		RYR1, ryanodine receptor 1 (skeletal)	5667	GCATCTTTGG [C/T] GATCAGGATG	S	C	T	G	G
G2313u5	WIAF-12780	J05200		RYR1, ryanodine receptor 1 (skeletal)	6600	GCTCGCTGCT [C/T] ATCGTGCAGA	S	C	T	L	L
G2313u6	WIAF-12781	J05200		RYR1, ryanodine receptor 1 (skeletal)	7191	AGCCTGAGTG [C/T] TTCGGACCCG	S	C	T	C	C
G2313u7	WIAF-12782	J05200		RYR1, ryanodine receptor 1 (skeletal)	7602	ACCACAAGGC [G/A] TCCATGGTGC	S	G	A	A	A
G2313u8	WIAF-12784	J05200		RYR1, ryanodine receptor 1 (skeletal)	9288	CAGACGCCCC [A/G] GCTGTGGTCA	S	A	G	P	P
G2313u9	WIAF-12786	J05200		RYR1, ryanodine receptor 1 (skeletal)	13690	TCCAAGAAG [G/A] AGGAAGCTGG	M	G	A	E	K
G2313u10	WIAF-12789	J05200		RYR1, ryanodine receptor 1 (skeletal)	3147	ACATCCGAGC [G/A] CGCGAAACC	S	G	A	A	A
G2314u1	WIAF-12771	J05272		IMPDH1, IMP (inosine monophosphate) dehydrogenase 1	1920	TGAAGATCGC [A/G] CAGGTGTCT	S	A	G	A	A
G2319u1	WIAF-12814	K03191		CYP1A1, cytochrome P450, subfamily I (aromatic compound-inducible), polypeptide 1	651	CCCTACAGG [T/C] ATGTGGTGGT	M	T	C	Y	H
G232u1	WIAF-11657	U58917		Homo sapiens IL-17 receptor mRNA, complete cds.	1490	TGAACATGAT [C/T] CTCGGGACT	S	C	T	I	I
G232u2	WIAF-11677	U58917		Homo sapiens IL-17 receptor mRNA, complete cds.	1293	GCAGGCCATC [T/C] CGGAGGCAGG	M	T	C	S	P
G232u3	WIAF-11658	U58917		Homo sapiens IL-17 receptor mRNA, complete cds.	1132	GGCTGCGCTG [C/T] GGCTGACCTG	M	C	T	A	V
G232u4	WIAF-11679	U58917		Homo sapiens IL-17 receptor mRNA, complete cds.	905	GCAGTGCCT [C/T] AATGACTGCC	S	C	T	L	L

[illegible]

G235u1	WIAF-11678	U83171		SCYA22, small inducible cytokine subfamily A (Cys-Cys), member 22	CAGGCCCTTA [C/T] GCGCACAACA	S	C	T	Y	Y
G2363a1	WIAF-10519	M37435	596	CSF1, colony stimulating factor 1 (macrophage)	GACAAAGGACT [G/T] GAATATTTTC	M	G	T	W	L
G2363a2	WIAF-13225	M37435	498	CSF1, colony stimulating factor 1 (macrophage)	AAGAGCATGA [C/T] AAGGCTGGG	S	C	T	D	D
G2363a3	WIAF-13226	M37435	712	CSF1, colony stimulating factor 1 (macrophage)	CAGTGACCCG [G/T] CTTCTGTCTC	M	G	T	A	S
G2369u1	WIAF-12854	M30773	857	PPP3R1, protein phosphatase 3 (formerly 2B), regulatory subunit B (19kD), alpha isoform (calcieneurin B, type I)	TTGATTGGA [C/T] AATTCTGGTT	S	C	T	D	D
G2369u2	WIAF-12855	M30773	1274	PPP3R1, protein phosphatase 3 (formerly 2B), regulatory subunit B (19kD), alpha isoform (calcieneurin B, type I)	ATGTGTGACT [C/T] TTATCAGAGA	-	C	T	-	-
G237u1	WIAF-11662	U86358	311	SCYA25, small inducible cytokine subfamily A (Cys-Cys), member 25	CACCACAACA [T/C] GCGACCTTC	M	T	C	M	T
G237u2	WIAF-11680	U86358	134	SCYA25, small inducible cytokine subfamily A (Cys-Cys), member 25	GTGCTCCGG [G/A] CGCTGGACT	M	G	A	R	H
G237u3	WIAF-11681	U86358	133	SCYA25, small inducible cytokine subfamily A (Cys-Cys), member 25	TGTGCTCCGG [C/T] GCGCTGGAC	M	C	T	R	C
G237u5	WIAF-11661	U86358	302	SCYA25, small inducible cytokine subfamily A (Cys-Cys), member 25	GCAAGGCTCC [A/G] CCAACAATG	M	A	G	H	R
G237u6	WIAF-11663	U86358	378	SCYA25, small inducible cytokine subfamily A (Cys-Cys), member 25	AGTTATCATC [A/G] TCCAAGTTTA	S	A	G	S	S
G2373u1	WIAF-12870	M26035	500	BZRP, benzodiazapine receptor (peripheral)	GCTGGCCTTC [G/A] CGACCACT	M	G	A	A	T
G2376u1	WIAF-13025	M57414	979	TACR2, tachykinin receptor 2	CTGTGTGCCA [T/C] GGGTCACACC	M	T	C	W	R
G238u1	WIAF-10177	X01394	239	TNF, tumor necrosis factor (TNF superfamily, member 2)	GCTCCAGGG [G/T] TGCTTGTTCC	S	G	T	R	R

G2381u1	WIAF-12894	M59941	730	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	CAGAGTTTG [C/T] TGGGACTCCC	S	C	T	C	C
G2381u2	WIAF-12896	M59941	1306	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	GGATCTGGAG [C/T] GAGTGGAGTG	S	C	T	S	S
G2381u3	WIAF-12900	M59941	1972	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	CGATGGGACC [G/A] GGACAGGCCG	S	G	A	P	P
G2381u4	WIAF-12901	M59941	1982	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	GGGACAGGCC [G/A] TGAAGTGGGA	M	G	A	V	M
G2381u5	WIAF-12942	M59941	773	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	CCAGAACCTG [G/C] AGTGTCTTCTT	M	G	C	E	Q
G2381u6	WIAF-12946	M59941	2458	CSF2RB, colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage)	CCCCACAGCC [C/A] GAGGGCCTCC	S	C	A	P	P
G2384u1	WIAF-12908	M61831	1000	AHCY, S-adenosylhomocysteine hydrolase	GCCGTGGAGA [A/C] GGTGAACATC	M	A	C	K	T
G2387u1	WIAF-12910	M63967	2585	ALDH5, aldehyde dehydrogenase 5	CTGCTGRAAC [T/G] CCTGGCAGAC	M	T	G	L	R
G2387u2	WIAF-12911	M63967	2996	ALDH5, aldehyde dehydrogenase 5	TATGGCCCAA [C/G] AGCAGGTGGG	M	C	G	T	R
G2387u3	WIAF-12954	M63967	2522	ALDH5, aldehyde dehydrogenase 5	GCCCGGGAAG [C/T] CTTCGGCCTG	M	C	T	A	V
G2387u4	WIAF-12955	M63967	2448	ALDH5, aldehyde dehydrogenase 5	ACCTACCAC [C/T] GGGGAGGTCA	S	C	T	T	T

G2387u5	WIAF-12956	M63967	2460	ALDH5, aldehyde dehydrogenase 5	GGGAGTTCAT [C/T] GGGCACGTGG	S	C	T	I	I
G2387u6	WIAF-12957	M63967	2991	ALDH5, aldehyde dehydrogenase 5	CGGGTATGG [C/T] CCAACAGCAG	S	C	T	G	G
G2387u7	WIAF-12958	M63967	3022	ALDH5, aldehyde dehydrogenase 5	CGCCACAGCAC [A/G] TGGATGTTGA	M	A	G	M	V
G2387u8	WIAF-12959	M63967	2943	ALDH5, aldehyde dehydrogenase 5	CCCTCATCAA [G/C] GAGGCAGGCT	M	G	C	K	N
G2388u1	WIAF-12888	M64590	588	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	TGCCACAGAC [G/A] ATTTTGGCGA	S	G	A	T	T
G2388u2	WIAF-12889	M64590	651	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	ACCAGCCTGA [G/A] GTGTCTCAGG	S	G	A	E	E
G2388u3	WIAF-12890	M64590	698	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	CAGACCATGG [T/C] GTGTGACATC	M	T	C	V	A
G2388u4	WIAF-12891	M64590	557	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	TATATTGCCA [T/C] GGGCTATTAT	M	T	C	M	T
G2388u5	WIAF-12938	M64590	587	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	GTGCCACAGA [C/G] GATTTTGGCG	M	C	G	T	R
G2388u6	WIAF-12939	M64590	518	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	CTGCATGCCA [T/C] TTCACAGCAA	M	T	C	I	T

002060" 22443960

G2388u7	WIAF-12940	M64590	810	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	GGAAATTCT [C/T] GTTGATCCCC	S	C	T	L	L
G2388u8	WIAF-12941	M64590	1481	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	CATTGTGGCT [G/A] CTCAGTGAAG	M	G	A	C	Y
G2388u9	WIAF-12947	M64590	1841	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	AAACTGAACA [G/A] TTCGTCTGAA	M	G	A	S	N
G2388u10	WIAF-12948	M64590	2325	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	GACAGGTCTA [C/T] CTAGACGGGG	S	C	T	Y	Y
G2388u11	WIAF-12949	M64590	2362	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	GTTGGGAATC [T/A] GTCGCCCTGG	M	T	A	C	S
G2388u12	WIAF-12950	M64590	3220	GLDC, glycine dehydrogenase (decarboxylating; glycine decarboxylase, glycine cleavage system protein P)	TTAGTCTCT [C/G] TCCCTAAGTT	-	C	G	-	-
G2391u1	WIAF-12998	M69238	623	ARNT, aryl hydrocarbon receptor nuclear translocator	TGGTGTATGT [G/C] TCTACTCCG	S	G	C	V	V
G2391u2	WIAF-13002	M69238	1072	ARNT, aryl hydrocarbon receptor nuclear translocator	TGCTAGTGG [C/T] CATTGGCAGA	M	C	T	A	V
G2391u3	WIAF-13021	M69238	966	ARNT, aryl hydrocarbon receptor nuclear translocator	ACCTCACTTC [G/A] TGGTGGTCCA	M	G	A	V	M
G2394u1	WIAF-13003	M73747	2061	TSHR, thyroid stimulating hormone receptor	TTGCTGGTAC [T/A] CTTCTATCCA	M	T	A	L	H

G2394u2	WIAF-13004	M73747	2248	TSHR, thyroid stimulating hormone receptor	TTACCCACGA [C/G] ATGAGGCAGG	M	C	G	D	E
G2396u1	WIAF-12995	M74542	1027	ALDH3, aldehyde dehydrogenase 3	CCCCAGTCC [C/G] CGGTGATGCA	M	C	G	P	A
G2396u2	WIAF-13019	M74542	1295	ALDH3, aldehyde dehydrogenase 3	GGCAAGAAGA [G/A] CTTGAGACT	M	G	A	S	N
G2403u1	WIAF-13583	M83670	280	CA4, carbonic anhydrase IV	TACGATAAGA [A/T] GCAAACGTGG	M	A	T	K	M
G2409u1	WIAF-10010	HT2156	1268	AGTR1, angiotensin receptor 1	CCACTCAAAC [C/T] TTTCACAAA	M	C	T	L	F
G2411u1	WIAF-13541	M97759	210	ADORA2B, adenosine A2b receptor	TGGCGGGCAA [C/T] GTGCTGTGT	S	C	T	N	N
G2422u1	WIAF-14077	S90469	375	POR, P450 (cytochrome) oxidoreductase	GCAGCTGCC [A/G] GAGATCGACA	S	A	G	P	P
G2422u2	WIAF-14078	S90469	852	POR, P450 (cytochrome) oxidoreductase	TCTGTGCTGC [A/G] GTCACCACCA	S	A	G	A	A
G2422u3	WIAF-14082	S90469	1496	POR, P450 (cytochrome) oxidoreductase	AAGGAGCCTG [T/C] CGGGGAGAAC	M	T	C	V	A
G2422u4	WIAF-14099	S90469	1443	POR, P450 (cytochrome) oxidoreductase	AGACCAAGGC [C/T] GGCCGCAITCA	S	C	T	A	A
G2422u5	WIAF-14100	S90469	1704	POR, P450 (cytochrome) oxidoreductase	GCCGCCGCTC [G/A] CATGAGGACT	S	G	A	S	S
G2427u1	WIAF-14079	U07919	1369	ALDH6, aldehyde dehydrogenase 6	ACTATGGACT [C/T] ACAGCAGCGG	S	C	T	L	L
G2427u2	WIAF-14096	U07919	1347	ALDH6, aldehyde dehydrogenase 6	ATAAAAGAG [C/T] GAATAGCACC	M	C	T	A	V
G243u1	WIAF-11684	X57522	926	TAP1, transporter 1, ABC (ATP binding cassette)	ATAGCCAGTG [C/G] AGTGTGGAG	M	C	G	A	G
G243u2	WIAF-11685	X57522	627	TAP1, transporter 1, ABC (ATP binding cassette)	ACCCTACCGC [C/T] TTCGTTGCA	S	C	T	A	A
G243u3	WIAF-11686	X57522	538	TAP1, transporter 1, ABC (ATP binding cassette)	CCTGCCGGGA [C/G] TTGCCTTGT	M	C	G	L	V
G243u4	WIAF-11687	X57522	798	TAP1, transporter 1, ABC (ATP binding cassette)	TGTTGGTCTC [C/G] TCCCTCTTGG	S	C	G	L	L
G243u5	WIAF-11689	X57522	1465	TAP1, transporter 1, ABC (ATP binding cassette)	TAGTATTTCG [G/T] GTATGCTGCT	M	G	T	G	C
G243u6	WIAF-11690	X57522	177	TAP1, transporter 1, ABC (ATP binding cassette)	AGATCCCGAG [A/G] CCGGCGGGG	S	A	G	R	R
G243u7	WIAF-11693	X57522	1067	TAP1, transporter 1, ABC (ATP binding cassette)	AACATCATGT [C/T] TCGGGTAACA	M	C	T	S	F
G243u8	WIAF-11665	X57522	1207	TAP1, transporter 1, ABC (ATP binding cassette)	GGTCACCCTG [A/G] TCACCTGGC	M	A	G	I	V

G2445u5	WIAF-14242	X95520		CNP, 2',3'-cyclic nucleotide 3' phosphodiesterase 1057	TGGAGTTGAT [C/T] TTTCAGTGCT	? C T ? ?
G2445u6	WIAF-14243	X95520		CNP, 2',3'-cyclic nucleotide 3' phosphodiesterase 1583	TCTACTGGCT [C/G] TCTAACTAAT	? C G ? ?
G2448u1	WIAF-13973	U46689		ALDH10, aldehyde dehydrogenase 10 1895 (Fatty aldehyde dehydrogenase)	TTGTCAAGGC [A/T] GAATAATTACT	S A T A A
G2457u1	WIAF-13898	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 1304 2A	GGTCCCGATG [C/T] ACACCTTGCA	M C T H Y
G2457u2	WIAF-13899	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 1934 2A	AAGAAGTAAT [G/T] GCACCGTCTC	M G T G C
G2457u3	WIAF-13900	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 2230 2A	TCGCTGTCTAT [A/G] TTCCTGGCTA	M A G I M
G2457u4	WIAF-13902	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 2916 2A	GGCATCTACA [G/A] CTGCATTCTAT	M G A S N
G2457u5	WIAF-13903	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 3251 2A	CTATGTATTTC [C/T] AGGGAACAACA	N C T Q *
G2457u6	WIAF-13917	U90277		GRIN2A, glutamate receptor, ionotropic, N-methyl D-aspartate 2756 2A	GGACATTGAC [A/G] ACAITGGCGGG	M A G N D
G2468u1	WIAF-13642	X04011		CYBB, cytochrome b-245, beta polypeptide (chronic granulomatous disease) 1017	AGGTGTCCAA [G/A] CTGGAGTGGC	S G A K K
G2473u1	WIAF-13670	X06990		ICAM1, intercellular adhesion molecule 1 (CD54), human rhinovirus receptor 1417	GGTCACCCGC [G/A] AGGTGACCGT	M G A E K
G2473u2	WIAF-13695	X06990		ICAM1, intercellular adhesion molecule 1 (CD54), human rhinovirus receptor 179	GACCAGCCCCA [A/T] GTTGTGTGGGC	M A T K M
G2480u1	WIAF-14148	X55330		AGA, aspartylglucosaminidase 800	TTGGCATGGT [T/G] GTAATCCATA	S T G V V
G2480u2	WIAF-14149	X55330		AGA, aspartylglucosaminidase 852	AAATGGTATA [A/T] AATTCAAAAAT	N A T K *

G2480u3	WIAF-14158	X55330	616	AGA, aspartylglucosaminidase	TTATCTACCA [G/C] TGCTTCTCAA	M	G	C	S	T
G2485u1	WIAF-13612	X59543	2301	RRM1, ribonucleotide reductase M1 polypeptide	ATTGATCAAA [G/A] CCAATCTTTG	M	G	A	S	N
G2485u2	WIAF-13613	X59543	2410	RRM1, ribonucleotide reductase M1 polypeptide	ATTTAAGGAC [G/A] AGACCCAGCAG	S	G	A	T	T
G2485u3	WIAF-13651	X59543	548	RRM1, ribonucleotide reductase M1 polypeptide	CAAGTCAACA [T/C] TGGATATTGT	S	T	C	L	L
G2485u4	WIAF-13652	X59543	199	RRM1, ribonucleotide reductase M1 polypeptide	TGCATGTGAT [C/T] AAGCGAGATG	S	C	T	I	I
G2485u5	WIAF-13653	X59543	1037	RRM1, ribonucleotide reductase M1 polypeptide	CAACACAGCT [C/A] GATATGTGGA	S	C	A	R	R
G2485u6	WIAF-13660	X59543	1955	RRM1, ribonucleotide reductase M1 polypeptide	GAAGATTGCA [A/C] AGTATGGTAT	M	A	C	K	Q
G2485u7	WIAF-13877	X59543	860	RRM1, ribonucleotide reductase M1 polypeptide	GAGTATGAAA [G/C] ATGACAGCAT	M	G	C	D	H
G2486u1	WIAF-14075	X59618	543	RRM2, ribonucleotide reductase M2 polypeptide	TCAGCACTGG [G/C] AATCCCTGAA	M	G	C	E	Q
G2486u2	WIAF-14076	X59618	189	RRM2, ribonucleotide reductase M2 polypeptide	TCGCTGCGCC [T/G] CCACTATGCT	-	T	G	-	-
G2486u3	WIAF-14092	X59618	524	RRM2, ribonucleotide reductase M2 polypeptide	TTGACCTCTC [C/G] AAGGACATTC	S	C	G	S	S
G2488u1	WIAF-13585	X63563	1633	POLR2B, polymerase (RNA) II (DNA directed) polypeptide B (140kD)	CCTTGATGGC [G/A] TATATTTTCA	S	G	A	A	A
G2488u2	WIAF-13586	X63563	2452	POLR2B, polymerase (RNA) II (DNA directed) polypeptide B (140kD)	CTGTAGACCG [C/T] GGCTTCTTCA	S	C	T	R	R
G2488u3	WIAF-13587	X63563	2740	POLR2B, polymerase (RNA) II (DNA directed) polypeptide B (140kD)	TCAGAACTAG [T/C] GAGACGGGCA	S	T	C	S	S
G2488u4	WIAF-13602	X63563	1411	POLR2B, polymerase (RNA) II (DNA directed) polypeptide B (140kD)	GGGTGATCA [A/G] AAGAAAGCTC	S	A	G	Q	Q
G2488u5	WIAF-13603	X63563	2386	POLR2B, polymerase (RNA) II (DNA directed) polypeptide B (140kD)	CAATTGTGGC [C/T] ATTGCATCAT	S	C	T	A	A
G2489u1	WIAF-14181	X63564	1346	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	TGGTGGACAA [T/C] GAGCTGCCTG	S	T	C	N	N

G2489u2	WIAF-14236	X63564		1847	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	TGAATCTTAG [C/T]GTGACAACCTC	?	C	T	?	?
G2489u3	WIAF-14237	X63564		2678	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	CTGAATACAA [C/T]AATTCAAGT	?	C	T	?	?
G2489u4	WIAF-14238	X63564		3059	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	AGCTGGGCTA [C/T]GGCAAGACG	?	C	T	?	?
G2489u5	WIAF-14239	X63564		3827	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	TGGGCCAGTC [C/T]GCTCGAGATG	?	C	T	?	?
G2489u6	WIAF-14240	X63564		3992	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	TGCCTGACTT [T/C]GATGTGGCCC	?	T	C	?	?
G2489u7	WIAF-14245	X63564		3938	POLR2A, polymerase (RNA) II (DNA directed) polypeptide A (220kD)	CCGAGAGCAC [G/A]GTGTGGCAG	?	G	A	?	?
G250u1	WIAF-11696	HT0155		1113	IL3RA, interleukin 3 receptor, alpha (low affinity)	CTGTGCTTTC [G/C]TGATCTGCAG	M	G	C	V	L
G251u1	WIAF-11666	HT0240		179	interleukin 1 beta convertase	TGGATAAGAC [C/T]CGAGCTTTGA	S	C	T	T	T
G251u2	WIAF-11694	HT0240		973	interleukin 1 beta convertase	GATGCTATTA [A/G]GAAAGCCAC	M	A	G	K	R
G251u3	WIAF-11695	HT0240		783	interleukin 1 beta convertase	CCCAGATATA [C/T]TACAACTCAA	S	C	T	L	L
G2513u1	WIAF-13736	HT27365		1721	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AACATCTCTAT [G/A]AAAAGCCAAA	M	G	A	M	I
G2513u2	WIAF-13737	HT27365		1741	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AACATATTGGG [A/T]AATGTGTTCA	M	A	T	E	V
G2513u3	WIAF-13738	HT27365		1697	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AATCTGTTCA [A/G]TACAGGGATT	S	A	G	Q	Q

002060-2416960

G2513u4	WIAF-13739	HT27365	1908	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	CTGTCAGATT [G/A]TAGCAATGAA	M	G	A	V	I
G2513u5	WIAF-13740	HT27365	2172	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	TATAGAGATA [C/T]ACGGAATTC	M	C	T	H	Y
G2513u6	WIAF-13744	HT27365	3019	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	TTGAGGGCC [A/G]AGGAGATCTG	M	A	G	Q	R
G2513u7	WIAF-13745	HT27365	3024	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	GGCCCAAGGA [G/A]ATCTGTTGAA	M	G	A	D	N
G2513u8	WIAF-13771	HT27365	1079	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	ACATTTTGA [T/C]CCTGAGCAAA	S	T	C	D	D
G2513u9	WIAF-13772	HT27365	1546	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AAGTTGCCTT [C/T]TGATCCAGAT	M	C	T	S	F
G2513u10	WIAF-13773	HT27365	1514	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AATTA AAAAG [A/T]ATGATCATTTG	M	A	T	R	S
G2513u11	WIAF-13774	HT27365	1445	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	AGGCTTTGG [C/T]AATAAACTCT	S	C	T	G	G
G2513u12	WIAF-13778	HT27365	2087	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	TTCAATATCAA [G/A]ATCATCAGTG	S	G	A	K	K
G2513u13	WIAF-13779	HT27365	2367	PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	TGAATGTTTG [C/T]AGCCTGGATA	N	C	T	Q	*

G2513u4 2446560

G2513u14	WIAF-13782	HT27365		PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	2719	CTCATCACCA [G/A] TGACAATACT	M	G	A	S	N
G2513u15	WIAF-13783	HT27365		PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	2567	TTGATGACAT [C/T] TTTAAATAG	S	C	T	I	I
G2513u16	WIAF-13784	HT27365		PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	2864	TAGAAATGGC [G/A] GACACAGTCC	S	G	A	A	A
G2513u17	WIAF-13785	HT27365		PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	2571	TGACATCTTT [A/T] AAATAGCGGT	N	A	T	K	*
G2513u18	WIAF-13786	HT27365		PLCB3, phospholipase C, beta 3 (phosphatidylinositol-specific)	2706	TCGTGTCATCT [C/T] GGCTCATCAC	M	C	T	R	W
G252u1	WIAF-10195	HT0425		FCER2, Fc fragment of IgE, low affinity II, receptor for (CD23A)	397	GAGGGCTGCC [C/T] GGAACGTCTC	M	C	T	R	W
G252u2	WIAF-10206	HT0425		FCER2, Fc fragment of IgE, low affinity II, receptor for (CD23A)	930	ATGGGAGGCA [T/C] GTGGACTACA	S	T	C	H	H
G253u1	WIAF-10175	HT0573		IFNB1, interferon, beta 1, fibroblast	228	GGCTTGAATA [C/T] TGCCCTCAAGG	S	C	T	Y	Y
G254u1	WIAF-10196	HT0611		IL4R, interleukin 4 receptor	466	TCAGTGGGA [T/C] AACTATACAC	S	T	C	D	D
G254u2	WIAF-10198	HT0611		IL4R, interleukin 4 receptor	1474	CATGCCCTTCT [T/C] CCACCTTCGG	S	T	C	L	L
G254u3	WIAF-10207	HT0611		IL4R, interleukin 4 receptor	1902	AGTGGCTATC [A/G] GGAGTTTGTA	M	A	G	Q	R
G260u1	WIAF-10186	HT1090		IL1R1, interleukin 1 receptor, type I	453	TGTTATAATG [C/G] ACAAGCCATA	M	C	G	A	G
G261u1	WIAF-10187	HT1101		IL7R, interleukin 7 receptor	434	CCTGAGTGTG [A/G] TCTATCGGA	M	A	G	I	V
G261u2	WIAF-10203	HT1101		IL7R, interleukin 7 receptor	517	TTTTAATGCA [T/C] GATGTAGCTT	S	T	C	H	H
G267u1	WIAF-11735	HT1877		IL2RB, interleukin 2 receptor, beta	881	TCCTCGTGGG [C/T] CTCAGCGGGG	S	C	T	G	G

G2513u14" 24.4.2000

G267u2	WIAF-11759	HT1877	IL2RB, interleukin 2 receptor, beta	379	AGTCAAGCAT [C/T] CTGGGCTGC	M	C	T	S	F
G268u1	WIAF-11758	HT1985	CD19 antigen	568	GCCTCCGTGT [G/C] TCCACCGAG	M	G	C	V	L
G268u2	WIAF-11734	HT1985	CD19 antigen	783	ACGATCGCCC [G/T] GCAGAGATA	S	G	T	P	P
G270u1	WIAF-11736	HT2415	IL6R, interleukin 6 receptor	530	AGGAGGTGGC [A/G] AGAGGCGTGC	S	A	G	A	A
G270u2	WIAF-11760	HT2415	IL6R, interleukin 6 receptor	1590	CATTGCCATT [G/A] TTCTGAGTT	M	G	A	V	I
G270u3	WIAF-11737	HT2415	IL6R, interleukin 6 receptor	1510	CCAGTCAAG [A/C] TTCTTCTTCA	M	A	C	D	A
G270u4	WIAF-11761	HT2415	IL6R, interleukin 6 receptor	1451	CTACTAATAA [A/T] GACGATGATA	M	A	T	K	N
G270u5	WIAF-11766	HT2415	IL6R, interleukin 6 receptor	1843	TTCCCCAGAT [A/G] GCTGGCTGGG	N	A	G	*	W
G270u6	WIAF-11767	HT2415	IL6R, interleukin 6 receptor	1829	ATACAGACTA [C/T] TTCTTCCCCA	S	C	T	Y	Y
G271u1	WIAF-11762	HT2531	CD2, CD2 antigen (p50), sheep red blood cell receptor	577	TCAGAGGGTC [A/G] TCACACACAA	M	A	G	I	V
G271u2	WIAF-11739	HT2531	CD2, CD2 antigen (p50), sheep red blood cell receptor	861	GGAGCCCCA [A/C] CAAATTCCAG	M	A	C	X	H
G271u3	WIAF-11768	HT2531	CD2, CD2 antigen (p50), sheep red blood cell receptor	818	CTGGAGACAA [G/A] AGCCACAGA	M	G	A	R	K
G271u4	WIAF-11738	HT2531	CD2, CD2 antigen (p50), sheep red blood cell receptor	736	CCTCTTGATG [G/A] TCTTTGTGGC	M	G	A	V	I
G273u1	WIAF-11763	HT3139	IL2RA, interleukin 2 receptor, alpha	667	ATCATGGTGC [C/T] TGGCTGCCAG	M	C	T	P	L
G273u2	WIAF-11764	HT3139	IL2RA, interleukin 2 receptor, alpha	956	AAAGTCCAAT [G/C] CAGCCAGTGG	M	G	C	M	I
G273u3	WIAF-11765	HT3139	IL2RA, interleukin 2 receptor, alpha	701	ACGATGACCC [G/A] CCAGAGATCC	S	G	A	P	P
G273u4	WIAF-11740	HT3139	IL2RA, interleukin 2 receptor, alpha	1133	AAATGACCCA [C/T] GGAAGACAA	S	C	T	H	H
G273u5	WIAF-11769	HT3139	IL2RA, interleukin 2 receptor, alpha	1163	AGCCCCAGCT [C/A] ATATGCACAG	S	C	A	L	L
G276u1	WIAF-10192	HT3670	CD4 antigen	644	CTGGTAGTAG [C/G] CCTCAGTGC	M	C	G	S	R
G276u2	WIAF-10193	HT3670	CD4 antigen	1535	CCTGCCAGTG [T/C] CCTCACGGT	S	T	C	C	C
G276u3	WIAF-10205	HT3670	CD4 antigen	1217	TGATGCTGAG [T/C] TTGAAACTGG	S	T	C	S	S

G277u1	WIAF-10007	D10232	851	RENBP, renin-binding protein	CACGTGATTG [A/G] CAAGTTCCTA	M	A	G	D	G
G277u2	WIAF-10032	D10232	842	RENBP, renin-binding protein	CTTCGAGCCC [A/G] CGTGATTGAC	M	A	G	H	R
G277u3	WIAF-10042	D10232	634	RENBP, renin-binding protein	GCTGGCGGGC [A/G] AATACGCAGA	M	A	G	K	E
G279u1	WIAF-10047	K01740	1658	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	ACTGATGTCC [G/A] TCCTTTGTAT	M	G	A	R	H
G279u2	WIAF-10049	K01740	2328	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	CCTTACTGAA [G/A] GTTCTAGTT	S	G	A	K	K
G279u3	WIAF-10050	K01740	4650	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	CTGTTCTCCC [G/A] AAACCAGACT	S	G	A	P	P
G279u4	WIAF-10061	K01740	6919	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	CCAGAGACAC [A/G] TGAAGTCAC	M	A	G	M	V
G279u5	WIAF-10080	K01740	480	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	TTAAGAACAT [G/A] GCTTCCCATC	M	G	A	M	I
G279u6	WIAF-10082	K01740	2129	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	TACATTCTAA [G/A] CATTGGAGCA	M	G	A	S	N
G279u7	WIAF-10084	K01740	2533	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	GTTTGCACAC [A/G] GAACACCTAT	M	A	G	R	G
G279u8	WIAF-10086	K01740	6639	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	ACCTCCAAT [T/C] ATTGCTCGAT	S	T	C	I	I
G279u9	WIAF-10087	K01740	5957	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	GAGAAATTATC [G/A] CTTCATGCA	M	G	A	R	H
G279a10	WIAF-10495	K01740	5829	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	TGACAGTACA [G/A] GAATTGCTC	S	G	A	Q	Q
G279a11	WIAF-10496	K01740	5852	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	TTTTTCACCA [T/G] CTTTGATGAG	M	T	G	I	S
G279a12	WIAF-10502	K01740	2492	F8C, coagulation factor VIIIC, procoagulant component (hemophilia)	ACCACAATTC [C/T] AGAAATGAC	M	C	T	P	L

G279a13	WIAF-10503	K01740	F8C, coagulation factor VIIIc, procoagulant component (hemophilia 6906 A)	TGCAAGTGA [C/T] TTCCAGAAGA	S	C	T	D	D
G279a14	WIAF-13120	K01740	F8C, coagulation factor VIIIc, procoagulant component (hemophilia 1980 A)	CAGAGAATAT [A/C] CAACGCTTTC	S	A	C	I	I
G279a15	WIAF-13121	K01740	F8C, coagulation factor VIIIc, procoagulant component (hemophilia 1982 A)	GAGAATATAC [A/C] ACCTTTTCTC	M	A	C	Q	P
G282u1	WIAF-10067	L25615	AVPR1A, arginine vasopressin receptor 1A	CGCTTTCTT [C/A] ATCATCCAGA	M	C	A	F	L
G282u2	WIAF-10070	L25615	AVPR1A, arginine vasopressin receptor 1A	TCGGCATGTT [T/C] GCGTCGGCCT	S	T	C	F	F
G282u3	WIAF-10071	L25615	AVPR1A, arginine vasopressin receptor 1A	GCCTGGCCGA [C/T] CTGGCCGTGG	S	C	T	D	D
G282u4	WIAF-10072	L25615	AVPR1A, arginine vasopressin receptor 1A	TCTCTCGCC [G/A] GTCCCGACGC	M	G	A	G	S
G282u5	WIAF-10073	L25615	AVPR1A, arginine vasopressin receptor 1A	AGACTCTGCA [A/G] CAGCCCGCGC	S	A	G	Q	Q
G282u6	WIAF-10092	L25615	AVPR1A, arginine vasopressin receptor 1A	CCTTGAATAG [C/A] TGCTGTATTC	M	C	A	S	R
G282a7	WIAF-10499	L25615	AVPR1A, arginine vasopressin receptor 1A	TGTAATCCT [G/A] GATATACATG	N	G	A	W	*
G284u1	WIAF-10182	M16827	ACADM, acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain	AATATCCTGT [A/G] GAAGAACTAA	S	A	G	V	V
G284a2	WIAF-10515	M16827	ACADM, acyl-Coenzyme A dehydrogenase, C-4 to C-12 straight chain	TTGTGGAAGC [A/G] GATACCCAG	S	A	G	A	A
G285u1	WIAF-10108	M28372	ZNF9, zinc finger protein 9 (a cellular retroviral nucleic acid binding protein)	CTCTTCCAGA [T/C] ATTGTATC	S	T	C	D	D
G289u1	WIAF-10041	M63012	172 PON1, paraoxonase 1	CTCTGAAGAC [A/T] TGGAGATACT	M	A	T	M	L
G290u1	WIAF-10085	M63959	LRPAP1, low density lipoprotein-related protein-associated protein 1 (alpha-2-macroglobulin receptor-associated protein 1)	CTCATACGCA [A/G] CCTCAATGTC	M	A	G	N	S

G290a2	WIAF-13122	M63959			LRPAP1, low density lipoprotein-related protein-associated protein 1 (alpha-2-macroglobulin receptor-associated protein 1)	223	ACACTGCA [T/A] CTTCCTCCG	M	T	A	H	Q
G292u1	WIAF-10180	M74096			ACADL, acyl-Coenzyme A dehydrogenase, long chain	1002	AGTCAACAT [A/C] AATAGCAGA	M	A	C	K	Q
G293u1	WIAF-10068	M74775			LIPA, lipase A, lysosomal acid, cholesterol esterase (Wolman disease)	723	AAGACTTAT [T/C] TGGAGACAA	M	T	C	F	S
G293a2	WIAF-10497	M74775			LIPA, lipase A, lysosomal acid, cholesterol esterase (Wolman disease)	107	TCAGGGTCT [G/A] GAGGAAACT	M	G	A	G	R
G293a3	WIAF-10498	M74775			LIPA, lipase A, lysosomal acid, cholesterol esterase (Wolman disease)	86	GGTCTCTGG [C/A] CCCTGCATTC	M	C	A	P	T
G295u1	WIAF-10057	U04270			KCNH2, potassium voltage-gated channel, subfamily H, member 2	1282	AAAGGACGA [A/T] CCCACAATGT	M	A	T	T	S
G295u2	WIAF-10062	U04270			KCNH2, potassium voltage-gated channel, subfamily H, member 2	1875	CGCACTGGCT [A/G] GCCTGCATCT	S	A	G	L	L
G295u3	WIAF-10064	U04270			KCNH2, potassium voltage-gated channel, subfamily H, member 2	2040	ACTTCACCTT [C/T] AGCAGCCTCA	S	C	T	F	F
G295u4	WIAF-10088	U04270			KCNH2, potassium voltage-gated channel, subfamily H, member 2	1650	CCGGCCGCAT [C/T] GCCGTCCACT	S	C	T	I	I
G295u5	WIAF-10090	U04270			KCNH2, potassium voltage-gated channel, subfamily H, member 2	2139	CCCTCATGTA [T/C] GCTAGCATCT	S	T	C	Y	Y
G2951u1	WIAF-14147	HT0030			ZNF42, zinc finger protein 42 (myeloid-specific retinoic acid-responsive)	1334	CCCTGCTCTG [A/G] TCACCACCCG	M	A	G	I	V
G2951u2	WIAF-14157	HT0030			ZNF42, zinc finger protein 42 (myeloid-specific retinoic acid-responsive)	1558	ACCAGCTTAC [G/A] CACACCGAGG	S	G	A	T	T

G2959u1	WIAF-13501	HT0134		1014	GRLF1, glucocorticoid receptor DNA binding factor 1	GTGGAGAGAC [T/C] CTGCATAGCT	S	T	C	T	T
G2959u2	WIAF-13518	HT0134		1853	GRLF1, glucocorticoid receptor DNA binding factor 1	GAGCATCTT [A/C] CAGCCTGTTT	M	A	C	Y	S
G296a1	WIAF-10514	U12778		961	ACADSE, acyl-Coenzyme A dehydrogenase, short/branched chain	TATTCCATAT [A/G] TTAAGAAAG	M	A	G	I	V
G2968u1	WIAF-12699	HT0244		1754	SMARCA1, SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 1	CAGAAGAAAC [C/T] AGTACGTGTA	M	C	T	P	L
G2968u2	WIAF-12716	HT0244		2624	SMARCA1, SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 1	TGGGAACGTT [G/T] CAATGAATTA	M	G	T	C	F
G297u1	WIAF-10109	U16660		402	ECH1, enoyl Coenzyme A hydratase 1, peroxisomal	ACATGGCTTC [G/A] GACATCTCTGC	S	G	A	S	S
G297u2	WIAF-10110	U16660		149	ECH1, enoyl Coenzyme A hydratase 1, peroxisomal	GCACAGAGG [A/C] GGCTTCCGGA	M	A	C	E	A
G2970u1	WIAF-12746	HT0281		682	BR140: bromodomain-containing protein, 140KD (peregryn)	ATGACATGGA [C/T] GAGGAGGACT	S	C	T	D	D
G2975u1	WIAF-12729	HT0334		1104	B-cell-specific transcription factor	AGTTTTCGG [G/A] AGTCCCTACA	S	G	A	G	G
G2975u2	WIAF-12730	HT0334		1185	B-cell-specific transcription factor	GCTCCCTCTA [C/T] TATTATAGCG	S	C	T	Y	Y
G2976a1	WIAF-12129	HT0340		1600	SATB1, special AT-rich sequence binding protein 1 (binds to nuclear matrix/scaffold- associating DNA's)	GTCCTGCCCC [C/A] CTCATCAGCA	S	C	A	P	P
G2976u2	WIAF-12743	HT0340		2116	SATB1, special AT-rich sequence binding protein 1 (binds to nuclear matrix/scaffold- associating DNA's)	TGGCCTCTCC [A/G] GCAGAGTCAG	S	A	G	P	P

G2978u1	WIAF-12721	HT0346	1140	MSX1, msh (Drosophila) homeo box homolog 1 (formerly homeo box 7)	CATAGAGGT [C/T] CCAGGTCCCC	-	C	T	-	-
G298u1	WIAF-10048	U33837	8995	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	CCGGACAGGA [G/A] GTCATTCCC	M	G	A	R	K
G298u2	WIAF-10051	U33837	13217	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	ATGACGCCAT [C/T] GAACTGCCTA	S	C	T	I	I
G298u3	WIAF-10077	U33837	6298	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	AACTCTTTCA [T/C] TGTGTTTCA	M	T	C	I	T
G298u4	WIAF-10078	U33837	6371	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	CCATGTTGCC [G/A] GTGCAGGCC	S	G	A	P	P
G298u5	WIAF-10079	U33837	6914	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	ACTCTGAAGT [G/A] ATTGCTTATG	S	G	A	V	V
G298u6	WIAF-10081	U33837	8718	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	GTCCCAATGC [G/A] CATCTGGCG	M	G	A	A	T
G298u7	WIAF-10083	U33837	9088	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	ACTTGCTCTG [A/G] AAATGAATTC	M	A	G	E	G
G298u8	WIAF-10096	U33837	6949	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	ACTCCTTATG [G/C] CATCACTGTT	M	G	C	G	A
G298u9	WIAF-10097	U33837	7149	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	TTGCTTGGAA [A/G] ACAATGGTGG	M	A	G	N	D
G298u10	WIAF-10100	U33837	8590	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	TACACAAAT [G/A] TCATAATTCA	M	G	A	C	Y
G298u11	WIAF-10101	U33837	12948	Human glycoprotein receptor gp330 precursor, mRNA, complete cds.	CATCTTTGAA [G/C] ACAGTTATA	M	G	C	D	H
G2980u1	WIAF-12723	HT0356	437 E(spl)	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	TCATGGCCAC [G/A] GACCCCCAGT	M	G	A	G	R

G2980u2	WIAF-12726	HT0356	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	2044 E(sp1)	AGTGCTGGC [A/G] GTGGGATGG	S	A	G	A	A
G2980u3	WIAF-12747	HT0356	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	379 E(sp1)	CCATGGCAGA [G/A] TTGAATGCCA	S	G	A	E	E
G2980u4	WIAF-12748	HT0356	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	276 E(sp1)	ATCGCCAAGA [G/A] ATTGAATACG	M	G	A	R	K
G2980u5	WIAF-12749	HT0356	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	1876 E(sp1)	GCCACACAGA [C/T] GGAGCCAGCT	S	C	T	D	D
G2980u6	WIAF-12750	HT0356	TLE1, transducin-like enhancer of split 1, homolog of Drosophila	1759 E(sp1)	CGCCCTGCTA [C/T] GCCCTGGCCA	S	C	T	Y	Y
G2981u1	WIAF-12720	HT0357	TLE2, transducin-like enhancer of split 2, homolog of Drosophila	2206 E(sp1)	ACAAATACAT [T/C] GTGACAGGCT	S	T	C	I	I
G2981u2	WIAF-12737	HT0357	TLE2, transducin-like enhancer of split 2, homolog of Drosophila	1036 E(sp1)	CGGACAGCGT [C/T] GCCCTGAGGA	S	C	T	V	V
G2981u3	WIAF-12740	HT0357	TLE2, transducin-like enhancer of split 2, homolog of Drosophila	2181 E(sp1)	CTGAGTTGTG [A/T] CATCTCCAGA	M	A	T	D	V
G2983u1	WIAF-12833	HT0360	TLE3, transducin-like enhancer of split 3, homolog of Drosophila	636 E(sp1)	TGTCACCTTC [G/C] GAAAGCCTCC	S	G	C	S	S
G2983u2	WIAF-12834	HT0360	TLE3, transducin-like enhancer of split 3, homolog of Drosophila	1944 E(sp1)	TGGACAACAC [G/A] GTGCGCTCCT	S	G	A	T	T

004000 2445960

G2983u3	WIAF-12848	HT0360	1710 E (spl)	TLE3, transducin-like enhancer of split 3, homolog of Drosophila	ACCTGGCCTC [G/A] CCCACGCCCC	S	G	A	S	S
G2985u1	WIAF-12724	HT0421	995 homeotic protein D3		GGCTTGCCA [G/A] CGCCAACCTG	M	G	A	S	N
G2985u2	WIAF-12725	HT0421	1003 homeotic protein D3		CAGCGCAAC [C/T] TGCAGGGCAG	S	C	T	L	L
G2986u1	WIAF-14124	HT0468	1197 CSDA, cold shock domain protein A		GCCGTGGATA [C/T] CGCGTCCCT	S	C	T	Y	Y
G2987u1	WIAF-12758	HT0474	2068 ZNF7, zinc finger protein 7 (KOX 4, clone HF.16)		ACTGGTTTAA [C/T] GAATATGGGA	S	C	T	Y	Y
G2987u2	WIAF-12773	HT0474	985 ZNF7, zinc finger protein 7 (KOX 4, clone HF.16)		GAGAGAGGCC [G/C] TACGAATGTG	S	G	C	P	P
G2987u3	WIAF-12775	HT0474	1278 ZNF7, zinc finger protein 7 (KOX 4, clone HF.16)		AGCCAGCAGT [C/T] GCAGCTGTT	M	C	T	S	L
G3005a1	WIAF-12133	HT0735	1441 homeotic protein 5.1		GAGGACGCG [C/T] CCCGGGCGTG	S	C	T	G	G
G3008a1	WIAF-12134	HT0753	1850 ATF4, activating transcription factor 4 (tax-responsive enhancer element B67)		TAAAGAGAG [G/A] GCGGATTCCC	S	G	A	R	R
G3008u2	WIAF-12798	HT0753	946 ATF4, activating transcription factor 4 (tax-responsive enhancer element B67)		CCCTTCGACC [C/A] CTCGGGTTTG	M	C	A	P	Q
G3008u3	WIAF-12812	HT0753	1482 ATF4, activating transcription factor 4 (tax-responsive enhancer element B67)		CACCTGCTTAC [G/A] TTGCCATGAT	M	G	A	V	I
G3008u4	WIAF-12813	HT0753	1847 ATF4, activating transcription factor 4 (tax-responsive enhancer element B67)		CTCTAAAAGA [G/C] AGGGCGGATT	M	G	C	E	D
G301u1	WIAF-10127	U71285	3639 MTR, 5-methyltetrahydrofolate-homocysteine methyltransferase		TGTGGAGACT [C/T] GCAGACATCG	S	C	T	L	L
G3012u1	WIAF-12794	HT0873	402 MAD, MAX dimerization protein		TGGTCCCACT [G/T] GGACCCGAAT	S	G	T	L	L
G3014u1	WIAF-14183	HT0899	274 homeotic protein 2, distal-less		AAAAGACTCA [G/A] TACTTGGCCT	S	G	A	Q	Q

G3020u1	WIAF-12797	HT0956		MLL3, myeloid/lymphoid or mixed-lineage leukemia (trithorax (Drosophila) homolog); translocated to, 3	852		GTGCTTCAA [A/G] GAACCTTCCA	S	A	G	K	K
G3023u1	WIAF-13724	HT0966		zinc finger, X-linked, duplicated	381 A		GCTGCAGCA [G/A] CAATATGACA	S	G	A	K	K
G3023u2	WIAF-13725	HT0966		zinc finger, X-linked, duplicated	220 A		GGCCAAACTC [G/A] GCGCCCAACA	M	G	A	G	S
G3023u3	WIAF-13726	HT0966		zinc finger, X-linked, duplicated	69 A		AGTCGCACGA [T/C] AAACCTGCGC	S	T	C	D	D
G3023u4	WIAF-13727	HT0966		zinc finger, X-linked, duplicated	249 A		ACTTCGAACC [C/T] GAGAGGCCTT	S	C	T	P	P
G3023u5	WIAF-13765	HT0966		zinc finger, X-linked, duplicated	661 A		CAGGTTCTCT [G/A] CTCGCAGTAG	M	G	A	A	T
G3023u6	WIAF-13766	HT0966		zinc finger, X-linked, duplicated	1302 A		TGACTCCTTC [G/T] AGCACCCCTT	S	G	T	S	S
G3027u1	WIAF-12800	HT1035		124 HOXB7, homeo box B7	124		TTATGCGAAT [G/A] CTTTATTTTC	M	G	A	A	T
G3027u2	WIAF-12816	HT1035		450 HOXB7, homeo box B7	450		GGGACTCGGA [C/T] TTGGCGGCG	S	C	T	D	D
G3028u1	WIAF-12806	HT1037		701 homeotic protein C8	701		AGACCCCTGA [A/G] CTGGAGAAGG	S	A	G	E	E
G3029u1	WIAF-14153	HT1100		441 zinc finger protein 8	441		TCAGACTCAG [G/A] GAAACTGGG	S	G	A	R	R
G3029u2	WIAF-14155	HT1100		1416 zinc finger protein 8	1416		GGCGTGACA [A/G] TCCTCGAGCA	S	A	G	Q	Q
G303u1	WIAF-10000	X13916		LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	4110		ATGAGCTGG [G/A] GCGGACACAC	M	G	A	G	E
G303u2	WIAF-10001	X13916		LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	4012		GCGAGCTCTG [C/T] GACCACTGCT	S	C	T	C	C
G303u3	WIAF-10002	X13916		LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	4702		GCCTGCCCG [C/T] ATTAGGCAG	S	C	T	R	R
G303u4	WIAF-10003	X13916		LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	6395		CTGGATCGCA [G/A] GCAACATCTA	M	G	A	G	S
G303u5	WIAF-10004	X13916		LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	6937		AAGGCACCA [C/T] GTGTGCCGG	S	C	T	N	N

G303u6	WIAF-10005	X13916	9391	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GGCTGAGGA [T/C] GACGGCCGGA	S	T	C	D	D
G303u7	WIAF-10011	X13916	766	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACTGCATGGA [C/T] GGCTCAGATG	S	C	T	D	D
G303u8	WIAF-10015	X13916	9040	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACCCGACCTG [C/T] GGCCCCAGTG	S	C	T	C	C
G303u9	WIAF-10019	X13916	11749	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	CCCTGCGCTG [C/T] AACATGTTCTG	S	C	T	C	C
G303u10	WIAF-10020	X13916	1917	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GACCAGTATG [G/A] GAAGCCGGGT	M	G	A	G	E
G303u11	WIAF-10021	X13916	4810	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	AGAAGCGCAT [C/T] CTTTGGATTG	S	C	T	I	I
G303u12	WIAF-10022	X13916	6367	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	TTGGCCGTGT [G/C] GAGGGCATTG	S	G	C	V	V
G303u13	WIAF-10023	X13916	6247	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	CTGTCGGCAT [C/T] GACTTCCACG	S	C	T	I	I
G303u14	WIAF-10024	X13916	8371	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACGCCTCAGA [T/C] GAGATGAACCT	S	T	C	D	D
G303u15	WIAF-10030	X13916	11395	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACGGCAGCGA [C/T] GAGGAGGCT	S	C	T	D	D

G303u16	WIAF-10031	X13916	12763	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACGTCTTTGA [G/A]GATTACATCT	S	G	A	E	E
G303u17	WIAF-10035	X13916	640	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	ACGGATCTGA [C/T]GAGGCCCTTG	S	C	T	D	D
G303u18	WIAF-10037	X13916	1609	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GCCGCCTTGT [C/T]TACTGGGCAG	S	C	T	V	V
G303u19	WIAF-10038	X13916	1629	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GATGCCCTATC [T/G]GGACTATATT	M	T	G	L	R
G303u20	WIAF-10039	X13916	2210	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	CACCAGCTAC [C/T]TCATTGGCCG	M	C	T	L	F
G303u21	WIAF-10043	X13916	7287	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GATGGCTCCA [G/A]GAGGATCACC	M	G	A	R	K
G303u22	WIAF-10044	X13916	8258	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	CTCTGACGAG [A/G]TCCCTTGCAA	M	A	G	I	V
G303u23	WIAF-10045	X13916	11871	LRP1, low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor)	GTGGGCACCG [A/G]GAAAGCGGCC	M	A	G	E	G
G303u1	WIAF-14097	HT1128	611	PSMC3, proteasome (prosome, 26S subunit, ATPase, 3	TGGGGATCCA [A/G]CCTCCAAAAG	S	A	G	Q	Q
G303u1	WIAF-12836	HT1182	137	TCF12, transcription factor 12 (HTF4, helix-loop-helix transcription factors 4)	ATAAGGGAGC [G/A]TGAGGAGTCT	M	G	A	R	H

G3034u2	WIAF-12837	HT1182	TCF12, transcription factor 12 (HTF4, helix-loop-helix transcription factors 4)	421	ATCTTCAATT [A/G] TGGGTTCCCTT	M	A	G	M	V
G3038u1	WIAF-12864	HT1373	NFKB1, nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (p105)	1700	AGAGAAGGCT [A/G] TGCAGCTTGC	M	A	G	M	V
G3038u2	WIAF-12881	HT1373	NFKB1, nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (p105)	1936	TGTACCAGAC [G/A] CCCTTGCACT	S	G	A	T	T
G3038u3	WIAF-12882	HT1373	NFKB1, nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (p105)	2641	AGCTGCAGCT [G/C] TATAAGTTAC	S	G	C	L	L
G3039u1	WIAF-13027	HT1375	GLI3, GLI-Kruppel family member GLI3 (Greig cephalopolysyndactyly syndrome)	3761	AACAGCCCCG [G/T] AAGTGGCACC	M	G	T	G	V
G3039u2	WIAF-13028	HT1375	GLI3, GLI-Kruppel family member GLI3 (Greig cephalopolysyndactyly syndrome)	3963	CGCCAAATGA [G/T] TCAGCTGGCA	M	G	T	E	D
G304u1	WIAF-12242	HT637	FABP3, fatty acid binding protein 3, muscle and heart (mammary- derived growth inhibitor)	158	CTCACCTTAA [A/G] AACACACAGC	M	A	G	K	R
G3043u1	WIAF-12867	HT1486	IRF2, interferon regulatory factor 2	842	GTGCCGAGGG [G/A] CGGCCACACT	S	G	A	G	G
G3047u1	WIAF-12875	HT1518	transcription factor 1, nucleolar	1233	TCCGTTTTCCT [C/T] GAGAGCCTGC	S	C	T	L	L
G3047u2	WIAF-12876	HT1518	transcription factor 1, nucleolar	1746	GGATTAAAGAA [G/A] GCAGCCGGAAG	S	G	A	K	K
G3047u3	WIAF-12877	HT1518	transcription factor 1, nucleolar	1829	TCCAAGAAGA [T/C] GAAATTCAG	M	T	C	M	T
G3048u1	WIAF-12884	HT1530	transcription factor USF	628	AGTGGAGCGT [C/T] GCCGCCGAGA	M	C	T	R	C
G305u1	WIAF-10150	HT0034	prolyl 4-hydroxylase, beta subunit/protein disulfide isomerase/thyroid hormone-binding protein, alt. transcript 1	777	CCCTTGTCAT [C/T] GAGTTCACCG	S	C	T	I	I

G305u2	WIAF-10154	HT0034		prolyl 4-hydroxylase, beta subunit/protein disulfide isomerase/thyroid hormone-binding protein, alt. transcript 1	186		TGGGGCCCA [C/A] AAGTACTGC	M	C	A	H	Q
G305u3	WIAF-10155	HT0034		prolyl 4-hydroxylase, beta subunit/protein disulfide isomerase/thyroid hormone-binding protein, alt. transcript 1	1428		GGACGGTCAT [T/C] GATTACAACG	S	T	C	I	I
G3050u1	WIAF-12860	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	2098		AACATTGCAA [T/C] GGCATTTTGA	S	T	C	N	N
G3050u2	WIAF-12861	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	2845		TAGGCCCTTC [T/C] GGGCTTGGAC	S	T	C	S	S
G3050u3	WIAF-12862	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	3409		CCTCGTCGTC [G/A] TCCTTCAGACA	S	G	A	S	S
G3050u4	WIAF-12874	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	1699		TCTCTTCTGT [G/C] TCACACACAG	S	G	C	V	V
G3050u5	WIAF-12878	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	2093		GTAAAAACAT [T/G] GCAATGGCAT	M	T	G	C	G
G3050u6	WIAF-12879	HT1558		FSRG1: female sterile homeotic-related gene 1 (mouse homolog)	2746		CTGGGGCCGA [C/T] GAAGATGACA	S	C	T	D	D
G3051u1	WIAF-12866	HT1569		MEF2B, MADS box transcription enhancer factor 2, polypeptide B 1423 (myocyte enhancer factor 2B)	1423		CTTGGCCGAC [G/A] GCTGGCCCCG	S	G	A	T	T
G3051u2	WIAF-13022	HT1569		MEF2B, MADS box transcription enhancer factor 2, polypeptide B 661 (myocyte enhancer factor 2B)	661		CAGACTACAG [C/T] GAGCCCCACG	S	C	T	S	S
G3057a1	WIAF-12142	HT1669		alpha-fetoprotein enhancer-binding protein 5565	5565		AGACTGCTCT [T/C] GAGGCTCATA	S	T	C	L	L
G3057a2	WIAF-12143	HT1669		alpha-fetoprotein enhancer-binding protein 5634	5634		CTCTGTCTGC [G/A] ATGCTCTTAG	S	G	A	A	A

G3057a3	WIAF-12144	HT1669		5664	alpha-fetoprotein enhancer-binding protein	GCGGACTCCA [G/T] ATGAAAGGAG	M	G	T	Q	H
G3057a4	WIAF-12145	HT1669		5703	alpha-fetoprotein enhancer-binding protein	GCTTTTCCCA [C/T] CTACCCCAA	S	C	T	H	H
G3057u5	WIAF-12885	HT1669		2227	alpha-fetoprotein enhancer-binding protein	TCTGGAGATC [C/T] ATATGAGGTC	M	C	T	H	Y
G3057u6	WIAF-12892	HT1669		3720	alpha-fetoprotein enhancer-binding protein	AGACCTTGCC [G/A] GCTCAGCTAC	S	G	A	P	P
G3057u7	WIAF-12893	HT1669		4137	alpha-fetoprotein enhancer-binding protein	CAAGGTTTAC [G/A] GACTACCAGC	S	G	A	T	T
G3057u8	WIAF-12897	HT1669		4783	alpha-fetoprotein enhancer-binding protein	GAAGACCAAC [A/C] CTCCCCAGCA	M	A	C	T	P
G3057u9	WIAF-12898	HT1669		5215	alpha-fetoprotein enhancer-binding protein	TCCAACCTCC [A/C] CAATGAACAC	M	A	C	T	P
G3057u10	WIAF-12904	HT1669		7266	alpha-fetoprotein enhancer-binding protein	CCCTGCAGGC [C/T] GCGTTGACTT	S	C	T	A	A
G3057u11	WIAF-12907	HT1669		8345	alpha-fetoprotein enhancer-binding protein	CCAACAGACG [A/C] CTATTGGGAG	M	A	C	D	A
G3057u12	WIAF-12943	HT1669		4257	alpha-fetoprotein enhancer-binding protein	TGGTGTGGTT [T/C] CAGAAATGCC	S	T	C	F	F
G3057u13	WIAF-12951	HT1669		7333	alpha-fetoprotein enhancer-binding protein	ACCAGGCTTT [T/A] CTCCTTATTA	M	T	A	S	T
G3057u14	WIAF-13030	HT1669		303	alpha-fetoprotein enhancer-binding protein	GCAGCCTGTC [G/A] GAGGACGAGT	S	G	A	S	S
G3057u15	WIAF-13031	HT1669		777	alpha-fetoprotein enhancer-binding protein	GCCTTCCAGA [G/A] GAGGACGAGG	S	G	A	E	E
G306u1	WIAF-10118	HT0040		1618	CPT2, carnitine palmitoyltransferase II	CTCTACTGCC [G/A] TCCACTTTGA	M	G	A	V	I
G307u1	WIAF-10076	HT0114		110	EDN2, endothelin 2	CGTTGCGCTA [G/A] CCCTGCTCGT	M	G	A	A	T
G3070u1	WIAF-12972	HT2085		625	pre-B-cell leukemia transcription factor 3	AGAAATATGA [A/G] CAGGCAATGA	S	A	G	E	E
G3070u2	WIAF-12973	HT2085		841	pre-B-cell leukemia transcription factor 3	GTAACCTCAG [T/C] AAACAGGCCA	S	T	C	S	S
G3071u1	WIAF-12886	HT2086		995	AGER, advanced glycosylation end product-specific receptor	CCTGCAGGC [T/C] GTGATGATCC	S	T	C	A	A
G3071u2	WIAF-12887	HT2086		1475	AGER, advanced glycosylation end product-specific receptor	GAGGCCAGAT [C/G] TACAGGCCAC	M	C	G	I	M
G3071u3	WIAF-12935	HT2086		933	AGER, advanced glycosylation end product-specific receptor	ACGCATGGTG [A/G] GCATCATCCA	M	A	G	S	G

000000 2445960

G3071u4	WIAF-12936	HT2086	1052	AGER, advanced glycosylation end product-specific receptor	GTAACCTTCAG [C/T] AAACAGGCCA	S	C	T	S	S
G3071u5	WIAF-12937	HT2086	836	AGER, advanced glycosylation end product-specific receptor	ACAAGTATGA [G/A] CAGCATGTA	S	G	A	E	E
G308u1	WIAF-10094	HT0192	484	ANX4, annexin IV (placental anticoagulant protein II)	ATGGACGGAG [C/G] CTTGAAGATG	M	C	G	S	R
G308u2	WIAF-10095	HT0192	333	ANX4, annexin IV (placental anticoagulant protein II)	GGGATGATGA [C/T] GCCCACGGTG	M	C	T	T	M
G3081u1	WIAF-12997	HT2188	689	PSMC2, proteasome (prosome, 26S subunit, ATPase, 2)	GGCATTGAGC [C/T] TCCCAAGGGC	M	C	T	P	L
G3083u1	WIAF-12976	HT2228	106	IGHMBP2, immunoglobulin mu binding protein 2	TGCTGGAGCT [T/C] GAGAGAGAGC	S	T	C	L	L
G3083u2	WIAF-12985	HT2228	2260	IGHMBP2, immunoglobulin mu binding protein 2	TGGAGTTTCAT [G/C] GCCAGCAAGA	M	G	C	M	I
G3083u3	WIAF-12986	HT2228	2060	IGHMBP2, immunoglobulin mu binding protein 2	GGGACCTCT [A/G] CGTCCACCAG	M	A	G	T	A
G3083u4	WIAF-12987	HT2228	2365	IGHMBP2, immunoglobulin mu binding protein 2	ACGACAGTTC [C/T] GGGGAAGGGA	S	C	T	S	S
G3083u5	WIAF-13005	HT2228	411	IGHMBP2, immunoglobulin mu binding protein 2	TTTGATGAGT [C/T] CCACGATTTC	M	C	T	S	P
G3083u6	WIAF-13006	HT2228	272	IGHMBP2, immunoglobulin mu binding protein 2	ATACGGGTCC [G/A] CGGCAGCTCT	M	G	A	A	T
G3083u7	WIAF-13010	HT2228	2581	IGHMBP2, immunoglobulin mu binding protein 2	TCAGGAGCGC [G/A] CAGGGGCGAGC	S	G	A	A	A
G3083u8	WIAF-13011	HT2228	2594	IGHMBP2, immunoglobulin mu binding protein 2	GGGGCAGCCC [G/A] CCAGCAAGGA	M	G	A	A	T
G3088u1	WIAF-12984	HT2318	884	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TGTGCACTA [C/T] GTCCCCCTCC	M	C	T	T	M
G3088u2	WIAF-12988	HT2318	2469	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TCTTGTCAAC [A/G] CGTCAACACC	S	A	G	P	P

G3088u3	WIAF-12989	HT2318	3056	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TTCTTGGTAC [T/C] GGACAGTCCC	S	T	C	T	T
G3088u4	WIAF-12991	HT2318	4008	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TTATCCGGCA [G/T] CACAACATCC	M	G	T	Q	H
G3088u5	WIAF-12992	HT2318	4880	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	CAAATCCATG [C/G] ACCGCCTAGC	M	C	G	A	G
G3088u6	WIAF-12993	HT2318	5148	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TTGACAGCAT [G/A] TCTAATTGGC	M	G	A	M	I
G3088u7	WIAF-12999	HT2318	5834	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	CCAGCTGATA [A/G] TTCATCAACA	M	A	G	N	S
G3088u8	WIAF-13000	HT2318	6065	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	CAAAGTCAAC [G/A] GCCAGTCACT	M	G	A	R	Q
G3088u9	WIAF-13001	HT2318	7652	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	CATAGGAATA [C/T] GGTACAGAA	M	C	T	T	M
G3088u10	WIAF-13008	HT2318	741	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TTCTGCAGCA [A/G] CCATCTGAAC	S	A	G	Q	Q
G3088u11	WIAF-13009	HT2318	948	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	CAGAACTGAG [C/T] ACCTTGTCAAC	S	C	T	S	S
G3088u12	WIAF-13012	HT2318	1909	HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	TGAAACTTTTA [C/T] TAAATCAAG	S	C	T	L	L

G3088u13	WIAF-13013	HT2318		HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	2803	TCTTCTGTCT [G/A]TACCTTCACT	M G A V I
G3088u14	WIAF-13015	HT2318		HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	3342	GGGTCTGCA [A/G]CCTCAGATTC	S A G Q Q
G3088u15	WIAF-13016	HT2318		HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	3542	CCTAAACATA [G/A]TGTTACCATA	M G A S N
G3088u16	WIAF-13017	HT2318		HIVEP1, human immunodeficiency virus type I enhancer-binding protein 1	4972	TGGGTCTTCT [A/G]AAAGTGAGGA	M A G K E
G3095u1	WIAF-12994	HT2435		TCF2, transcription factor 2, hepatic; LF-B3; variant hepatic nuclear factor	701	CGGCTCTGTA [C/T]ACCTGGTAAG	S C T Y Y
G3095u2	WIAF-13018	HT2435		TCF2, transcription factor 2, hepatic; LF-B3; variant hepatic nuclear factor	362	GGGCGGAGCC [C/T]GACACCAAGC	S C T P P
G3095u3	WIAF-13020	HT2435		TCF2, transcription factor 2, hepatic; LF-B3; variant hepatic nuclear factor	1620	CCAGTTCTCC [C/T]AGCAGCTGCA	N C T Q *
G3100a1	WIAF-12147	HT2483		ZNF141, zinc finger protein 141 (clone pHZ-44)	526	GAATGAGTGT [A/G]AGTTGCAGAA	M A G K E
G3102u1	WIAF-12975	HT2508		NRF1, nuclear respiratory factor 1	259	CGCCTTCTTC [G/T]CCCGAGGACA	S G T S S
G3103u1	WIAF-13617	HT2511		E2F2, E2F transcription factor 2	1106	CCTTGGACCA [G/T]CTCATCCAGA	M G T Q H
G3103u2	WIAF-13659	HT2511		E2F2, E2F transcription factor 2	1154	CTGAGGACAA [G/A]GCCAACAAAGA	S G A K K
G311u1	WIAF-10291	HT0402		alpha-2-macroglobulin	1339	GTCCCTTGTTA [C/T]GGCTACCAGT	S C T Y Y
G311u2	WIAF-10292	HT0402		alpha-2-macroglobulin	1201	TCATATTTCAT [C/T]AGAGGAAATG	S C T I I
G311u3	WIAF-10293	HT0402		alpha-2-macroglobulin	3041	TACTCCAGAG [G/A]TCAAGTCCAA	M G A V I

G311u4	WIAF-10294	HT0402	3676 A2M, alpha-2-macroglobulin	TCACATCCTA [T/C] GTGCTCTCG	S	T	C	Y	Y
G311u5	WIAF-10296	HT0402	3364 A2M, alpha-2-macroglobulin	ATATCACCAT [C/T] GCCCTTCTGG	S	C	T	I	I
G311u6	WIAF-10297	HT0402	3203 A2M, alpha-2-macroglobulin	CCAAGCTCGA [G/T] CCTACATCTT	M	G	T	A	S
G311a7	WIAF-10494	HT0402	1122 A2M, alpha-2-macroglobulin	TCACACTTTC [G/A] ACAGGGAATT	M	G	A	R	Q
G3119u1	WIAF-13947	HT2654	GLI, glioma-associated oncogene homolog (zinc finger protein)	TTTCTGGGGG [G/A] TTCCAGGTT	M	G	A	G	D
G3119u2	WIAF-13959	HT2654	GLI, glioma-associated oncogene homolog (zinc finger protein)	AGTCCCGGGA [G/A] GAACCTTGG	S	G	A	E	E
G3119u3	WIAF-13965	HT2654	GLI, glioma-associated oncogene homolog (zinc finger protein)	TGGGGAACA [G/C] AATTCTCTCAA	M	G	C	E	Q
G312u1	WIAF-10006	HT0428	PLAU, plasminogen activator, urokinase	CTCACCACAA [C/T] GACATTTGCT	S	C	T	N	N
G312u2	WIAF-10029	HT0428	PLAU, plasminogen activator, urokinase	GGCCTAAAGC [C/T] CCTTGTCCAA	M	C	T	P	L
G312a3	WIAF-10521	HT0428	PLAU, plasminogen activator, urokinase	TGATTACCCA [A/C] AGAAGGAGGA	M	A	C	K	Q
G3125u1	WIAF-13675	HT2674	GRF2F2, general transcription factor IIF, polypeptide 2 (30kD subunit)	ACATCACAAA [A/G] CAACCTGTGG	S	A	G	K	K
G313u1	WIAF-10129	HT0462	platelet-derived growth factor, alpha polypeptide (GB:M21574)	CATGCGTGTG [G/A] ACTCAGACAA	M	G	A	D	N
G313u2	WIAF-10130	HT0462	platelet-derived growth factor, alpha polypeptide (GB:M21574)	ATGAGAAAGG [T/G] TTCATTGAAA	S	T	G	G	G
G313u3	WIAF-10133	HT0462	platelet-derived growth factor, alpha polypeptide (GB:M21574)	GGAGATCCAC [T/C] CCGAGACAG	M	T	C	S	P
G313u4	WIAF-10135	HT0462	platelet-derived growth factor, alpha polypeptide (GB:M21574)	CTCGCAACGT [C/T] CTCCTGGCAC	S	C	T	V	V
G314u1	WIAF-10069	HT0467	ALOX15, arachidonate 15-lipoxygenase	TCAGGAGGGA [G/A] CTGGCTGCC	S	G	A	E	E

G3141u1	WIAF-13934	HT27498		878	NFATC3, nuclear factor of activated T-cells, cytoplasmic 3	CCAGAGGATA[G/A]CTGGCTACTC	M	G	A	S	N
G3141u2	WIAF-13936	HT27498		1189	NFATC3, nuclear factor of activated T-cells, cytoplasmic 3	GCCTGCCTCA[T/C]GCAATGGAA	M	T	C	C	R
G3141u3	WIAF-13938	HT27498		2241	NFATC3, nuclear factor of activated T-cells, cytoplasmic 3	CTCTGGGGG[T/C]TTCCTTCAG	S	T	C	G	G
G3141u4	WIAF-13944	HT27498		702	NFATC3, nuclear factor of activated T-cells, cytoplasmic 3	ATCCCTCTGA[C/T]GAGCAGCCC	S	C	T	D	D
G3159u1	WIAF-13891	HT2757		523	SP4, Sp4 transcription factor	CTTCAAAAAGA[G/A]AATAACGTTT	S	G	A	E	E
G3159u2	WIAF-13892	HT2757		1514	SP4, Sp4 transcription factor	ACAGAAATGTT[C/T]AACTTCAAGC	N	C	T	Q	*
G3159u3	WIAF-13893	HT2757		2236	SP4, Sp4 transcription factor	TGTTTGTGG[C/T]AAAAGATTCA	S	C	T	G	G
G3165u1	WIAF-13860	HT27636		437	transcription factor B-ATF	AGCAGCTCAC[A/G]GAGGAACCTGA	S	A	G	T	T
G3165u2	WIAF-13861	HT27636		512	transcription factor B-ATF	CCAGCAGCC[C/G]TCGCCCCCG	S	C	G	P	P
G3173u1	WIAF-13556	HT2772		1686	ZNF74, zinc finger protein 74 (Cos52)	TGCACAGCGA[G/A]GGGAAGCCCT	S	G	A	E	E
G3175u1	WIAF-13948	HT2776		2037	transcriptional regulator, via glucocorticoid receptor	TGTTTCGGACC[A/G]GAAGCACCCA	S	A	G	P	P
G3182u1	WIAF-14036	HT2783		1614	MHC2TA, MHC class II transactivator	ATCCTAGACG[C/G]CTTCGAGGAG	M	C	G	A	G
G3182u2	WIAF-14037	HT2783		2791	MHC2TA, MHC class II transactivator	TGAGCGACAC[G/A]GTGGGCTGT	S	G	A	T	T
G3182u3	WIAF-14059	HT2783		1657	MHC2TA, MHC class II transactivator	TGCACAGCAC[G/A]TCGGGACCGG	S	G	A	T	T
G3182u4	WIAF-14060	HT2783		1606	MHC2TA, MHC class II transactivator	TTCTGCTCAT[C/T]CTAGACGCTT	S	C	T	I	I
G3183u1	WIAF-13950	HT27861		392	zinc finger protein C2H2-150	TACTCTAGAG[G/A]AGCCTGTGG	M	G	A	E	K
G3184u1	WIAF-13864	HT27862		271	zinc finger protein C2H2-171	GAAACTCCAG[T/G]TCAAAGACTT	M	T	G	F	V
G3184u2	WIAF-13865	HT27862		248	zinc finger protein C2H2-171	CTGCTTGAAT[T/C]CATGTATGAR	M	T	C	F	S
G320u1	WIAF-10136	HT0791		552	ANX7, annexin VII (synexin)	CCAACCTCGA[T/C]GCTATAAGAG	S	T	C	D	D

G320u2	WIAF-10137	HT0791	1350 ANX7, annexin VII (synexin)	TTGACCTTGT [A/G]CAAATAAAC	S	A	G	V	V
G320u1	WIAF-14186	HT27930	485 zinc finger protein ZNF37A	GTCAGAAGTC [A/G]GCCCTAATTG	S	A	G	S	S
G3218u1	WIAF-13526	HT28104	zinc finger protein ZNF169, 187 Krueppel-type	CCCGACAGCT [C/T]ATTAAGAAG	M	C	T	H	Y
G323u1	WIAF-10066	HT0915	Homo sapiens inducible nitric oxide synthase (NOS) mRNA, complete cds.	ACTTCTGTGA [C/T]GTCCAGCGCT	S	C	T	D	D
G325u1	WIAF-10106	HT0962	FBN1, fibrillin 1 (Marfan syndrome)	TGTGAATGCC [C/T]GCCTGGCCAT	M	C	T	P	L
G325u2	WIAF-10113	HT0962	FBN1, fibrillin 1 (Marfan syndrome)	AGATAGCTCC [T/G]TCCTGTGGCT	S	T	G	P	P
G325u3	WIAF-10114	HT0962	FBN1, fibrillin 1 (Marfan syndrome)	GATCTGCAAT [A/C]ATGACCGCTG	M	A	C	N	H
G325u4	WIAF-10116	HT0962	FBN1, fibrillin 1 (Marfan syndrome)	GAACTGCACA [G/C]ACATTGACGA	M	G	C	D	H
G325u5	WIAF-10117	HT0962	FBN1, fibrillin 1 (Marfan syndrome)	TCCTGCATGAA [C/T]GGCGCTTGCG	S	C	T	N	N
G326u1	WIAF-10036	HT1009	KLKB1, kallikrein B plasma, 1854 (Fletcher factor) 1	GCAACACAA [C/T]GGAATGTGGC	S	C	T	N	N
G327u1	WIAF-10052	HT1011	1599 HRG, histidine-rich glycoprotein	AAGCCAGACA [A/T]TCAGCCCTTT	M	A	T	N	I
G327u2	WIAF-10054	HT1011	1083 HRG, histidine-rich glycoprotein	CCACTATTGC [C/T]CATGTCCTGC	M	C	T	P	L
G327u3	WIAF-10055	HT1011	1140 HRG, histidine-rich glycoprotein	GCCCAAAGAC [A/G]TTCTCATAAT	M	A	G	H	R
G328u1	WIAF-10145	HT1087	255 SAAL, serum amyloid A1	GTGCTGGGC [T/C]GCAGAAGTGA	S	T	C	A	A
G328a2	WIAF-10511	HT1087	248 SAAL, serum amyloid A1	CCTGGGGTG [C/T]CTGGGCTGCA	M	C	T	A	V
G328a3	WIAF-10512	HT1087	305 SAAL, serum amyloid A1	TTCTTTGGCC [A/G]TGGTGGGAG	M	A	G	H	R
G328a4	WIAF-13126	HT1087	295 SAAL, serum amyloid A1	TATCCAGAGA [T/C]TCTTTGGCCA	M	T	C	F	L
G328a5	WIAF-13127	HT1087	82 SAAL, serum amyloid A1	CTTGGTCCTG [G/A]GTGTCAGCAG	M	G	A	G	S
G329u1	WIAF-10140	HT1141	PLCG1, phospholipase C, gamma 1 (formerly subtype 148)	CTGACCTTCA [T/C]CAAGAGGCGC	M	T	C	I	T
G329u2	WIAF-10162	HT1141	PLCG1, phospholipase C, gamma 1 (formerly subtype 148)	TATGCCCGGA [C/A]ACCATGAACA	M	C	A	D	E
G329u3	WIAF-10163	HT1141	PLCG1, phospholipase C, gamma 1 (formerly subtype 148)	GTTCAATGCTC [A/G]GCTTCCTCCG	M	A	G	S	G

G3295u1	WIAF-14017	HT3460	1229	FUBP, far upstream element binding protein	CCATAAAAAG [C/T] ATAAGCCAGC	S	C	T	S	S
G3296u1	WIAF-14168	HT3466	6289	transcription factor TFIIC, RNA polymerase III, alpha subunit	CAGCCTGGAC [G/A] AGAGCCCAT	M	G	A	E	K
G3296u2	WIAF-14179	HT3466	235	transcription factor TFIIC, RNA polymerase III, alpha subunit	GGGCATCAGC [T/A] TCTATGAGGA	M	T	A	F	I
G3298u1	WIAF-13523	HT3504	1803	DNA-binding protein HRFX2	ACTTTGCCAA [C/T] GTCCAGGAGC	S	C	T	N	N
G3298u2	WIAF-13524	HT3504	1743	DNA-binding protein HRFX2	GGCGGTGCT [G/A] CAGAACACGT	S	G	A	L	L
G3298u3	WIAF-13528	HT3504	2002	DNA-binding protein HRFX2	GTCTTGCTG [A/G] AATGGTCTCTT	M	A	G	K	E
G33u1	WIAF-10254	X82540	1044	INHBC, inhibin, beta C	AAGGCCAACA [C/T] AGTGCAGGC	M	C	T	T	I
G33u2	WIAF-10255	X82540	1136	INHBC, inhibin, beta C	CAGCAACATT [G/A] TCAAGACTGA	M	G	A	V	I
G33u3	WIAF-10256	X82540	1185	INHBC, inhibin, beta C	GGGTGCAGTT [A/G] GTCTATGTGT	N	A	G	*	W
G33u4	WIAF-10259	X82540	892	INHBC, inhibin, beta C	TTTTTGTTGA [C/T] TTCCGTGAGA	S	C	T	D	D
G3303u1	WIAF-13566	HT3523	981	POU6F1, POU domain, class 6, transcription factor 1	CAGGCCAGGA [G/A] ATCACTGAAA	S	G	A	E	E
G3304u1	WIAF-13932	HT3544	970	SP2, Sp2 transcription factor	TCAACACCT [C/T] GTGAACGCCA	S	C	T	L	L
G3304u2	WIAF-13935	HT3544	1891	SP2, Sp2 transcription factor	AGAAGCAGGT [T/G] TCCACATCC	S	T	G	V	V
G3304u3	WIAF-13943	HT3544	920	SP2, Sp2 transcription factor	TGTGGTGAAG [T/C] TGACAGGTGG	S	T	C	L	L
G3311u1	WIAF-13839	HT3585	757	GATA3, GATA-binding protein 3	CCCACTCCCG [T/C] GGCAGCATGA	S	T	C	R	R
G3311u2	WIAF-13840	HT3585	901	GATA3, GATA-binding protein 3	TCGATGCAA [G/A] TCCAGGCCCA	S	G	A	K	K
G3316u1	WIAF-13818	HT3607	282	zinc finger protein HKE-T1, Kruppel-like	AAAGATTTC [A/G] GTCAGAGTTC	M	A	G	S	G
G3319u1	WIAF-14214	HT3613	1086	SMARCA3, SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 3	AAACTCTTAC [A/G] GCCATTGCAG	S	A	G	T	T
G3319u2	WIAF-14221	HT3613	1261	SMARCA3, SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 3	TAGATGTAGT [G/C] AACAAACCCAG	M	G	C	E	Q
G3320u1	WIAF-13692	HT3622	624	BCL6, B-cell CLL/lymphoma 6 (zinc finger protein 51)	ATTTCGGGA [G/C] GGCAACATCA	M	G	C	E	D

002000 2445960

G3320u2	WIAF-13717	HT3622	1062	BCL6, B-cell CLL/lymphoma 6 (zinc finger protein 51)	ACAGCGGGCC [G/A] ACTTTGGAGG	S	G	A	P	P
G3321u1	WIAF-13761	HT3641	235	STAT2, signal transducer and activator of transcription 2, 113kD	TCTTGGATCA [G/C] CTGAACATATG	M	G	C	Q	H
G3321u2	WIAF-13762	HT3641	774	STAT2, signal transducer and activator of transcription 2, 113kD	CAAAAAGCCT [G/C] CACFAGAGCT	M	G	C	C	S
G3328u1	WIAF-13543	HT3681	1550	transcription factor znf6	CCACATGGT [A/G] TCAGAGAGG	S	A	G	V	V
G3328u2	WIAF-13544	HT3681	1389	transcription factor znf6	AGAGATTTA [G/C] AGGAAGATGA	M	G	C	E	Q
G3336u1	WIAF-13848	HT3732	216	XBPL1, X-box binding protein 1	ACCTGAGCCC [C/T] GAGGAGAAGG	S	C	T	P	P
G334u1	WIAF-10008	HT1220	893	THBS1, thrombospondin 1	TACATTGGCC [A/C] CAAGACAAAG	M	A	C	H	P
G334u2	WIAF-10009	HT1220	2000	THBS1, thrombospondin 1	TCACAGCCCT [T/C] CGGCCAGGGT	M	T	C	F	S
G334u3	WIAF-10016	HT1220	1521	THBS1, thrombospondin 1	CCAGATGAA [T/C] GGGAAACCTG	S	T	C	N	N
G334u4	WIAF-10017	HT1220	2210	THBS1, thrombospondin 1	GGTGGCCCA [A/G] TGAGAACCTG	M	A	G	N	S
G334u5	WIAF-10018	HT1220	2979	THBS1, thrombospondin 1	GTGAGACCGA [T/C] TTCGCCGAT	S	T	C	D	D
G334u6	WIAF-10033	HT1220	1136	THBS1, thrombospondin 1	TGTCACCTGTC [A/G] GAACTCAGTT	M	A	G	Q	R
G334u7	WIAF-10034	HT1220	1859	THBS1, thrombospondin 1	AGTGAAATG [G/A] CATCCAGTGC	M	G	A	G	D
G3343u1	WIAF-13545	HT3770	1104	ZNF76, zinc finger protein 76 (expressed in testis)	GCAGTGCCCA [C/T] GGCAGAGCTGG	S	C	T	H	H
G3343u2	WIAF-13561	HT3770	425	ZNF76, zinc finger protein 76 (expressed in testis)	GAGCAGTATG [C/A] CAGCAAGGTT	M	C	A	A	D
G3343u3	WIAF-13562	HT3770	143	ZNF76, zinc finger protein 76 (expressed in testis)	CACCAGGTGA [C/T] GGTACAGAAA	M	C	T	T	M
G3343u4	WIAF-13563	HT3770	646	ZNF76, zinc finger protein 76 (expressed in testis)	GAAGAGCCAC [G/T] TTCGTACCCA	M	G	T	V	F
G3343u5	WIAF-13564	HT3770	611	ZNF76, zinc finger protein 76 (expressed in testis)	AGCTGTGGAA [A/G] GGCCTTTGCC	M	A	G	K	R
G3344u1	WIAF-13664	HT3772	925	zinc finger protein MAZ	AGCTGTGCA [C/T] TCGGACGAGA	S	C	T	H	H
G3345u1	WIAF-13508	HT3823	315	TCF6L1, transcription factor 6-like 1 (mitochondrial transcription factor 1-like)	TTCCGATTTTC [T/C] AAAGAACAAC	S	T	C	S	S

G3345u2	WIAF-13509	HT3823	167	TCF6L1, transcription factor 6-like 1 (mitochondrial transcription factor 1-like)	GGCGTGTGA [G/C]TGCCTGGGA	M	G	C	S	T
G3345u3	WIAF-13531	HT3823	625	TCF6L1, transcription factor 6-like 1 (mitochondrial transcription factor 1-like)	TTATAACGTT [T/G]ATGTAGTGA	M	T	G	Y	D
G3352u1	WIAF-13589	HT4005	1190	MITF, microphthalmia-associated transcription factor	CTCGGAACGT [G/A]GACTGAGGCC	M	G	A	G	E
G3352u2	WIAF-13604	HT4005	1156	MITF, microphthalmia-associated transcription factor	TCTCAGGGAT [G/A]GCACCATCAC	M	G	A	G	S
G3353u1	WIAF-13937	HT4010	360	GTF2H3, general transcription factor IIH, polypeptide 3 (34kD subunit)	ATCTAATGAC [C/A]AAAAATGACA	S	C	A	T	T
G3358u1	WIAF-13671	HT4187	398	ETV5, ets variant gene 5 (ets-related molecule)	GATGATGAAC [A/G]GTTTGTCCTCA	M	A	G	Q	R
G3358u2	WIAF-13672	HT4187	223	ETV5, ets variant gene 5 (ets-related molecule)	TCAGCAAGTC [C/T]CTTTTATGGT	M	C	T	P	S
G3358u3	WIAF-13673	HT4187	1236	ETV5, ets variant gene 5 (ets-related molecule)	GACTGGAAGG [C/G]AAAGTCAAAC	S	C	G	G	G
G3358u4	WIAF-13674	HT4187	1678	ETV5, ets variant gene 5 (ets-related molecule)	TTACCTCCTG [G/A]ACATGGACCG	M	G	A	D	N
G3358u5	WIAF-13706	HT4187	414	ETV5, ets variant gene 5 (ets-related molecule)	TCCCAGATTT [T/C]CAGTCTGATA	S	T	C	F	F
G3358u6	WIAF-13707	HT4187	1238	ETV5, ets variant gene 5 (ets-related molecule)	CTGGAAGGCA [A/G]AGTCAAAACAG	M	A	G	K	R
G336u1	WIAF-10152	HT1258	566	ACAT1, acetyl-Coenzyme A acetyltransferase 1 (acetoacetyl Coenzyme A thiolase)	AGAGCATGTC [C/A]AATGTTCCAT	S	C	A	S	S
G3369u1	WIAF-14047	HT4302	614	zinc finger protein DB1	ATCTCAATCG [A/G]CACAACTCT	S	A	G	R	R
G337u1	WIAF-10268	HT1259	464	EDNRB, endothelin receptor type B	AAAGGAGACA [G/T]GACGCAGGA	M	G	T	R	M
G337u2	WIAF-10298	HT1259	1281	EDNRB, endothelin receptor type B	TGAAGCTCAC [T/A]CTTTATAATC	S	T	A	T	T

G3373u1	WIAF-14203	HT4342	MTF1, metal-regulatory transcription factor 1	1253	CTCAACAGAC [A/G] GCTTCCTTGA	S	A	G	T	T
G3390u1	WIAF-14182	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	680	AGAGCCAGAG [C/T] TCTACCTCGA	M	C	T	L	F
G3390u2	WIAF-14184	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	1026	GCTCAGACAG [G/A] GAACCTTGAG	M	G	A	G	E
G3390u3	WIAF-14185	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	1423	AAAAGCCTTA [T/C] GTGTGCCGG	S	T	C	Y	Y
G3390u4	WIAF-14197	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	811	CTGGGGATCC [A/G] GGCCCGGGG	S	A	G	P	P
G3390u5	WIAF-14198	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	1420	GGGAAAAGCC [T/G] TATGTGTGCC	S	T	G	P	P
G3390u6	WIAF-14199	HT4483	ZNF133, zinc finger protein 133 (clone PHZ-13)	2143	CAAGCTCTAAT [C/T] ACACACAAGC	S	C	T	I	I
G3391u1	WIAF-13631	HT4484	ZNF136, zinc finger protein 136 (clone PHZ-20)	391	AGCATTGTAT [A/G] TGGAGAAGTC	M	A	G	Y	C
G3396u1	WIAF-13978	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1283	CACAGCTCCT [C/T] GCTCAGCCAG	M	C	T	S	L
G3396u2	WIAF-13979	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1296	TCAGCCAGCA [C/T] GAAAGGACGC	S	C	T	H	H
G3396u3	WIAF-13980	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1028	AGTCACAGCT [C/T] GTCCCTCACC	M	C	T	S	L
G3396u4	WIAF-13981	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1057	GCGAATCCAC [A/G] CTGGGGAGAA	M	A	G	T	A
G3396u5	WIAF-13982	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1152	CAGGAGAGAA [A/G] CCTATGAAT	S	A	G	K	K
G3396u6	WIAF-13983	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1243	AAAGCCGTAT [G/C] GGTGCAATGA	M	G	C	G	R
G3396u7	WIAF-13984	HT4491	ZNF135, zinc finger protein 135 (clone PHZ-17)	1045	CACCAACAT [C/T] AGCGAATCCA	N	C	T	Q	*
G340u1	WIAF-10139	HT1386	CYP27A1, cytochrome P450, subfamily XXVIIA (steroid 27-hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1	459	CCTATGGGCC [G/A] TTCACACGG	S	G	A	P	P
G340u2	WIAF-10160	HT1386	CYP27A1, cytochrome P450, subfamily XXVIIA (steroid 27-hydroxylase, cerebrotendinous xanthomatosis), polypeptide 1	801	TCCCAAGTG [G/A] ACTGCCCCG	N	G	A	W	*

G341u1	WIAF-10121	HT1388	MUT, methylmalonyl Coenzyme A mutase	912	GAGCTGGCCT [A/G] TACTTTAGCA	M	A	G	Y	C
G341u2	WIAF-10128	HT1388	MUT, methylmalonyl Coenzyme A mutase	2087	TGCTGTGGGC [G/A] TAAGCACCT	M	G	A	V	I
G3410u1	WIAF-13749	HT4550	1720 zinc finger homeodomain protein		TGAGTCCTCT [G/T] TTTCATCAGC	M	G	T	V	F
G3410u2	WIAF-13750	HT4550	2843 zinc finger homeodomain protein		AAACATCAAT [T/C] GATGACAC	M	T	C	L	S
G3410u3	WIAF-13751	HT4550	2745 zinc finger homeodomain protein		AGATATTCCA [A/T] AAGAGTAGTT	M	A	T	Q	H
G3410u4	WIAF-13775	HT4550	236 zinc finger homeodomain protein		AGAGAGGGA [A/C] TGCTAAGAAC	M	A	C	N	T
G3410u5	WIAF-13776	HT4550	195 zinc finger homeodomain protein		TGCCAACAGA [C/T] CAGACAGTGT	S	C	T	D	D
G3410u6	WIAF-13777	HT4550	606 zinc finger homeodomain protein		ATAACTTTAG [T/C] TGCTCCCTGT	S	T	C	S	S
G3410u7	WIAF-13793	HT4550	2073 zinc finger homeodomain protein		CAGTTTTACC [A/G] GTGGGATCAA	S	A	G	P	P
G343u1	WIAF-10120	HT1552	561 HK1, hexokinase 1		CTTGCCAAACA [A/G] TCCAAAATAG	S	A	G	Q	Q
G343u2	WIAF-10124	HT1552	159 HK1, hexokinase 1		ACAAATATCT [G/C] TATGCCATGC	S	G	C	L	L
G348u1	WIAF-10269	HT1906	PECAM1, platelet/endothelial cell adhesion molecule (CD31 antigen)	2212	TGACGATGTC [A/G] GAAACCATGC	S	A	G	G	G
G348u2	WIAF-10277	HT1906	PECAM1, platelet/endothelial cell adhesion molecule (CD31 antigen)	1656	GCCATTCCCA [C/T] GCCAAAATGT	S	C	T	H	H
G348u3	WIAF-10283	HT1906	PECAM1, platelet/endothelial cell adhesion molecule (CD31 antigen)	577	AGAGTACCAG [C/G] TGTTGGTGGA	S	C	G	V	V
G348a5	WIAF-13119	HT1906	PECAM1, platelet/endothelial cell adhesion molecule (CD31 antigen)	?	ATTGTTCC [C/G]	?	C	G		
G351u1	WIAF-10123	HT1990	OSBP, oxysterol binding protein	1047	TGCTGGCAGA [G/A] TCAGATGAAT	S	G	A	E	E
G351u2	WIAF-10132	HT1990	OSBP, oxysterol binding protein	1023	TGGCCAAGGC [C/A] AAAGCTGTGA	S	C	A	A	A
G355u1	WIAF-10146	HT2143	THBS4, thrombospondin 4	1670	AACTGCCTGA [G/A] TGCTTAAAT	M	G	A	S	N
G355u2	WIAF-10165	HT2143	THBS4, thrombospondin 4	1186	TCGAAATGGA [G/C] CGTGGCTTCC	M	G	C	A	P

G355a3	WIAF-10510	HT2143	1962 THBS4, thrombospondin 4	ACTGCCCCAC [C/G] GTCATTAAACA	S	C	G	T	T
G355a4	WIAF-13125	HT2143	1963 THBS4, thrombospondin 4	CTGCCCCACC [G/a] TCATTAAACAG	M	G	a	V	I
G3552u1	WIAF-12701	HT28101	1006 CLCN2, chloride channel 2	AAGAGACTAT [T/C] ACAGCCCTCT	S	T	C	I	I
G3552u2	WIAF-12731	HT28101	1823 CLCN2, chloride channel 2	CGGCCACCAG [C/T] AGTACCAGGT	N	C	T	Q	*
G3552u3	WIAF-12736	HT28101	2254 CLCN2, chloride channel 2	GGAGCGCAGA [G/C] TCGGCAGGCA	M	G	C	E	D
G3555u1	WIAF-12744	HT2896	334 calyculin	GCCCTCAAGG [G/a] CTGAAATAA	M	G	A	G	D
G357u1	WIAF-10267	HT2244	4300 C4B, complement component 4B	ATGAGTACGA [T/C] GAGCTTCCAG	S	T	C	D	D
G357u2	WIAF-10280	HT2244	5095 C4B, complement component 4B	TCATGGGTCT [G/a] GATGGGGCCA	S	G	A	L	L
G357u3	WIAF-10295	HT2244	2996 C4B, complement component 4B	CTCAGATCCA [T/C] TGGACACTTT	S	T	C	L	L
G359u1	WIAF-10026	HT2411	PIAT, plasminogen activator, tissue	CGCAGGCTGA [A/G] GTGGGAGTAC	M	A	G	T	M
G359a2	WIAF-10520	HT2411	PIAT, plasminogen activator, tissue	AGGCTTGTG [T/C] CCTTCTATT	S	T	C	S	S
G3592u1	WIAF-12759	HT4214	743 CLCN4, chloride channel 4	CTTCTAACGA [G/a] ACCACTTTTG	S	G	A	E	E
G3592u2	WIAF-12761	HT4214	835 CLCN4, chloride channel 4	GCTTACATTC [T/G] GAATTACTTA	M	T	G	L	R
G361u1	WIAF-10053	HT2479	cystathionine beta synthase, alt. transcript 1	TGGTCTACTA [C/T] GACACCACCG	S	C	T	Y	Y
G361u2	WIAF-10056	HT2479	cystathionine beta synthase, alt. transcript 1	TCATFCCCAC [G/a] GTGCTGGACA	S	G	A	T	T
G362u1	WIAF-10058	HT2638	ADRB2, adrenergic, beta-2-, receptor, surface	GGCACCCCAAT [G/a] GAAGCCATGC	M	G	A	G	R
G362u2	WIAF-10059	HT2638	ADRB2, adrenergic, beta-2-, receptor, surface	TCATGGGCT [G/a] GCAGTGGTGC	S	G	A	L	L
G362u3	WIAF-10060	HT2638	ADRB2, adrenergic, beta-2-, receptor, surface	CGTCACGCAG [G/C] AAAGGGACGA	M	G	C	E	Q
G362u4	WIAF-10093	HT2638	ADRB2, adrenergic, beta-2-, receptor, surface	AGGCTATGG [G/C] AATGGTACT	S	G	C	G	G
G3620u1	WIAF-12808	HT97200	ACATN, acetyl-Coenzyme A transporter	CACCTCTCTGG [A/G] TATGAAGAGC	M	A	G	D	G
G3627u1	WIAF-12820	HT97387	NAPG, N-ethylmaleimide-sensitive factor attachment protein, gamma	GCAGAAACTA [C/T] CAGAGCCGT	M	C	T	P	S
G366u1	WIAF-10046	HT2764	987 BDKRB2, bradykinin receptor B2	GCCTCTTCA [T/C] GGCCTACAGC	M	T	C	M	T
G366a2	WIAF-10500	HT2764	820 BDKRB2, bradykinin receptor B2	AGATCCAGAC [G/a] GAGAGAGGG	S	G	A	T	T

G366a3	WIAF-10501	HT2764	961 BDKRB2, bradykinin receptor B2	GCATCATCGA [T/C] GTAATCACAC	S	T	C	D	D
G367u1	WIAF-10156	HT27685	ACACA, acetyl-Coenzyme A	ATCATCCATA [T/C] GACGAGCAC	N	T	C	*	C
G370u1	WIAF-10281	HT27888	6965 carboxylase alpha	AAAATCTCC [G/A] TTGAAGATT	S	G	A	P	P
G370u2	WIAF-10282	HT27888	3250 LEPR, leptin receptor	TCACCAAGTG [C/T] TTCTCTAGCA	S	C	T	C	C
G370u3	WIAF-10284	HT27888	3229 LEPR, leptin receptor	CAATATCAAG [T/C] GAAATATCA	M	T	C	V	A
G370u4	WIAF-10285	HT27888	1005 LEPR, leptin receptor	CAGAGAATAA [C/T] CTTCAATCC	S	C	T	N	N
G370u5	WIAF-10299	HT27888	1894 LEPR, leptin receptor	TTCTGACAAG [T/C] GTTGGGTCTA	S	T	C	S	S
G370u6	WIAF-10300	HT27888	1222 LEPR, leptin receptor	CTATGAAAAA [G/C] GAGAAAAATG	M	G	C	K	N
G371u1	WIAF-10107	HT27943	2161 LEPR, leptin receptor						
G371a2	WIAF-12093	HT27943	349 CRAT, carnitine acetyltransferase	TCATCTACTC [G/C] AGCCACGGCG	S	G	C	S	S
G372a1	WIAF-10506	HT28247	287 CRAT, carnitine acetyltransferase	GGAGAACTGG [C/T] TGCTGAGTG	S	C	T	L	L
G374u1	WIAF-10103	HT28496	HADHA, hydroxyacyl-Coenzyme A						
G374u2	WIAF-10104	HT28496	dehydrogenase/3-ketoacyl-Coenzyme A						
G374u3	WIAF-10105	HT28496	A thiolase/enoyl-Coenzyme A						
G374u4	WIAF-10115	HT28496	hydratase (trifunctional protein),						
G374u5	WIAF-10119	HT28496	1099 alpha subunit	TGGAGCTCCA [C/A] AGAAGGATGT	M	C	A	Q	K
G374a6	WIAF-12094	HT28496	4435 FASN, fatty acid synthase	CACCTCCAC [G/A] TCCCGGAGT	M	G	A	V	I
G377u1	WIAF-10142	HT2996	5996 FASN, fatty acid synthase	CTGGACAGGG [T/C] GACCCGAGAG	M	T	C	V	A
G377u2	WIAF-10143	HT2996	5644 FASN, fatty acid synthase	CAAGAGTAC [A/G] TCATCGCTGG	M	A	G	I	V
G380u1	WIAF-10122	HT3159	6387 FASN, fatty acid synthase	TGGCACACAT [C/T] CTGGCAATCC	S	C	T	I	I
G380u2	WIAF-10126	HT3159	567 FASN, fatty acid synthase	GGGCAATCAA [C/T] GTCTCTCTGA	S	C	T	N	N
G380u4	WIAF-11605	HT3159	5520 FASN, fatty acid synthase	ACATGGCCCA [A/G] GGGGAAGCACA	S	A	G	Q	Q
G383u1	WIAF-10125	HT33546	PCCB, propionyl Coenzyme A	GGACCCGGCT [T/C] CCGTCCGTGA	M	T	C	S	P
G385u1	WIAF-10141	HT3383	929 carboxylase, beta polypeptide						
G387u2	WIAF-10143	HT2996	PCCB, propionyl Coenzyme A	CACCTTTGTG [G/A] TGATACCAAC	M	G	A	G	D
G380u1	WIAF-10122	HT3159	1416 carboxylase, beta polypeptide	TCTACCTTGA [C/T] GGCAGGTGTG	S	C	T	D	D
G380u2	WIAF-10126	HT3159	831 INSR, insulin receptor	GGCAGGATGC [A/G] TGTGTTCCA	S	A	G	A	A
G380u4	WIAF-11605	HT3159	1698 INSR, insulin receptor	GCCTGCCAC [G/A] AGTCCGAGG	S	G	A	T	T
G383u1	WIAF-10125	HT33546	2382 INSR, insulin receptor						
G385u1	WIAF-10141	HT3383	phospholipase C, beta 3, alt. transcript 2	AGACGGGCG [G/A] AGGCTCCCC	M	G	A	R	Q
G387u2	WIAF-10143	HT2996	3633 PRCP, prolylcarboxypeptidase						
G380u1	WIAF-10122	HT3159	1505 (angiotensinase C)	ATGACAGTGC [A/G] GGAAAGCAGC	S	A	G	A	A

G385u2	WIAF-10157	HT3383	1360	PRCP, prolylcarboxypeptidase (angiotensinase C)	ATCACAGACA [C/G] TCTGGTGTCA	M	C	G	T	S
G387u1	WIAF-11729	HT3439	2697	SREBF2, sterol regulatory element binding transcription factor 2	CACTCTCCAG [G/C] AGCTCCGTGC	M	G	C	R	S
G387u2	WIAF-11770	HT3439	1901	SREBF2, sterol regulatory element binding transcription factor 2	GCTGTGCGG [C/G] CAACCTACAA	M	C	G	A	G
G388u1	WIAF-10270	HT3440	245	SELP1G, selectin P ligand	CTCCAGAAAT [G/A] CTGAGGAACA	M	G	A	M	I
G390u1	WIAF-10276	HT3568	2049	NOS3, nitric oxide synthase 3 (endothelial cell)	TTGCTCGTGC [C/G] GTGGACACAC	S	C	G	A	A
G391u1	WIAF-10013	HT3630	6205	VWF, von Willebrand factor	AGGACCTGGA [G/C] GTGATTCTCC	M	G	C	E	D
G391u2	WIAF-10265	HT3630	4554	VWF, von Willebrand factor	GCCCCTGAGA [A/G] CAAGGCCTTC	M	A	G	N	S
G391u3	WIAF-10266	HT3630	7489	VWF, von Willebrand factor	TGGCCTCAAC [C/T] GCCACCAATG	S	C	T	T	T
G391u4	WIAF-10272	HT3630	2470	VWF, von Willebrand factor	ACTGTACCAT [G/A] AGTGGAGTCC	M	G	A	M	I
G391u5	WIAF-10273	HT3630	2615	VWF, von Willebrand factor	GCTCGAGTGT [A/G] CCAAAACGTG	M	A	G	T	A
G391u6	WIAF-10274	HT3630	2635	VWF, von Willebrand factor	GCCAGAACTA [T/C] GACCTGGAGT	S	T	C	Y	Y
G391u7	WIAF-10275	HT3630	4045	VWF, von Willebrand factor	TCTCGGAACC [G/A] CCGTTGCACG	S	G	A	P	P
G391u8	WIAF-10278	HT3630	4446	VWF, von Willebrand factor	AACCTTGTCC [G/A] CTACGTCCAG	M	G	A	R	H
G391u9	WIAF-10279	HT3630	5152	VWF, von Willebrand factor	GCCCTAATGC [C/T] AACGTGCAGG	S	C	T	A	A
G391u10	WIAF-10286	HT3630	3448	VWF, von Willebrand factor	TTACCAGTGA [C/T] GTCTTCCAGG	S	C	T	D	D
G391u11	WIAF-10287	HT3630	4891	VWF, von Willebrand factor	ACATGGTGAC [C/T] GTGGAGTACC	S	C	T	T	T
G391u12	WIAF-10288	HT3630	4805	VWF, von Willebrand factor	CAGGAGCAAG [G/A] AGTTTCATGA	M	G	A	E	K
G391u13	WIAF-10289	HT3630	4943	VWF, von Willebrand factor	CCTGCAGCGG [G/T] TCGAGAGAT	M	G	T	V	L
G391u14	WIAF-10290	HT3630	4915	VWF, von Willebrand factor	TCAGCGAGGC [A/C] CAGTCCAAAG	S	A	C	A	A

G391a15	WIAF-10517	HT3630	6194 VWF, von Willebrand factor	AAACAAGGAG [C/T] AGGACCTGGA	N	C	T	Q	*
G391a16	WIAF-13222	HT3630	6419 VWF, von Willebrand factor	TCACCTTGGT [C/T] ACATCTTCAC	M	C	T	H	Y
G3941u1	WIAF-14123	HT3464	1265 mannosidase, alpha, lysosomal	CAGGTGTGCA [A/G] CCAGCTGGAG	M	A	G	N	S
G3941u2	WIAF-14135	HT3464	965 mannosidase, alpha, lysosomal	ACCAACCACA [C/T] TGTGATGACC	M	C	T	T	I
G395u1	WIAF-10271	HT4158	ECB1, endothelin converting enzyme 1	TCACTGCCGA [T/C] CAGCTCAGGA	S	T	C	D	D
G395a2	WIAF-13110	HT4158	ECB1, endothelin converting enzyme 1	CATCTACAAC [A/T] TGATAGGATA	M	A	T	M	L
G3959u1	WIAF-13634	HT4490	ADTB1, adaptin, beta 1 (beta prime)	TGAAGAAGCT [G/A] GTATACCTCT	S	G	A	L	L
G3959u2	WIAF-13640	HT4490	ADTB1, adaptin, beta 1 (beta prime)	TTCTTGGCGG [T/C] GGCTTTGACA	S	T	C	G	G
G3959u3	WIAF-13641	HT4490	ADTB1, adaptin, beta 1 (beta prime)	AGGTCCACGC [G/A] CCACTCAGCC	S	G	A	A	A
G3967u1	WIAF-13997	HT2958	ACTC, actin, alpha, cardiac muscle	GAGGCACCAC [T/C] ATGTACCTGT	S	T	C	T	T
G3968u1	WIAF-14159	HT1986	1747 ACTN3, actinin, alpha 3	CGAGGTGAC [C/T] GAGAGCGAGG	N	C	T	R	*
G3968u2	WIAF-14164	HT1986	1900 ACTN3, actinin, alpha 3	GGTGCCAGC [C/T] GTGACCAGAC	M	C	T	R	C
G3968u3	WIAF-14165	HT1986	2184 ACTN3, actinin, alpha 3	ACACCGTCTA [C/T] AGCATGGAGC	S	C	T	Y	Y
G3968u4	WIAF-14167	HT1986	2557 ACTN3, actinin, alpha 3	GATCTTGGCA [G/A] GAGACAAGAA	M	G	A	G	R
G3968u5	WIAF-14175	HT1986	1212 ACTN3, actinin, alpha 3	GGCTGTCTC [G/A] GAGATCCGGC	S	G	A	S	S
G3979u1	WIAF-13884	HT0623	776 GPC1, glypican 1	TGCTGTGCC [T/G] GATGACTACC	S	T	G	P	P
G3979u2	WIAF-13885	HT0623	680 GPC1, glypican 1	TGTACTACCG [C/T] GTGCGCAACC	S	C	T	R	R
G3979u3	WIAF-13886	HT0623	1361 GPC1, glypican 1	AGCTGGTCTC [T/C] GAAGCCCAAGG	S	T	C	S	S
G3979u4	WIAF-13887	HT0623	1163 GPC1, glypican 1	AGAGTGTCTC [C/T] GGCGGCTGTC	S	C	T	I	I
G3979u5	WIAF-13888	HT0623	1670 GPC1, glypican 1	ACGCCAGTGA [C/T] GACGGGAGCG	S	C	T	D	D
G3979u6	WIAF-13905	HT0623	1069 GPC1, glypican 1	CTTGCCAAAC [A/T] GGCGGACCTG	M	A	T	Q	L
G3979u7	WIAF-13906	HT0623	1514 GPC1, glypican 1	TCATGGGTGA [C/T] GGCTGGGCA	S	C	T	D	D
G3979u8	WIAF-13907	HT0623	1720 GPC1, glypican 1	GACCTCTGCG [G/C] CCGGAAGTCT	M	G	C	G	A
G3979u9	WIAF-13908	HT0623	1676 GPC1, glypican 1	GTGACGACGG [C/T] AGCGGCTCGG	S	C	T	G	G
G3979u10	WIAF-13909	HT0623	1719 GPC1, glypican 1	TGACCTCTGC [G/A] GCCGGAAGGT	M	G	A	G	S
G399u1	WIAF-10102	HT48511	450 AQP3, aquaporin 3	TCRGGACCTT [T/C] GCCGACCAAC	S	T	C	F	F
G399u2	WIAF-10111	HT48511	192 AQP3, aquaporin 3	GCTCCGTGGC [C/T] CAGTTTGTGC	S	C	T	A	A
G399u3	WIAF-10112	HT48511	165 AQP3, aquaporin 3	CCCTCATCCT [C/G] GTGATGTTTG	S	C	G	L	L
G3997u1	WIAF-13649	HT27682	MFAP2, microfibrillar-associated protein 2	TGTGTGCCCA [C/T] GAGGAGTCTC	S	C	T	H	H
G3997u2	WIAF-13650	HT27682	MFAP2, microfibrillar-associated protein 2	CCATACACAG [G/T] CCTTGCAAC	M	G	T	R	S

G4038u9	WIAF-14026	HT4211		LAMB3, laminin, beta 3 (nicein (125kD), kalinin (140kD), BM600 (125kD))	1553	GGCTGTGAAC [C/T] GGTGCTGTC	M	C	T	P	L
G4038u10	WIAF-14029	HT4211		LAMB3, laminin, beta 3 (nicein (125kD), kalinin (140kD), BM600 (125kD))	3562	CCTGACAGGA [C/T] TGGAGAAGCG	S	C	T	L	L
G4038u11	WIAF-14030	HT4211		LAMB3, laminin, beta 3 (nicein (125kD), kalinin (140kD), BM600 (125kD))	3546	TGCTGCGCTC [A/G] GCGGACCTGA	S	A	G	S	S
G4045u1	WIAF-13571	HT0652		1266 adducin, beta subunit	1266	TGGAGCAGGA [G/T] AAGACCGGC	M	G	T	E	D
G4050u1	WIAF-14106	HT1466		1366 villin	1366	CGTTGGCAG [G/A] GCAGCCAGGC	M	G	A	G	S
G4050u2	WIAF-14107	HT1466		1468 villin	1468	GGTCCCAATG [G/A] GCAAGGAGCC	M	G	A	G	S
G4050u3	WIAF-14108	HT1466		1932 villin	1932	CCACAGAGAT [C/T] CCTGACTTCA	S	C	T	I	I
G4050u4	WIAF-14110	HT1466		2438 villin	2438	TTTGGGATGA [C/T] TCCAGCTGCC	M	C	T	T	I
G4057u1	WIAF-13648	HT33633		371 CNN3, calponin 3, acidic	371	TTCAGGCTTA [T/C] GGTATGAAGC	S	T	C	Y	Y
G4066u1	WIAF-13676	HT4301		654 troponin T, beta, skeletal	654	AGATTGACAA [G/A] TTCGAGTTTG	S	G	A	K	K
G4066u2	WIAF-13677	HT4301		774 troponin T, beta, skeletal	774	GCAAGTCGG [C/T] GGGGCGCTGGA	S	C	T	G	G
G4066u3	WIAF-13708	HT4301		625 troponin T, beta, skeletal	625	GGAGCTCTGG [G/C] AGACCCCTGCA	M	G	C	E	Q
G4080u1	WIAF-14142	HT1396		HSPG2, heparan sulfate	13130	GATTCTCCTC [G/A] GGCATCACAG	S	G	A	S	S
G4080u2	WIAF-14150	HT1396		HSPG2, heparan sulfate	10340	TTGAGTTCCA [C/T] TGTGCTGTGC	S	C	T	H	H
G4080u3	WIAF-14151	HT1396		HSPG2, heparan sulfate	12392	AATGCTATGA [T/C] AGCTCCCCAT	S	T	C	D	D
G4080u4	WIAF-14152	HT1396		HSPG2, heparan sulfate	3416	TGCTGTGCC [C/T] GAGGAAACCG	S	C	T	P	P
G4080u5	WIAF-14154	HT1396		HSPG2, heparan sulfate	4588	GTGCCGCTGG [T/C] GGCCAGCATC	M	T	C	V	A
G4080u6	WIAF-14156	HT1396		HSPG2, heparan sulfate	9582	GGACAGCCAC [G/A] CGGTGCTGCA	M	G	A	A	T
G4096u1	WIAF-13890	HT4237		394 motor protein	394	CRAAGAAATC [G/A] ATTCAAGTCGG	S	G	A	S	S
G4096u2	WIAF-13910	HT4237		455 motor protein	455	ATCTAAACAG [C/T] CTGCCTCACA	M	C	T	P	S
G4096u3	WIAF-13911	HT4237		1150 motor protein	1150	CTAAGGTTGT [A/G] TCTCAGTATC	S	A	G	V	V
G4109u1	WIAF-14034	HT28223		1238 phosphoglucosyltransferase-related protein	1238	TACAGCGTGG [C/T] GAAGACGGAT	M	C	T	A	V
G4109u2	WIAF-14035	HT28223		1043 phosphoglucosyltransferase-related protein	1043	ATTATTGCTG [C/A] CCGAAGCAG	M	C	A	A	D
G4112u1	WIAF-13615	HT4401		374 KIF5A, kinesin family member 5A	374	AGATGTCCTT [G/A] CTGGCTACAA	M	G	A	A	T
G4112u2	WIAF-13623	HT4401		2767 KIF5A, kinesin family member 5A	2767	AGAGAGTTAA [G/T] GCCCTGGAGG	M	G	T	K	N

2825.1027-001

G417lu2	WIAF-13689	HT1587	1036 KIF5B, kinesin family member 5B	TTAGTAAAC [T/C] GGAGCTGAAG	S	T	C	T	T
G4176u1	WIAF-14204	HT33754	TNR, tenascin R (restrictin, 130 janusin)	GCTCATTGGC [G/A] TCAACCTGAT	M	G	A	V	I
G4176u2	WIAF-14205	HT33754	TNR, tenascin R (restrictin, 463 janusin)	CTGTCCATGT [G/T] CCAGTTCAGC	M	G	T	A	S
G4176u3	WIAF-14206	HT33754	TNR, tenascin R (restrictin, 249 janusin)	ACTACAACAC [G/A] TCCAGCAAAG	S	G	A	T	T
G4176u4	WIAF-14208	HT33754	TNR, tenascin R (restrictin, 2009 janusin)	CTGTCCCCCA [G/A] GGGCATTGGT	M	G	A	R	K
G4176u5	WIAF-14209	HT33754	TNR, tenascin R (restrictin, 2175 janusin)	CAGCCTCCTC [G/A] GAGACCTCCA	S	G	A	S	S
G4176u6	WIAF-14210	HT33754	TNR, tenascin R (restrictin, 3318 janusin)	AATCCACCGA [C/T] GGAAGCCGCA	S	C	T	D	D
G4176u7	WIAF-14211	HT33754	TNR, tenascin R (restrictin, 3221 janusin)	CCGGCAAAAC [T/C] GACAGCCAGT	M	T	C	L	P
G4176u8	WIAF-14217	HT33754	TNR, tenascin R (restrictin, 1635 janusin)	TCTCGGACAC [C/T] GTGGCTTTTG	S	C	T	T	T
G4178u1	WIAF-14138	HT0224	2827 ACTN2, actinin, alpha 2	GCTGCTTCT [C/T] TTCGCACTC	M	C	T	S	F
G4178u2	WIAF-14139	HT0224	2818 ACTN2, actinin, alpha 2	CTGGATTACG [C/T] TGGTTCTCT	M	C	T	A	V
G418u1	WIAF-11750	L07594	TGFBR3, transforming growth factor, beta receptor III (betaglycan, 300kD)	GAGTGCACTT [C/T] CCTATCCCGC	S	C	T	F	F
G418u2	WIAF-11751	L07594	TGFBR3, transforming growth factor, beta receptor III (betaglycan, 300kD)	AGAAGACGTT [C/T] ACCAAGCCCC	S	C	T	F	F
G418u3	WIAF-11752	L07594	TGFBR3, transforming growth factor, beta receptor III (betaglycan, 300kD)	AATTTCCTCA [C/T] CAAITTTCCA	M	C	T	P	S
G418u4	WIAF-11771	L07594	TGFBR3, transforming growth factor, beta receptor III (betaglycan, 300kD)	TGTGTGAAT [G/T] TCACCTGTCA	S	G	T	L	L
G418u5	WIAF-11744	L07594	TGFBR3, transforming growth factor, beta receptor III (betaglycan, 300kD)	CTGATGAGCT [T/C] CTGTTTAGCC	M	T	C	F	S

004060" 22425360

G418u6	WIAF-11772	L07594	1470	TGFB3, transforming growth factor, beta receptor III (betaglycan, 300kD)	AGCTACGGAT [C/T] CTGCTGGACC	S	C	T	I	I
G418u7	WIAF-11773	L07594	1170	TGFB3, transforming growth factor, beta receptor III (betaglycan, 300kD)	TCTTGAAGTG [C/A] AAAAAAGTCTG	N	C	A	C	*
G418u8	WIAF-11745	L07594	1463	TGFB3, transforming growth factor, beta receptor III (betaglycan, 300kD)	CCTCCTGAGC [T/C] ACGGATCCTG	M	T	C	L	P
G418u9	WIAF-11746	L07594	2211	TGFB3, transforming growth factor, beta receptor III (betaglycan, 300kD)	ATGTTGAGGT [A/G] TCTGTTACTA	S	A	G	V	V
G418u1	WIAF-14207	HT2008	425	SPTBN1, spectrin, beta, non-erythrocytic 1	CTCTGCGCGG [C/T] TTTTGTAGCG	M	C	T	L	F
G418u2	WIAF-14213	HT2008	3565	SPTBN1, spectrin, beta, non-erythrocytic 1	AGACAGCGAT [C/T] GCCTCGGAGG	S	C	T	I	I
G418u3	WIAF-14218	HT2008	1258	SPTBN1, spectrin, beta, non-erythrocytic 1	ACCTTCTGGA [A/G] TGGATTGAAC	S	A	G	E	E
G418u4	WIAF-14219	HT2008	1780	SPTBN1, spectrin, beta, non-erythrocytic 1	AGCTCGAGGC [C/T] GAGAAATTACC	S	C	T	A	A
G418u5	WIAF-14220	HT2008	3637	SPTBN1, spectrin, beta, non-erythrocytic 1	ACATCAAGAA [T/C] GAGATCGACA	S	T	C	N	N
G4183u1	WIAF-13976	HT2640	404	TPM4, tropomyosin 4	CCAGCACAT [T/C] GCGGAAGAGG	S	T	C	I	I
G4185u1	WIAF-13554	HT3451	257	MFAP1, microfibrillar-associated protein 1	AAGCCACAGC [T/G] ATGCCCTTAT	M	T	G	Y	D
G4185u2	WIAF-13555	HT3451	1108	MFAP1, microfibrillar-associated protein 1	CCAACAAAGC [T/G] GTTAAGGGCA	S	T	G	A	A
G4185u3	WIAF-13570	HT3451	274	MFAP1, microfibrillar-associated protein 1	CTATGGAGTC [C/T] TCAGATGAGG	S	C	T	S	S
G4196u1	WIAF-13665	HT97558	941	NUP88, nucleoporin 88kD	GGGTCCATTG [C/A] CCATGCATCT	M	C	A	A	D
G4196u2	WIAF-13666	HT97558	1092	NUP88, nucleoporin 88kD	ATGACCACAC [G/A] TCAGAAAAGT	S	G	A	A	T
G4196u3	WIAF-13667	HT97558	1551	NUP88, nucleoporin 88kD	TCCATCCAGC [G/A] TCTCCTCCCC	S	G	A	A	A
G4196u4	WIAF-13668	HT97558	2220	NUP88, nucleoporin 88kD	AGGTGAACA [T/C] ATAGGGAAA	S	T	C	H	H
G4196u5	WIAF-13669	HT97558	2205	NUP88, nucleoporin 88kD	CCATCTGAA [A/G] GAGGAGGTG	S	A	G	K	K
G4208u1	WIAF-13921	HT1122	1329	VCL, vinculin	TGATCCTAAA [G/C] AAAGAGATGA	M	G	C	E	Q
G4208u2	WIAF-13922	HT1122	2438	VCL, vinculin	CCATCTCCC [A/G] ATGGTGATGG	S	A	G	P	P
G4208u3	WIAF-13941	HT1122	818	VCL, vinculin	GGGATGAAGA [T/C] GCCTGGGCCA	S	T	C	D	D
G4208u4	WIAF-13942	HT1122	1556	VCL, vinculin	AAGCACAGCG [G/A] TGGATTGATA	S	G	A	R	R

002060 2445960

G4213u1	WIAF-13605	HT2813	163	NUP153, nucleoporin 153kD	GCAGGGTGG [T/C] TACAAAGATA	S	T	C	L	L	L
G4213u2	WIAF-13606	HT2813	742	NUP153, nucleoporin 153kD	GAATCTTCA [A/G] TCCTTAAAC	M	A	G	I	V	
G4213u3	WIAF-13609	HT2813	1800	NUP153, nucleoporin 153kD	TTAGACCTGC [A/C] GAAATCCTGA	S	A	C	A	A	
G4213u4	WIAF-13627	HT2813	1829	NUP153, nucleoporin 153kD	AGTGTCTTAG [A/C] TATTCTGAAA	M	A	C	D	A	
G4213u5	WIAF-13632	HT2813	3258	NUP153, nucleoporin 153kD	CTTTGGCAA [C/T] GTGGAGCCTG	S	C	T	N	N	
G4213u6	WIAF-13635	HT2813	4162	NUP153, nucleoporin 153kD	CTCTGGAACA [A/G] CTCCTAATTC	M	A	G	T	A	
G4218u1	WIAF-13854	HT1681	1122	phosphatidylinositol glycan, class A	AACCTTATTA [T/C] TTTATGTGAG	M	T	C	I	T	
G4223u1	WIAF-14160	HT1684	1434	CD36L2, CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 2 (lysosomal integral membrane protein II)	ATTAGATGAC [T/C] TTGTTGAAAC	M	T	C	F	L	
G4223u2	WIAF-14173	HT1684	696	CD36L2, CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 2 (lysosomal integral membrane protein II)	GTGGTCCAG [G/A] TGCACCTTCT	M	G	A	V	M	
G4223u3	WIAF-14174	HT1684	986	CD36L2, CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 2 (lysosomal integral membrane protein II)	CAGACAGTG [C/T] AATATGATTA	S	C	T	C	C	
G4223u4	WIAF-14176	HT1684	1437	CD36L2, CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 2 (lysosomal integral membrane protein II)	AGATGACTTT [G/A] TTGAACGGG	M	G	A	V	I	
G4227u1	WIAF-14056	HT1929	912	proteoglycan 2	ATGCTCCAA [G/A] AAAGATGGG	S	G	A	K	K	
G4227u2	WIAF-14057	HT1929	1254	proteoglycan 2	GGAACTTTGC [G/A] TACTGGGCTG	S	G	A	A	A	
G4227u3	WIAF-14058	HT1929	1321	proteoglycan 2	CCGAGGAGGC [T/C] ACTGGCGTGC	M	T	C	Y	H	
G4229u1	WIAF-13961	HT1689	74	SDC4, syndecan 4 (amphiglycan, ryudocan)	GCTGCTGCTG [T/C] TCTTCGTAGG	M	T	C	F	L	
G4230u1	WIAF-13525	HT4995	602	TRAM protein	CCATAACCTG [A/C] TGACATTCA	M	A	C	M	L	
G4243u1	WIAF-14169	HT2901	406	KRT17, keratin 17	AGCTGGAGGT [G/A] AAGATCCGTG	S	G	A	V	V	
G4243u2	WIAF-14170	HT2901	478	KRT17, keratin 17	ACAGGACAAT [T/C] GAGGAGCTGC	S	T	C	I	I	
G4243u3	WIAF-14171	HT2901	389	KRT17, keratin 17	GGAGGAGGCC [A/G] ACACCTGAGCT	M	A	G	N	D	
G4243u4	WIAF-14178	HT2901	564	KRT17, keratin 17	CTGGCTGCTG [A/C] TGACTTCGGC	M	A	C	D	A	
G4244u1	WIAF-14086	HT1056	386	clathrin, light polypeptide a	ATCGATTGCA [G/C] TCAGAGCCTG	M	G	C	Q	H	
G4246u1	WIAF-14044	HT97492	259	SIN, sarcophilin	GTCTTATCAG [T/C] ACTGAGAGGC	M	T	C	Y	H	
G4246u2	WIAF-14045	HT97492	189	SIN, sarcophilin	ACACCCGGGA [G/A] CTGTTTCTCA	S	G	A	E	E	

002060-2412360

G4254u1	WIAF-13546	HT3393	86	TNNI2, troponin I, skeletal, fast	ACCTGAAGAG [C/T]GTGATGCTGC	S	C	T	S	S
G4254u2	WIAF-13553	HT3393	530	TNNI2, troponin I, skeletal, fast	TCGAGGAGAA [G/C]TCTGGCATGG	M	G	C	K	N
G4255u1	WIAF-13644	HT2907	562	CRYAB, crystallin, alpha B	AGTCCACAG [G/A]AAATACCGGA	S	G	A	R	R
G4255u2	WIAF-13645	HT2907	367	CRYAB, crystallin, alpha B	CCTCCTTCT [G/A]CGGGCACCCA	S	G	A	L	L
G4255u3	WIAF-13872	HT2907	271	CRYAB, crystallin, alpha B	CCAGCCGCCT [C/T]TTTGACCACT	S	C	T	L	L
G4255u4	WIAF-13873	HT2907	580	CRYAB, crystallin, alpha B	GGATCCCAGC [T/C]GATGTAGACC	S	T	C	A	A
G4257u1	WIAF-14052	HT1694	394	PIGF, phosphatidylinositol glycan, class F	TAGAGTTGSC [A/G]TTGGAACAT	S	A	G	A	A
G4257u2	WIAF-14053	HT1694	252	PIGF, phosphatidylinositol glycan, class F	TATTTAGTAG [T/C]GAAACCAAT	M	T	C	V	A
G4257u3	WIAF-14069	HT1694	291	PIGF, phosphatidylinositol glycan, class F	TCATTATCAC [A/G]CAAGTAACCT	M	A	G	H	R
G4264u1	WIAF-13519	HT0968	1720	TJP1, tight junction protein 1 (zona occludens 1)	CGGTCACTGG [C/T]TTCCAGCCAG	M	C	T	A	V
G4264u2	WIAF-13520	HT0968	2272	TJP1, tight junction protein 1 (zona occludens 1)	CATGCTGATG [A/G]TCACACACCT	M	A	G	D	G
G4264u3	WIAF-13529	HT0968	5408	TJP1, tight junction protein 1 (zona occludens 1)	AGCCTCCTGA [A/T]GCTGATGGTG	M	A	T	E	D
G434u1	WIAF-11748	M21121	286	SCYA5, small inducible cytokine A5 (RANTES)	TACATCAACT [C/T]TTTGGAGATG	M	C	T	S	F
G434u2	WIAF-11749	M21121	137	SCYA5, small inducible cytokine A5 (RANTES)	GCTTTGCCTA [C/T]ATTGCCCGCC	S	C	T	Y	Y
G435u1	WIAF-11741	M31933	754	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	GTCACCTGGA [T/C]TGCTGTAGCG	M	T	C	I	T
G435u2	WIAF-11743	M31933	395	FCGR2B, Fc fragment of IgG, low affinity IIB, receptor for (CD32)	GGGAGTACAC [G/A]TGCCAGACTG	S	G	A	T	T

00000000000000000000

G435u3	WIAF-11742	M31933	FCR2B, Fc fragment of IgG, low 673 affinity IIB, receptor for (CD32)		TACAGGCTGT [T/A] CTCATCCAAG	M	T	A	F	Y
G436u1	WIAF-13728	HT0900	GBE1, glucan (1,4-alpha-), branching enzyme 1 (glycogen branching enzyme, Andersen disease, glycogen storage disease type IV)	1176	TTACGTCCAT [G/A] CTTTATCATC	M	G	A	M	I
G436u2	WIAF-13729	HT0900	GBE1, glucan (1,4-alpha-), branching enzyme 1 (glycogen branching enzyme, Andersen disease, glycogen storage disease type IV)	1609	GAGTGTCTCTG [A/G] CTCCTTTTAC	M	A	G	T	A
G4373u1	WIAF-13559	HT0940	HSD17B2, hydroxysteroid (17-beta) dehydrogenase 2	1117	GCCAGCAAGG [A/T] CTTCTCTCCG	M	A	T	D	V
G4373u2	WIAF-13560	HT0940	HSD17B2, hydroxysteroid (17-beta) dehydrogenase 2	1195	CCAGGGAAG [G/A] CGCTTACTTG	M	G	A	G	D
G438u1	WIAF-11830	M63121	TNFRSF1A, tumor necrosis factor receptor superfamily, member 1A	583	ACCGTGTGTG [G/A] CTGAGGAAG	M	G	A	G	D
G438u2	WIAF-11790	M63121	TNFRSF1A, tumor necrosis factor receptor superfamily, member 1A	618	TTATTGGAGT [G/A] AAAACCTTTT	M	G	A	E	K
G440u1	WIAF-11806	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	261	TGCTAAAGCT [A/G] AGAGGGCTGC	S	A	G	L	L
G440u2	WIAF-11807	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	2089	CAGGCTGCAG [G/A] CAGTTCAGCG	M	G	A	A	T
G440u3	WIAF-11808	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	2155	TGCCCAGCTC [C/T] AGGAGGGACA	N	C	T	Q	*
G440u4	WIAF-11818	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	1789	GAACAACATT [G/A] CTTATGGGCT	M	G	A	A	T
G440u5	WIAF-11819	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	1565	AAGGGCTGA [C/T] GTTTACCCCTA	M	C	T	T	M
G440u6	WIAF-11820	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	1254	TGCATTGGG [G/T] GTGCAGATGC	S	G	T	G	G
G440u7	WIAF-11788	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	1231	GTACCTGCTC [A/G] TAAGAGGGGT	M	A	G	I	V
G440u8	WIAF-11821	M74447	TAP2, transporter 2, ABC (ATP binding cassette)	1404	TGCTCAGCAA [C/T] GTGGGAGCTG	S	C	T	N	N

0000002445960

G440u9	WIAF-11783	M74447	2187	TAP2, transporter 2, ABC (ATP binding cassette)	CCGCGCTGGT [T/G] CAGCAGCGGC	S	T	G	V
G440u10	WIAF-11786	M74447	1825	TAP2, transporter 2, ABC (ATP binding cassette)	TGATAAGGTG [A/G] TGGCGGCTGC	M	A	G	M
G440u1	WIAF-14007	HT97396	839	A33	GCCAATCAAA [G/T] GAGGGCTCAC	M	G	T	K
G440u1	WIAF-14013	HT1215	109	ACP2, acid phosphatase 2, lysosomal	CGCCCCACCC [G/A] GGCCCCGAGT	M	G	A	R
G440u2	WIAF-14016	HT1215	1271	ACP2, acid phosphatase 2, lysosomal	ACCGCCACGT [C/T] GCAGATGGGG	S	C	T	V
G440u1	WIAF-13661	HT3564	872	ACPP, acid phosphatase, prostate	ACAAAAAACT [T/C] ATCATGTATT	S	T	C	L
G440u2	WIAF-13662	HT3564	839	ACPP, acid phosphatase, prostate	ATCACATGAA [G/A] AGAGCAACTC	S	G	A	K
G440u3	WIAF-13881	HT3564	741	ACPP, acid phosphatase, prostate	AGAATTGTCA [G/T] AATTGTCCCT	N	G	T	E
G441u1	WIAF-10166	M77349	698	TGPBI, transforming growth factor, beta-induced, 68kD	GTGCCCCGCT [C/G] CTGAAAGCCG	S	C	G	L
G441u2	WIAF-10168	M77349	1028	TGPBI, transforming growth factor, beta-induced, 68kD	GGCTGTCTGT [A/G] GAGACCCCTGG	S	A	G	V
G441u3	WIAF-10169	M77349	1667	TGPBI, transforming growth factor, beta-induced, 68kD	ACACAGTCTT [T/C] GCTCCCAAA	S	T	C	F
G441u4	WIAF-10171	M77349	1463	TGPBI, transforming growth factor, beta-induced, 68kD	GTAATAGCCT [C/T] TGCATTGAGA	S	C	T	L
G441u1	WIAF-14005	HT97468	492	acyl-CoA	GCTGACCAAT [A/G] AGGCCACCCCT	M	A	G	K
G441u2	WIAF-14008	HT97468	1076	acyl-CoA	TGCCCGAGAC [C/T] GAGGACGAGA	S	C	T	T
G4412u1	WIAF-13576	HT1882	657	ACADS, acyl-Coenzyme A dehydrogenase, C-2 to C-3 short chain	GCAAAACAAG [G/A] GCATCAGTGC	M	G	A	G
G4412u2	WIAF-13579	HT1882	1022	chain	TGACCTGSGC [C/T] GCTGCCATGC	S	C	T	R
G4415u1	WIAF-14080	HT2503	2170	acyl-Coenzyme A:cholesterol acyltransferase	TCATTATATT [C/T] GAGCAGATTTC	S	C	T	F
G4415u2	WIAF-14081	HT2503	1993	acyl-Coenzyme A:cholesterol acyltransferase	TTTCAGTCC [C/T] TATTTCGT	S	C	T	P

G4415u3	WIAF-14098	HT2503	2006	acyl-Coenzyme A:cholesterol acyltransferase	TTTTCTGTTT [C/G] AACATTGGCG	M	C	G	Q	E
G4415u4	WIAF-14101	HT2503	2365	acyl-Coenzyme A:cholesterol acyltransferase	GGGGTTATGT [C/T] GCTATGAAGT	S	C	T	V	V
G4417u1	WIAF-13819	HT0542	356	AOAH, acyloxyacyl hydrolase (neutrophil)	TCCAGCCAAC [G/A] ATGACCAGTC	M	G	A	D	N
G4417u2	WIAF-13820	HT0542	340	AOAH, acyloxyacyl hydrolase (neutrophil)	TTCAGTCCTC [G/A] GCCTCTCCAG	S	G	A	S	S
G4417u3	WIAF-13824	HT0542	1595	AOAH, acyloxyacyl hydrolase (neutrophil)	GCTAAATAAA [G/A] ACATGACCTA	M	G	A	D	N
G4417u4	WIAF-13841	HT0542	382	AOAH, acyloxyacyl hydrolase (neutrophil)	CCAGCCTCTC [G/A] AATGGGCACA	S	G	A	S	S
G4417u5	WIAF-13842	HT0542	458	AOAH, acyloxyacyl hydrolase (neutrophil)	CAACTCGAAG [G/A] TCCAGGCCTC	M	G	A	V	I
G4417u6	WIAF-13843	HT0542	1201	AOAH, acyloxyacyl hydrolase (neutrophil)	GATTTCTGGA [C/T] TCCACTGTTG	S	C	T	D	D
G4417u7	WIAF-13844	HT0542	1321	AOAH, acyloxyacyl hydrolase (neutrophil)	ACCTGAAGAA [A/G] TTTATAGAAA	S	A	G	K	K
G4417u8	WIAF-13845	HT0542	1404	AOAH, acyloxyacyl hydrolase (neutrophil)	GATGCTGCA [G/A] TGGGAAGAGT	M	G	A	S	N
G4417u9	WIAF-13846	HT0542	1759	AOAH, acyloxyacyl hydrolase (neutrophil)	AATTTACAAA [C/T] TTCAATCTTT	S	C	T	N	N
G4417u10	WIAF-13847	HT0542	1644	AOAH, acyloxyacyl hydrolase (neutrophil)	CTCCAGGTCA [G/A] CCCCTGCCAC	M	G	A	S	N
G442u1	WIAF-11828	M94582	933	IL8RA, interleukin 8 receptor, alpha	CACATCGACC [G/A] GGCTCTGGAT	M	G	A	R	Q
G442u2	WIAF-11829	M94582	721	IL8RA, interleukin 8 receptor, alpha	TCATCGTGCC [A/G] CTGCTGATCA	S	A	G	P	P
G442u3	WIAF-11780	M94582	1027	IL8RA, interleukin 8 receptor, alpha	GCCATGGACT [C/T] CTCAGATTC	S	C	T	L	L
G442u4	WIAF-11792	M94582	78	IL8RA, interleukin 8 receptor, alpha	ATGGAGAGTG [A/G] CAGCTTTGAA	M	A	G	D	G
G4423u1	WIAF-13752	HT2216	71	ADSL, adenylosuccinate lyase	GCTATGCCAG [C/T] CCGAGAGTGT	S	C	T	S	S
G4423u2	WIAF-13794	HT2216	126	ADSL, adenylosuccinate lyase	ATGGCGGCAG [C/T] TGTGGCTGTG	S	C	T	L	L
G4423u3	WIAF-13795	HT2216	674	ADSL, adenylosuccinate lyase	AGCTTGACAA [G/A] ATGTGTACAG	S	G	A	K	K

G4428u1	WIAP-13954	HT97524	ADFP, adipose differentiation- 57 related protein; adipophilin	TGGTCAACCT [G/A] CCCTTGGTGA	S	G	A	L	L
G4434u1	WIAP-13506	HT0863	551 ARF3, ADP-ribosylation factor 3	TCTGGAGACA [C/T] TACTTCCAGA	S	C	T	H	H
G444u1	WIAP-10172	U28694	CCR3, chemokine (C-C motif) receptor 3	CGAGATCTTT [T/G] TCATAATCCT	M	T	G	F	V
G444u2	WIAP-10181	U28694	CCR3, chemokine (C-C motif) 214 receptor 3	TCCTCATAAA [A/G] TACAGGAGGC	S	A	G	K	K
G4440u1	WIAP-14054	HT1392	ADRBK1, adrenergic, beta, 136 receptor kinase 1	GCAAGAAGAT [A/C] CTGCTGCCCG	S	A	C	I	I
G445u1	WIAP-10183	U40373	Human cell surface glycoprotein CD44 mRNA, complete cds.	TAGAAGGGCA [C/T] GIGGTGATTC	S	C	T	H	H
G4456u1	WIAP-13629	HT0626	ALDOC, aldolase C, fructose- 796 bisphosphate	CCCTGCTCAA [G/A] CCCAACATGG	S	G	A	K	K
G446u1	WIAP-11832	U64198	IL12RB2, interleukin 12 receptor, beta 2	TGAAGCCTTC [C/G] CATGTAATTT	S	C	G	S	S
G446u2	WIAP-11795	U64198	IL12RB2, interleukin 12 receptor, beta 2	TTTCTCAAC [G/A] CATTACTTCC	S	G	A	T	T
G446u3	WIAP-11833	U64198	IL12RB2, interleukin 12 receptor, beta 2	TGCAAGGTAA [A/G] GCCAATTGGA	S	A	G	K	K
G446u4	WIAP-11835	U64198	IL12RB2, interleukin 12 receptor, beta 2	CTCCTCGCCA [G/C] GTCTCTGCAA	M	G	C	Q	H
G446u5	WIAP-11793	U64198	IL12RB2, interleukin 12 receptor, beta 2	GTGGAGCAGA [G/A] ATCTTCGTG	S	G	A	E	E
G446u6	WIAP-11794	U64198	IL12RB2, interleukin 12 receptor, beta 2	AGTTCCACAG [G/C] AAATGAGAGG	M	G	C	G	A
G446a7	WIAP-13128	U64198	IL12RB2, interleukin 12 receptor, beta 2	GGTGACTTGG [C/G] AGCCTCCAG	M	C	G	Q	E
G446a8	WIAP-13129	U64198	IL12RB2, interleukin 12 receptor, beta 2	TCTAAACTGG [C/G] TACGGAGTCG	M	C	G	L	V
G447u1	WIAP-11796	X03663	CSF1R, colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	CCAGTGTCCC [C/T] GAGCTGGTCC	S	C	T	P	P
G447u2	WIAP-11836	X03663	CSF1R, colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	ACAACAACAC [T/C] AAGCTGGCAA	S	T	C	T	T

G447u3	WIAF-11837	X03663	886	CSF1R, colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	GCTGAAAAGTG [C/A] AGAAAGTCAT	M	C	A	Q	K
G447u4	WIAF-11797	X03663	2425	CSF1R, colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	GAAGAAATAT [G/A] TCCGCAGGGA	M	G	A	V	I
G4473u1	WIAF-13904	HT1352	860	FUC1, fucosidase, alpha-L-1, tissue	TTCAAGCCAC [A/G] GAGCTTGCCA	M	A	G	Q	R
G4473u2	WIAF-13916	HT1352	440	FUC1, fucosidase, alpha-L-1, tissue	ACAAACTGGC [C/T] GAGTCCTGTG	M	C	T	P	L
G4479u1	WIAF-13637	HT1995	2465	AMPD2, adenosine monophosphate deaminase 2 (isoform L)	GCTCAATGA [G/T] CCTGGTCCAT	-	G	T	-	-
G4479u2	WIAF-13866	HT1995	1258	AMPD2, adenosine monophosphate deaminase 2 (isoform L)	TGGATGTGA [T/C] GCGACAGGA	S	T	C	H	H
G4479u3	WIAF-13867	HT1995	1280	AMPD2, adenosine monophosphate deaminase 2 (isoform L)	CAC TTTCAT [C/T] GCTTGACAA	M	C	T	R	C
G4479u4	WIAF-13868	HT1995	1201	AMPD2, adenosine monophosphate deaminase 2 (isoform L)	TGCGGGAGGT [C/T] TTTGAGAGCA	S	C	T	V	V
G4479u5	WIAF-13869	HT1995	1579	AMPD2, adenosine monophosphate deaminase 2 (isoform L)	GTACCAAGGG [C/T] CAGCTGGCCA	S	C	T	G	G
G4492u1	WIAF-14084	HT3390	866	ANX11, annexin XI (56kD autoantigen)	CCTGGGGAGT [C/T] GCTCCAACAA	M	C	T	R	C
G4492u2	WIAF-14085	HT3390	850	ANX11, annexin XI (56kD autoantigen)	AGGCCATCAT [T/C] GACTGCCTGG	S	T	C	I	I
G450u1	WIAF-10170	X85740	1196	CCR4, chemokine (C-C motif) receptor 4	TCCAAATTTA [C/T] TCTGCTGACA	S	C	T	Y	Y
G4502u1	WIAF-13510	HT4840	165	ASS, argininosuccinate synthetase	AGGCTATGA [C/T] GTCATTGCCT	S	C	T	D	D
G4502u2	WIAF-13511	HT4840	369	ASS, argininosuccinate synthetase	GGCCCTGCAT [C/T] GCCCGCAAAC	S	C	T	I	I
G4502u3	WIAF-13512	HT4840	73	ASS, argininosuccinate synthetase	AAFTCCAGAC [G/A] CTAFTGCCAG	-	G	A	-	-

G4502u4	WIAF-13513	HT4840	129 ASS, argininosuccinate synthetase	TGGACACCTC[G/C]TGCATCCTCG	S	G	C	S	S
G4502u5	WIAF-13514	HT4840	285 ASS, argininosuccinate synthetase	AGTTTGTGGA[G/A]GAGTTCACTCT	S	G	A	E	E
G4502u6	WIAF-13515	HT4840	234 ASS, argininosuccinate synthetase	AGGCACGTGAA[G/A]CTTGGGGCCA	S	G	A	K	K
G4502u7	WIAF-13516	HT4840	316 ASS, argininosuccinate synthetase	CCAGTCCAGC[G/A]CACTGTATGA	M	G	A	A	T
G4502u8	WIAF-13537	HT4840	426 ASS, argininosuccinate synthetase	TGTCCTCCACGG[C/T]GCCACAGGAA	S	C	T	G	G
G4502u9	WIAF-13538	HT4840	530 ASS, argininosuccinate synthetase	GAATTCCTACA[A/G]CCGGTTCAAG	M	A	G	N	S
G4502u10	WIAF-13539	HT4840	750 ASS, argininosuccinate synthetase	TTCTCGAGAT[C/T]GAGTTCAAAA	S	C	T	I	I
G4502u11	WIAF-13540	HT4840	960 ASS, argininosuccinate synthetase	ATGCTCATTT[A/G]GACATCGAGG	S	A	G	L	L
G4508u1	WIAF-13663	HT28557	1767 ARSD, arylsulfatase D	CAGTTTTCCTCA[T/C]GAGCAACATC	M	T	C	M	T
G4508u2	WIAF-13693	HT28557	433 ARSD, arylsulfatase D	TTCACTGGAA[C/T]GCAGGCTCAG	S	C	T	N	N
G4508u3	WIAF-13694	HT28557	747 ARSD, arylsulfatase D	GGTTTCTTCT[C/G]TGCTCTCCGG	M	C	G	S	C
G4508u4	WIAF-13696	HT28557	1012 ARSD, arylsulfatase D	CCACGAGTGC[A/G]TTCCTGGGGA	S	A	G	A	A
G4508u5	WIAF-13697	HT28557	1302 ARSD, arylsulfatase D	CGAGTGAATG[G/A]AGAGCCACAG	M	G	A	G	E
G4508u6	WIAF-13698	HT28557	1285 ARSD, arylsulfatase D	GGTGTCTCCC[G/A]GCCGGCCGAG	S	G	A	P	P
G4508u7	WIAF-13699	HT28557	1807 ARSD, arylsulfatase D	AGCCGTGCTG[C/T]GGACATTTCC	S	C	T	C	C
G4508u8	WIAF-13718	HT28557	483 ARSD, arylsulfatase D	GCAAGAATCT[T/C]GCAGCAGCAT	M	T	C	L	S
G4518u1	WIAF-13809	HT3430	ASPA, aspartoacylase 515 (aminoacylase 2, Canavan disease)	ACAACACCAC[C/T]TCTAACATGG	S	C	T	T	T
G4518u2	WIAF-13810	HT3430	ASPA, aspartoacylase 851 (aminoacylase 2, Canavan disease)	AAGTTGATTA[C/T]CCCCGGGATG	S	C	T	Y	Y
G4518u3	WIAF-13811	HT3430	ASPA, aspartoacylase 787 (aminoacylase 2, Canavan disease)	CATCATTTCA[A/G]TGAAGGAAAA	M	A	G	N	S
G4518u4	WIAF-13837	HT3430	ASPA, aspartoacylase 618 (aminoacylase 2, Canavan disease)	ACCTGTCTAC[G/A]TTTATCTGAT	M	G	A	V	I
G452a1	WIAF-10509	HT0695	553 APOA4, apolipoprotein A-IV	ACCAGGTCA[A/G]CACGAGGCC	M	A	G	N	S
G452a2	WIAF-13124	HT0695	563 APOA4, apolipoprotein A-IV	ACACGAGGC[C/T]GAGCAGCTGC	S	C	T	A	A

G4524u1	WIAF-14120	HT1541			ATP5A1, ATP synthase, H ⁺ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle	726	CTCAATTGCT [A/G] TTGACACAAT	M	A	G	I	V
G4524u2	WIAF-14131	HT1541			ATP5A1, ATP synthase, H ⁺ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle	153	ATCTTTTCATT [G/T] CTGCAAGGAA	M	G	T	A	S
G4526u1	WIAF-14130	HT4994			ATP5D, ATP synthase, H ⁺ transporting, mitochondrial F1 complex, delta subunit	400	TCCATCGCAG [T/C] GAACGCCGAC	M	T	C	V	A
G453u1	WIAF-10138	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	1747	CTGCCGCCCA [C/T] GCTGCTGGG	M	C	T	T	M
G453u2	WIAF-10147	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	2957	TTTTCCTTT [A/G] AAGTGGATGG	S	A	G	L	L
G453u3	WIAF-10148	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	3608	AGCCGGAGCC [A/G] GAGCTGGAAC	S	A	G	P	P
G453u4	WIAF-10149	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	457	CAGGGCCTGG [T/G] CGTCACACCC	M	T	G	V	G
G453u5	WIAF-10151	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	1505	AGCTGACACT [G/C] GTTCGCGTGA	S	G	C	L	L
G453u6	WIAF-10153	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	3446	ACCCAAACC [C/T] GAGTTGCTG	S	C	T	P	P
G453u7	WIAF-10161	HT0768			PDGFRB, platelet-derived growth factor receptor, beta polypeptide	2030	TTTGGCAGAA [G/A] AAGCCACGTT	S	G	A	K	K
G4533u1	WIAF-13616	HT1618			ATP synthase, H ⁺ transporting, subunit D, vacuolar	343	GTTACATGAT [C/T] GACAACGTGA	S	C	T	I	I
G4534u1	WIAF-13569	HT3556			ATP6E, ATPase, H ⁺ transporting, lysosomal (vacuolar proton pump) 31kD	654	TAAAGGTTTC [C/T] AACACCCTGG	S	C	T	S	S

G4574u3	WIAP-13807	HT0198	726	beta-1,4 N-acetylgalactosaminyltransferase	AACTACAACCT [G/T]GTCACCTTACA	S	G	T	L	L
G4574u4	WIAP-13836	HT0198	559	beta-1,4 N-acetylgalactosaminyltransferase	AGGGCTCAGC [C/A]TTCAGGCAGC	M	C	A	L	I
G4575u1	WIAP-13626	HT0341	1251	GCNT1, glucosaminyl (N-acetyl) transferase 1, core 2 (beta-1,6-N-acetylglucosaminyltransferase)	AGTATGATCT [A/G]TCTGACATGC	S	A	G	L	L
G4577u1	WIAP-13971	HT1495	1268	SIAT1, sialyltransferase 1 (beta-galactoside alpha-2,6-sialyltransferase)	ATTTCITTTAA [C/T]AACTACAAGA	S	C	T	N	N
G458u1	WIAP-10063	HT2968	1464	ALB, albumin	GTGAGAAGA [C/A]TATCTATCCG	M	C	A	D	E
G458u2	WIAP-10089	HT2968	1470	ALB, albumin	AGACTATCT [A/C]TCCGTGGTCC	S	A	C	L	L
G458u3	WIAP-10091	HT2968	1707	ALB, albumin	TTGTTGAGCT [C/T]GTGAACACA	S	C	T	L	L
G458a4	WIAP-10504	HT2968	889	ALB, albumin	CAGGGCGGAC [C/T]TTGCCAAGTA	M	C	T	L	F
G458a5	WIAP-10508	HT2968	1475	ALB, albumin	TATCTATCCG [T/A]GGTCTGTAAC	M	T	A	V	E
G458a6	WIAP-12091	HT2968	1330	ALB, albumin	CCAGAAATGCG [C/T]TATTAGTTTCG	S	C	T	L	L
G458a7	WIAP-12092	HT2968	1408	ALB, albumin	CCTAGGAAAA [G/a]TGGCAGCAA	M	G	a	V	M
G4592u1	WIAP-14126	HT2128	985	branched-chain keto acid dehydrogenase E1, alpha polypeptide	ACCAGCCCTT [T/C]CTCATCGAGG	S	T	C	F	F
G4593u1	WIAP-13574	HT97373	1743	BARD1, BRCA1 associated RING domain 1	GCTAGCCACT [G/C]CTCAGTAATG	M	G	C	C	S
G4593u2	WIAP-13592	HT97373	1167	BARD1, BRCA1 associated RING domain 1	TGTTCTTTAC [C/T]ACCTTCATGC	M	C	T	P	L
G4593u3	WIAP-13593	HT97373	1591	BARD1, BRCA1 associated RING domain 1	AGAATGGGCA [C/T]GTGGATATAG	S	C	T	H	H
G4593u4	WIAP-13594	HT97373	2030	BARD1, BRCA1 associated RING domain 1	AAAGTATGAA [A/G]TTCCTGAAGG	M	A	G	I	V
G4593u5	WIAP-13595	HT97373	2006	BARD1, BRCA1 associated RING domain 1	AAGAAAAGTA [T/C]GTGAACAGGA	M	T	C	C	R
G4599u1	WIAP-13920	HT4273	1803	CDH13, cadherin 13, H-cadherin (heart)	TCGTACCCGA [C/T]GTCTCCTACG	S	C	T	D	D
G4614u1	WIAP-13733	HT4835	91	S100A3, S100 calcium-binding protein A3	AGGATGGCCA [G/A]GCCTCTGGAG	M	G	A	R	K
G4614u2	WIAP-13734	HT4835	203	S100A3, S100 calcium-binding protein A3	TGCTGCAGAA [G/A]GAGCTGGCCA	S	G	A	K	K
G4614u3	WIAP-13769	HT4835	344	S100A3, S100 calcium-binding protein A3	TCTACTGCCA [C/T]GAGTACTTCA	S	C	T	H	H

000000-2246960

G467a5	WIAF-13308	X72861	ADRB3, adrenergic, beta-3-, receptor	1899	TCTGTTGATC[A/C]GAACCTGTGG	-	A	C	-	-
G467lu1	WIAF-13956	HT1925	NDUFB7, NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7 (18kD, B18)	161	TGGTGGCCAC[A/G]CAGCAGGAGA	S	A	G	T	T
G4673u1	WIAF-13889	HT0191	1349 CDC25A, cell division cycle 25A	1349	TCTGGGGCCA[G/C]CCCCAAAGAG	M	G	C	S	T
G4674u1	WIAF-13821	HT1393	261 CDC25B, cell division cycle 25B	261	ACGACCTCGC[C/T]GGGCTGGCA	S	C	T	A	A
G4674u2	WIAF-13822	HT1393	1297 CDC25B, cell division cycle 25B	1297	GATGTGGCC[C/T]TATTGACGGG	S	C	T	L	L
G4674u3	WIAF-13823	HT1393	1083 CDC25B, cell division cycle 25B	1083	ATAAGCGGAG[G/A]CGGAGCCTGA	S	G	A	R	R
G4674u4	WIAF-13827	HT1393	1446 CDC25B, cell division cycle 25B	1446	AGAGCCCCAT[C/T]GCGCCCTGTA	S	C	T	I	I
G468a1	WIAF-13309	L37019	ASIP, agouti (mouse)-signaling 192 protein	192	AAATCCAAAC[C/A]GATCGGCAGA	M	C	A	P	Q
G469lu1	WIAF-13753	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	179	TATAGCCTGA[T/A]TTTTGTGTGG	M	T	A	I	N
G469lu2	WIAF-13754	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	134	AAGGATGCAG[T/C]GGTGTCTCTT	M	T	C	V	A
G469lu3	WIAF-13755	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	193	TGTGTTGGGC[C/T]TCAGCGGGAA	M	C	T	L	F
G469lu4	WIAF-13756	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	770	AAAATAGCTG[C/T]AGCCTTGGTG	M	C	T	A	V
G469lu5	WIAF-13759	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	1130	TCTCAGAACT[A/C]CCCTAACCAAG	M	A	C	Y	S
G469lu6	WIAF-13796	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	482	AGGCTGAGGA[C/A]CCGGGCCAAG	M	C	A	T	N
G469lu7	WIAF-13797	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	259	GATGTTGAG[A/G]TCTATCTGCT	M	A	G	I	V
G469lu8	WIAF-13798	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	434	ATGAGCCTGG[A/G]CAAGTACCTG	M	A	G	D	G
G469lu9	WIAF-13799	HT97602	CMKBR9, chemokine (C-C motif) receptor 9	755	CAGGGCCGGG[C/T]TTTAAAAATA	M	C	T	A	V
G4699u1	WIAF-14040	HT4277	BAAT, bile acid Coenzyme A: amino acid N-acyltransferase (glycine N- choloyltransferase)	1426	TTCCAGATGT[G/T]ACCAGTCAAC	S	G	T	V	V

G4726u1	WIAF-14128	HT48614	AOC3, amine oxidase, copper containing 3 (vascular protein 1)	1606	TCCACCCAG [T/C] GGGGCCATAG	S	T	C	S	S
G4726u2	WIAF-14129	HT48614	AOC3, amine oxidase, copper containing 3 (vascular protein 1)	2242	TTCTAACAC [A/G] GTGACTGTGG	S	A	G	T	T
G4726u3	WIAF-14141	HT48614	AOC3, amine oxidase, copper containing 3 (vascular protein 1)	659	CCTGCCTAT [C/T] ACCGACGCC	M	C	T	H	Y
G4744u1	WIAF-13683	HT2599	CTH, cystathionase (cystathionine gamma-lyase)	564	ATATTGTCCA [T/C] AACATGGAG	S	T	C	H	H
G4748u1	WIAF-14144	HT1061	CYBA, cytochrome b-245, alpha polypeptide	242	GGGACAGAAG [C/T] ACATGACGC	M	C	T	H	Y
G4748u2	WIAF-14145	HT1061	CYBA, cytochrome b-245, alpha polypeptide	265	TGCTGAAGCT [G/C] TTCGGGCCT	S	G	C	L	L
G4750u1	WIAF-14116	HT48417	CYB5, cytochrome b-5	156	TGAAGTACTA [C/T] ACCCTAGAG	S	C	T	Y	Y
G4751u1	WIAF-13770	HT1285	UQCRC2, ubiquinol-cytochrome c reductase core protein II	495	AGAATTTTCT [C/A] GTTGGGAAGT	M	C	A	R	S
G4788u1	WIAF-13931	HT28249	1864 DSC3, desmocollin 3	1864	CTGTGATCC [T/C] GATGAACCTG	S	T	C	P	P
G4788u2	WIAF-13933	HT28249	2000 DSC3, desmocollin 3	2000	TGGATTTCAA [G/T] AATATACCAT	N	G	T	E	*
G4788u3	WIAF-13945	HT28249	2524 DSC3, desmocollin 3	2524	ACACTTACTC [G/A] GAGTGGCACA	S	G	A	S	S
G479u1	WIAF-12567	U36310	GPD2, glycerol-3-phosphate dehydrogenase 2 (mitochondrial)	894	GGGAAACTGC [A/G] TGTGAGCGC	M	A	G	H	R
G479u2	WIAF-12574	U36310	GPD2, glycerol-3-phosphate dehydrogenase 2 (mitochondrial)	1657	CTGGCAAAAG [G/T] TGGCCTATTG	M	G	T	R	S
G479u3	WIAF-12575	U36310	GPD2, glycerol-3-phosphate dehydrogenase 2 (mitochondrial)	1131	GTTATTTTCT [T/C] CTTACCTGG	M	T	C	F	S
G480u1	WIAF-12175	HT336	GRB2, growth factor receptor- bound protein 2	250	AATGAAACCA [C/A] ATCCCTGGTT	M	C	A	H	N
G4819u1	WIAF-13985	HT97576	EYA1, eyes absent (Drosophila) homolog 1	1804	CCCTGCACCA [T/C] GCCTTGGAAC	S	T	C	H	H
G482u1	WIAF-12181	J04501	GYS1, glycogen synthase 1 (muscle)	1186	CTGACGTCTT [T/C] CTGAGGCAT	S	T	C	F	F

G482u2	WIAF-12195	J04501	Gyl, glycogen synthase 1 (muscle)	1406		CCTTCCCGAC [A/G] TGAACAAGAT	M	A	G	M	V
G4827u1	WIAF-14177	HT97477	58 elongation			CGAGCTGGCC [A/G] TGATGGTAT	M	A	G	H	R
G483a1	WIAF-12113	HT4341	1850 GSY2			TTACCAGCAT [G/T] CCAGACACCT	M	G	T	A	S
G483u2	WIAF-12148	HT4341	1130 GSY2			GTTTTCATT [A/C] TGCTTGCCAA	M	A	C	M	L
G483u3	WIAF-12149	HT4341	880 GSY2			GCTTGAATGT [T/G] AAGAAATTTT	S	T	G	V	V
G483u4	WIAF-12150	HT4341	1115 GSY2			CATCAGATG [G/A] TGCTGTTTT	M	G	A	V	M
G483u5	WIAF-12156	HT4341	1230 GSY2			GAAAAGTTTG [G/A] AAAAAACTC	M	G	A	G	E
G483u6	WIAF-12159	HT4341	2033 GSY2			TGAGAGATAC [G/A] ATGAGGAAGA	M	G	A	D	N
G483u7	WIAF-12160	HT4341	1836 GSY2			TACTTAGGCA [G/C] ATATTACCAG	M	G	C	R	T
G483u8	WIAF-12161	HT4341	1678 GSY2			CTTACGGTAT [T/C] TACATCGTTG	S	T	C	I	I
G483u9	WIAF-12177	HT4341	790 GSY2			GGCTCAGT [G/C] TTCACCACGG	S	G	C	V	V
G483u10	WIAF-12188	HT4341	1728 GSY2			TGCAATCAGC [T/C] GACTAAGTTT	M	T	C	L	P
G484u1	WIAF-12151	HT5111	487 GSY3			CATCAAAGTG [A/G] TTGGCAATGG	M	A	G	I	V
G484u2	WIAF-12187	HT5111	1141 GSY3			AACCCGGGAA [C/T] AAATCCGAGA	N	C	T	Q	*
G489u1	WIAF-12152	HT2607	IRS1, insulin receptor substrate 1181			AAGAATGGC [G/A] GCACAAGTCG	M	G	A	R	Q
G489u2	WIAF-12184	HT2607	IRS1, insulin receptor substrate 1031			ATGGCGAGCC [C/T] TCCGGAGAGC	M	C	T	P	L
G492a1	WIAF-13345	L08603	307 MC4R, melanocortin 4 receptor			AGAAACCATT [A/G] TCATCACCCCT	M	A	G	I	V
G493u1	WIAF-12154	X67594	MC1R, melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)	346		CGCGCTGGTG [G/T] TGGCCACCCT	M	G	T	V	L
G493u2	WIAF-12167	X67594	MC1R, melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)	646		GACCTGCGC [C/T] GGGCGCGGCA	M	C	T	R	W
G493u3	WIAF-12170	X67594	MC1R, melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)	1110		AGGTGCTGAC [A/G] TGCTCTGGT	S	A	G	T	T
G493u4	WIAF-12186	X67594	MC1R, melanocortin 1 receptor (alpha melanocyte stimulating hormone receptor)	442		CGGAGCAAC [G/T] TGCTGGAGAC	M	G	T	V	L
G498u1	WIAF-11809	J04127	CYP19, cytochrome P450, subfamily XIX (aromatization of androgens)	1305		CTTATAGGTA [C/T] TTTCAGCCAT	S	C	T	Y	Y

20250624 22:59:56

G498u2	WIAF-11810	J04127		CYP19, cytochrome P450, subfamily CYP19, cytochrome P450, subfamily 1377 XIX (aromatization of androgens)	TGAAAGCCAT [C/T] CTCGTACAC	S	C	T	I	I
G498u3	WIAF-11811	J04127		CYP19, cytochrome P450, subfamily CYP19, cytochrome P450, subfamily 1406 XIX (aromatization of androgens)	CGATTCCACG [T/C] GAAGACATTG	M	T	C	V	A
G498u4	WIAF-11838	J04127		CYP19, cytochrome P450, subfamily CYP19, cytochrome P450, subfamily 1055 XIX (aromatization of androgens)	ATTGGTGAGA [G/A] AGACATAAAG	M	G	A	R	K
G498u5	WIAF-11800	J04127		CYP19, cytochrome P450, subfamily CYP19, cytochrome P450, subfamily 1001 XIX (aromatization of androgens)	ATTGCAAAGC [A/G] CCTAATGTT	M	A	G	H	R
G499u1	WIAF-11785	HT1439		2142 ESRL, estrogen receptor 1	TCCCTGCCAC [A/G] GTCGTAGAGC	S	A	G	T	T
G499u2	WIAF-11801	HT1439		443 ESRL, estrogen receptor 1	CCCTTGAACC [G/A] TCGCAGCTC	M	G	A	R	H
G500u1	WIAF-11803	X99101		793 ESRL, estrogen receptor 1	CATGATCAGC [T/C] GGGCCAGAA	M	T	C	W	R
G500u2	WIAF-11816	X99101		489 ESRL, estrogen receptor 1	GGAAAGTTTA [C/T] GAAGTGGAA	S	C	T	Y	Y
G500u3	WIAF-11817	X99101		474 ESRL, estrogen receptor 1	AGGCTGCCG [A/G] CTTGGAGT	S	A	G	R	R
G505u1	WIAF-11824	HT1113		1063 PRLR, prolactin receptor	GCTTTGAAGG [G/A] CTATAGCATG	M	G	A	G	D
G505u2	WIAF-11827	HT1113		2083 PRLR, prolactin receptor	GCAACATCAA [G/A] CAAGTGCAGG	M	G	A	S	N
G505u3	WIAF-11787	HT1113		582 PRLR, prolactin receptor	GAGGACATAC [A/G] TCATGATGTT	M	A	G	I	V
G505u4	WIAF-11802	HT1113		792 PRLR, prolactin receptor	CCTGTATGAA [A/C] TTCGATTAAA	M	A	C	I	L
G509u1	WIAF-11789	M32313		SPD5A1, steroid-5-alpha- reductase, alpha polypeptide 1 (3- oxo-5 alpha-steroid delta 4- dehydrogenase alpha 1)	CACTGTGGC [A/G] TGTACAATGG	S	A	G	A	A
G510a1	WIAF-13348	U17280		STAR, steroidogenic acute regulatory protein	CCAAATGTCAA [G/A] GAGATCAAGG	S	G	A	K	K
G520u1	WIAF-10224	HT0488		1139 inhibin, beta B	CCAAATGAT [T/C] GTGGAGGAGT	S	T	C	I	I
G520u1	WIAF-13507	D31770		ACVR2, activin A receptor, type II	CTTATTTCCTCC [G/A] GAGATGGAAG	S	G	A	P	P
G520u2	WIAF-13532	D31770		ACVR2, activin A receptor, type II	CAGCTTGCAAT [T/G] GCTGACTTTG	M	T	G	I	M
G520u3	WIAF-13533	D31770		ACVR2, activin A receptor, type II	CTGACTTTGG [G/C] TTGGCCTTAA	S	G	C	G	G
G520u4	WIAF-13534	D31770		ACVR2, activin A receptor, type II	TCCTTTGGAA [T/C] GAACGTGTGC	S	T	C	N	N
G523u1	WIAF-12155	HT4996		538 OXTR, oxytocin receptor	TGAGCGGAA [C/T] CGGTGTGTC	S	C	T	N	N

002000" 22445960

G523u2	WIAF-12180	HT4996	1057 OXTR, oxytocin receptor	TCTGCGAGAA [C/T] TTGCGGCTCA	S	C	T	N	N
G524a1	WIAF-13349	L05144	PK1, phosphoenolpyruvate carboxykinase 1 (soluble)	TGGACAGCCT [G/A] CCCAGGCAG	S	G	A	L	L
G528u1	WIAF-11831	V00572	988 PGK1, phosphoglycerate kinase 1	AAGCCACTGT [G/C] GCTTCTGGCA	S	G	C	V	V
G53u1	WIAF-10307	HT0508	723 DNA repair protein XRCC1	CCAGCGACCC [G/A] GCAGGACCTA	S	G	A	P	P
G53u2	WIAF-10308	HT0508	746 DNA repair protein XRCC1	TATGAGCTGT [C/T] TACCCCTCCAG	M	C	T	A	V
G53u3	WIAF-10309	HT0508	1884 DNA repair protein XRCC1	GGATCCCAAC [G/A] CGTTCGATG	S	G	C	T	S
G53u4	WIAF-10362	HT0508	425 DNA repair protein XRCC1	AACCCCAACC [G/A] CGTTCGATG	M	G	A	R	H
G534a1	WIAF-13310	U28281	1284 SCTR, secretin receptor	GCTTCCTCAA [T/C] GGGAGGTGC	S	T	C	N	N
G534a2	WIAF-13311	U28281	1404 SCTR, secretin receptor	AGCAGAGCCA [G/A] GCACCTGCA	S	G	A	Q	Q
G535u1	WIAF-12157	HT5001	1158 SHC1	ATGCTCTTGG [G/C] GTGCCCTCCAC	S	G	C	R	R
G535u2	WIAF-12196	HT5001	774 SHC1	ATGAGGAGGA [G/A] GAAGGCCAC	S	G	A	E	E
G536u1	WIAF-13923	M20747	SLC2A4, solute carrier family 2 (facilitated glucose transporter), member 4	GCCTGGCCAA [C/T] GCTGTGCCT	S	C	T	N	N
G538u1	WIAF-11812	M55531	SLC2A5, solute carrier family 2 (facilitated glucose transporter), member 5	GCAGCAGAGT [C/T] GCCACATCAT	S	C	T	V	V
G538u2	WIAF-11813	M55531	SLC2A5, solute carrier family 2 (facilitated glucose transporter), member 5	GACGCTTGTTG [C/T] TTGCCCTGGC	M	C	T	L	F
G538u3	WIAF-11791	M55531	SLC2A5, solute carrier family 2 (facilitated glucose transporter), member 5	ACAGGGAGGT [G/A] GCCGAGATCC	S	G	A	V	V
G539u1	WIAF-12158	K03195	Human (HepG2) glucose transporter gene mRNA, complete cds.	TCATGCTGGC [T/C] GTGGGAGAG	S	T	C	A	A
G539u2	WIAF-12191	K03195	Human (HepG2) glucose transporter gene mRNA, complete cds.	CCATCGCGCT [A/G] GCACGTGTGG	S	A	G	L	L
G540a1	WIAF-12114	HT960	1100 SOS1	AGTAAAGATC [A/C] AGAAGACAAG	M	A	C	Q	P
G540u2	WIAF-12165	HT960	933 SOS1	ATGATCGTTT [C/T] CTTAGTCAGT	S	C	T	F	F
G540u3	WIAF-12178	HT960	399 SOS1	TAGTAGCAGT [C/T] TTAGAATACA	S	C	T	V	V
G540u4	WIAF-12193	HT960	195 SOS1	CTCAGCCCG [A/C] AGTGCTTCAG	S	A	C	R	R
G540u5	WIAF-12197	HT960	1329 SOS1	GTTCGTAATGA [A/G] TTTATATATGG	S	A	G	E	E
G540u6	WIAF-12198	HT960	1339 SOS1	ATTTATATATG [G/A] AAGGAACCTCT	M	G	A	E	K
G543a1	WIAF-13312	J00306	1373 SST, somatostatin	AAGCAGGAAC [T/C] GCCCAAGTAC	M	T	C	L	P

G543a2	WIAF-13313	J00306	1603 SST, somatostatin	AGTATTGTCC [A/G] TATCAGACCT	-	A	G	-
G544u1	WIAF-12174	HT27489	SUR, sulfonylurea receptor (hyperinsulinemia)	CCATTGACAT [G/C] GCCACGGAAA	M	G	C	M I
G546u1	WIAF-13618	HT225	TKT, transketolase (Wernicke-Korsakoff syndrome)	GCTACATTGC [C/T] GAGAGAACA	S	C	T	A A
G551u1	WIAF-11709	HT1118	TNFRSF1B, tumor necrosis factor receptor superfamily, member 1B	GCTGCAGCAA [A/G] TGCTCGCCGG	S	A	G	K K
G551u2	WIAF-11710	HT1118	TNFRSF1B, tumor necrosis factor receptor superfamily, member 1B	TTCTGCACCTG [C/T] AGCCCGGCT	S	C	T	C C
G551u3	WIAF-11719	HT1118	TNFRSF1B, tumor necrosis factor receptor superfamily, member 1B	GATCTGTAAC [G/A] TGTGTGCCAT	M	G	A	V M
G551u4	WIAF-11673	HT1118	TNFRSF1B, tumor necrosis factor receptor superfamily, member 1B	AATGCAAGCA [T/G] GGATGCAGTC	M	T	G	M R
G551u5	WIAF-11720	HT1118	TNFRSF1B, tumor necrosis factor receptor superfamily, member 1B	CCAAGCACCT [C/T] CTTCTGTCTC	M	C	T	S F
G552u1	WIAF-12229	HT5108	TRAP3	GCGCTGCCC [G/A] CTCATGTGA	S	G	A	P P
G555u1	WIAF-12211	U94592	UCP2, uncoupling protein 2 (mitochondrial, proton carrier)	CGCGCTACAG [T/C] CAGGCCCCAG	M	T	C	V A
G556u1	WIAF-11804	AF001787	UCP2, uncoupling protein 2 (mitochondrial, proton carrier)	TCGGCCCTCTA [T/C] GACTCCGTCA	S	T	C	Y Y
G556u2	WIAF-11805	AF001787	UCP2, uncoupling protein 2 (mitochondrial, proton carrier)	TGCACCACAG [G/A] AGCCATGGCG	M	G	A	G E
G556u3	WIAF-11823	AF001787	UCP2, uncoupling protein 2 (mitochondrial, proton carrier)	TACGGGAATC [A/G] CCGTTTGGAA	S	A	G	S S
G556u4	WIAF-11782	AF001787	UCP2, uncoupling protein 2 (mitochondrial, proton carrier)	ATCCTGACCA [T/C] GGTGGGACT	M	T	C	M T
G561a1	WIAF-12111	HT1176	IDE, insulin-degrading enzyme	ACTGTGGCAT [C/A] CAGATATACT	S	C	A	I I
G561u2	WIAF-12222	HT1176	IDE, insulin-degrading enzyme	ATATTAACCT [C/G] ATGGCTGCAA	M	C	G	F L

G562u1	WIAF-12223	HT27503	680	tumor necrosis factor receptor type 1 associated protein	CCTGTAGTGA [A/C] TCGGCGCTG	M	A	C	N	T
G562u2	WIAF-12224	HT27503	900	tumor necrosis factor receptor type 1 associated protein	CGCTGCAGCG [C/A] CTGCTGGAGG	S	C	A	R	R
G573u1	WIAF-12199	HT28094	469	SSTR1, somatostatin receptor 1	GGACCGCTAC [G/C] TGGCCGTGGT	M	G	C	V	L
G573u2	WIAF-12208	HT28094	480	SSTR1, somatostatin receptor 1	TGGCCGTGGT [G/A] CATCCCATCA	S	G	A	V	V
G573u3	WIAF-12209	HT28094	879	SSTR1, somatostatin receptor 1	TGCAGCTGGT [T/C] AACGTGTTTG	S	T	C	V	V
G574u1	WIAF-11822	HT4058	1054	SSTR5, somatostatin receptor 5	GCCACGGAGC [C/T] GCGTCCAGAC	M	C	T	P	L
G575u1	WIAF-12200	HT28095	99	SSTR3, somatostatin receptor 3	ACGTGTGCGC [G/A] GGCCCAAGCC	S	G	A	A	A
G575u2	WIAF-12217	HT28095	453	SSTR3, somatostatin receptor 3	CCACCGGCTC [G/A] GCCCGCTGGC	S	G	A	S	S
G585u1	WIAF-12204	HT1022	1133	PYGL, phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)	AGCTGAATGA [T/C] ACTCACCCTC	S	T	C	D	D
G585u2	WIAF-12205	HT1022	1988	PYGL, phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)	AGCTGATCAC [T/C] TCAGTGGCAG	S	T	C	T	T
G585u3	WIAF-12225	HT1022	1883	PYGL, phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)	TGTACAACCG [C/T] ATTAAGAAAG	S	C	T	R	R
G585u4	WIAF-12226	HT1022	2037	PYGL, phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)	AAGCAAGTTG [A/G] AAGTCATCTT	M	A	G	K	E
G585u5	WIAF-12231	HT1022	1387	PYGL, phosphorylase, glycogen; liver (Hers disease, glycogen storage disease type VI)	GATGTGGACC [C/G] TCTGAGAAGG	M	C	G	P	R
G586a1	WIAF-12112	HT1878	2410	PFKM, phosphofructokinase, muscle	CCGGGGAAGC [T/G] GCCGTCTAAA	S	T	G	A	A

002060 24425960

G586u2	WIAF-12206	HT1878	375 PFKM, phosphofructokinase, muscle	GGACGACTCC [G/A] AGCTGCTTAC	M	G	A	R	Q
G586u3	WIAF-12207	HT1878	322 PFKM, phosphofructokinase, muscle	TGGAGGCAC [G/A] GTGATTGGAA	S	G	A	T	T
G586u4	WIAF-12227	HT1878	334 PFKM, phosphofructokinase, muscle	TGATTGGAAG [T/C] GCCCGGTGCA	S	T	C	S	S
G586u5	WIAF-12228	HT1878	408 PFKM, phosphofructokinase, muscle	CGTGGATCA [C/G] CAATCTCTGT	M	C	G	T	S
G586u6	WIAF-12235	HT1878	717 PFKM, phosphofructokinase, muscle	CACTGTGGAT [A/G] CCTGGCCCTT	M	A	G	Y	C
G587u1	WIAF-12615	HT3847	366 phosphofructokinase, liver	ATGGCAGCCT [T/C] ACAGGTGCCA	S	T	C	L	L
G589u1	WIAF-12210	L39211	CPT1A, carnitine 1327 palmitoyltransferase I, liver	CAGCGTTCTT [C/T] GTGACGTTAG	S	C	T	F	F
G589u2	WIAF-12215	L39211	CPT1A, carnitine 2080 palmitoyltransferase I, liver	AATATCTCGC [T/C] GTGGAGTCCC	S	T	C	A	A
G589u3	WIAF-12216	L39211	CPT1A, carnitine 679 palmitoyltransferase I, liver	ACTTCAAACG [G/T] ATGACAGCAC	S	G	T	R	R
G589u4	WIAF-12218	L39211	CPT1A, carnitine 1844 palmitoyltransferase I, liver	CCTCACATAC [G/C] AGGCCTCCAT	M	G	C	E	Q
G592u1	WIAF-11814	X96586	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	TCCGGGATCT [C/T] AGTAAGCCAG	S	C	T	L	L
G592u2	WIAF-11815	X96586	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	AAGTATATCA [T/G] TTTCAAATAT	M	T	G	F	V
G592u3	WIAF-11834	X96586	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	GTAGCCATGC [T/C] TACGCAATAC	M	T	C	L	P

002000" 22145900

G592u4	WIAF-11784	X96586	1889	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	CACGAGCACT[A/G]TAAATCCAC	M	A	G	Y	C
G592u5	WIAF-11798	X96586	1677	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	CCATGCTTAC[G/A]CAAACTCTGG	S	G	A	T	T
G592u6	WIAF-11799	X96586	2429	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	TGCCATTACAG[G/C]GATTGTATGT	M	G	C	G	A
G592a7	WIAF-13156	X96586	2205	NSMAF, neutral sphingomyelinase (N-SMase) activation associated factor	ATTCTGCATC[G/A]TGGACTCTA	S	G	A	S	S
G594u1	WIAF-10065	HT3921	1153	annexin V, alt. transcript 2	TTGTGAAATC[T/A]ATTGGAAGTA	S	T	A	S	S
G594u2	WIAF-10098	HT3921	567	annexin V, alt. transcript 2	CGAAGTAATG[C/T]TCAGCGCCAG	M	C	T	A	V
G594u3	WIAF-10099	HT3921	774	annexin V, alt. transcript 2	ATTGCTTCAA[G/C]GACACCTGAA	M	G	C	R	T
G594a4	WIAF-10505	HT3921	424	annexin V, alt. transcript 2	GAGTAGTCGC[C/T]ATGCGACAGG	-	C	T	-	-
G594a5	WIAF-13123	HT3921	571	annexin V, alt. transcript 2	GTAATGCTCA[G/C]CGCCAGGAAA	M	G	C	Q	H
G595u1	WIAF-12203	HT27983	1008	NRIP1, nuclear receptor interacting protein 1	TGCAAGATTA[C/T]AGGCTGTTC	N	C	T	Q	*
G595u2	WIAF-12220	HT27983	785	NRIP1, nuclear receptor interacting protein 1	CCCTCAGTCA[T/C]GATTCCTTAA	S	T	C	H	H
G595u3	WIAF-12232	HT27983	1231	NRIP1, nuclear receptor interacting protein 1	GTTGGCAGTT[A/T]CCAGCTCCCA	M	A	T	Y	F
G595u4	WIAF-12261	HT27983	2048	NRIP1, nuclear receptor interacting protein 1	GCAGTACTCA[G/A]TCTGAAAGC	S	G	A	Q	Q
G595u5	WIAF-12274	HT27983	2376	NRIP1, nuclear receptor interacting protein 1	TCCTGAACCA[C/T]GGCTTTCTGG	M	G	T	G	W
G595u6	WIAF-12275	HT27983	3498	NRIP1, nuclear receptor interacting protein 1	ACTATATTAC[A/G]TGCTTCAAAA	M	A	G	M	V
G595u7	WIAF-12276	HT27983	3671	NRIP1, nuclear receptor interacting protein 1	ACAATAGCCA[T/C]ATGGGAAATA	S	T	C	H	H
G595u8	WIAF-12294	HT27983	2020	NRIP1, nuclear receptor interacting protein 1	ATCAAAATGGA[A/G]TTCCCCACCA	M	A	G	N	S

	WIAP-12295	HT27983	3140	NR1P1, nuclear receptor interacting protein 1	ATTGTGCCC [G/A] CACAGAAGTA	S	G	A	P
G595u9	WIAP-12295	HT27983	3140	NR1P1, nuclear receptor interacting protein 1	ATTGTGCCC [G/A] CACAGAAGTA	S	G	A	P
G596u1	WIAP-10144	HT3537	3299	PC, pyruvate carboxylase	TGCGGTCCAT [C/T] TTGGTCAAGG	S	C	T	I
G596u2	WIAP-10158	HT3537	2662	PC, pyruvate carboxylase	ACCAACTGC [A/C] CTTCAGGCC	M	A	C	H
G596u3	WIAP-10159	HT3537	2156	PC, pyruvate carboxylase	CCATCTCAT [C/A] ACGGGGACG	N	C	A	Y
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598a1	WIAP-12118	HT48666	5585	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GGGACCTATG [C/T] TGATAAACTG	M	C	T	A
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u2	WIAP-12236	HT48666	4456	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	CCTGTTAATA [T/C] TAGGAGTAAG	S	T	C	L
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u3	WIAP-12237	HT48666	6356	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GTTAATGAAG [G/T] CACGTGTGTT	M	G	T	G
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u4	WIAP-12240	HT48666	12219	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GTACCTTTGT [C/T] ATCCAGGCCA	S	C	T	V
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u5	WIAP-12241	HT48666	12480	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	CCAGGCAGAT [C/G] GAGGCCTTAC	M	C	G	I
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u6	WIAP-12244	HT48666	12975	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GAGTAATCAT [T/A] CAGAGATGTGG	S	T	A	I
				HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1					
G598u7	WIAP-12245	HT48666	1424	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	TCCAATAATC [A/T] GTCAACTTTA	M	A	T	Q

G598u8	WIAF-12250	HT48666	5854	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	TTCAAAAGCA [A/T] TTCAATCAAA	M	A	T	I	F
G598u9	WIAF-12251	HT48666	6754	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	TATTCAGCTC [G/A] TCCGTATCCT	M	G	A	V	I
G598u10	WIAF-12252	HT48666	7635	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	ATCTTTACCT [C/T] GGTGCTATGA	S	C	T	L	L
G598u11	WIAF-12254	HT48666	9189	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GTGAAAATCC [A/G] TACTACCTGT	S	A	G	P	P
G598u12	WIAF-12255	HT48666	10119	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	TTCTGGCATT [G/C] CTAGCAGACA	M	G	C	L	F
G598u13	WIAF-12257	HT48666	11109	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	ATCCATCTAT [T/C] GTAATGCGCA	S	T	C	I	I
G598u14	WIAF-12258	HT48666	13513	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	CTATGGACCT [C/T] AGATAACTGT	N	C	T	Q	*
G598u15	WIAF-12259	HT48666	13697	HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	ACCATCACAG [A/G] GATGTGCCAG	M	A	G	E	G

G598u16	WIAF-12265	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	1098		CCCTTTACGA [G/A] GCAGCATTAT	S	G	A	E	E
G598u17	WIAF-12272	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	6079		TATCTGGGAG [A/G] CACCCATTGC	M	A	G	T	A
G598u18	WIAF-12273	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	9551		AAGAGCTCCT [C/T] TGGGAGAATA	M	C	T	S	F
G598u19	WIAF-12277	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	666		GTCTTTGCAA [C/T] GATGTCATTC	S	C	T	N	N
G598u20	WIAF-12278	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	882		GCTCATTGGC [A/G] TATCTTCTTG	S	A	G	R	R
G598u21	WIAF-12279	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	893		TATCTTCTTG [A/T] ATGGATAGAA	M	A	T	E	V
G598u22	WIAF-12280	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	13276		AGAAGTCAGC [A/G] TTCACACGGT	M	A	G	I	V
G598u23	WIAF-12283	HT48666		HERC1, hect (homologous to the E6 AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	6519		CCTGTGTGTT [A/T] GACATGGAAG	M	A	T	L	F

002000-2242560

G598u24	WIAF-12284	HT48666	8386	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GGGTTCTCT [C/T] TTCGCAGAT	M	C	T	L	F
G598u25	WIAF-12286	HT48666	10266	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	CAGCTCAGCA [A/T] CTCGTGCGCA	M	A	T	Q	H
G598u26	WIAF-12287	HT48666	10099	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	CTTGTCTGTA [A/G] CACAGGCCCT	M	A	G	T	A
G598u27	WIAF-12289	HT48666	11835	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	AGAACTGTCT [G/C] CCTGACCCCTG	S	G	C	L	L
G598u28	WIAF-12290	HT48666	12689	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	TTAAACCACA [C/T] TTGGCAGTG	M	C	T	T	I
G598u29	WIAF-12291	HT48666	14655	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	ACGTGGACAA [C/T] GCCGAGGGCT	S	C	T	N	N
G598u30	WIAF-12296	HT48666	393	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	ATCCCCATT [T/C] GCCGGGCGAC	S	T	C	F	F
G598u31	WIAF-12297	HT48666	479	HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	GGCAAGGTGA [A/G] GCAGCAGCAG	M	A	G	K	R

G598u32	WIAF-12298	HT48666		HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	1197	ATGCTCCCAT [T/C] GTCTCCGAAA	S	T	C	I	I
G598u33	WIAF-12300	HT48666		HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	3595	TCCAGAGGAA [C/T] AGGACACTGC	N	C	T	Q	*
G598u34	WIAF-12301	HT48666		HERC1, hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1	3661	CATCTCTCAA [T/C] TGGATAAATG	S	T	C	L	L
G601u1	WIAF-12246	HT27734		PRKMK5, protein kinase, mitogen-activated, kinase 5 (MAP kinase 5)	106	TGGAGAACCA [G/A] GTGCTGGTAA	S	G	A	Q	Q
G601u2	WIAF-12247	HT27734		PRKMK5, protein kinase, mitogen-activated, kinase 5 (MAP kinase 5)	351	GTAATGGAC [A/G] GTTAATAGAG	M	A	G	Q	R
G601u3	WIAF-12292	HT27734		PRKMK5, protein kinase, mitogen-activated, kinase 5 (MAP kinase 5)	617	AGCATATCAT [G/C] TCCCGAGTGG	M	G	C	V	L
G603u1	WIAF-12248	HT4291		mitogen-activated protein (MAP) kinase p38	1336	AGTCATCAGC [T/C] TTGTGCCACC	M	T	C	F	L
G603u2	WIAF-12281	HT4291		mitogen-activated protein (MAP) kinase p38	1230	CTCAGTACCA [C/T] GATCCTGATG	S	C	T	H	H
G610u1	WIAF-12249	HT48690		protein kinase, mitogen-activated, p38Beta (MAP kinase p38Beta)	1012	CCGAGCCATA [T/C] GATGAGAGCG	S	T	C	Y	Y
G610u2	WIAF-12263	HT48690		protein kinase, mitogen-activated, p38Beta (MAP kinase p38Beta)	799	AAATCTCTTC [G/A] GAACACGCC	S	G	A	S	S
G610u3	WIAF-12264	HT48690		protein kinase, mitogen-activated, p38Beta (MAP kinase p38Beta)	848	GCCCCAGAAG [G/A] ACCTGAGCAG	M	G	A	D	N
G610u4	WIAF-12282	HT48690		protein kinase, mitogen-activated, p38Beta (MAP kinase p38Beta)	439	TCCTGGTTTA [C/T] CAGCTGTGTC	S	C	T	Y	Y

G612u1	WIAF-12344	HT1436	1513	RAF1, v-raf-1 murine leukemia viral oncogene homolog 1	TTTGATGCA [A/G] AGAATCAT	M	A	G	K	E
G614u1	WIAF-12267	HT321	603	BRAF, v-raf murine sarcoma viral oncogene homolog B1	GACAGTCTAA [A/G] GAAAGCACTG	M	A	G	K	R
G614u2	WIAF-12268	HT321	2282	BRAF, v-raf murine sarcoma viral oncogene homolog B1	CCAAACAGAG [G/A] ATTATTAGTCT	M	G	A	D	N
G614u3	WIAF-12299	HT321	973	BRAF, v-raf murine sarcoma viral oncogene homolog B1	AGGAAGAGGC [G/A] TCCTTAGCAG	S	G	A	A	A
G616u1	WIAF-12253	HT48746	498	TRAF-interacting protein (I-TRAF)	AAGAAGACAA [G/T] AGGTTTCTTC	N	G	T	E	*
G616u2	WIAF-12269	HT48746	1338	TRAF-interacting protein (I-TRAF)	GCATATACCT [C/G] GAGTATGTGA	M	C	G	R	G
G616u3	WIAF-12285	HT48746	377	TRAF-interacting protein (I-TRAF)	ATAACAATTA [T/C] GGCTGTGTCC	S	T	C	Y	Y
G616u4	WIAF-12288	HT48746	1032	TRAF-interacting protein (I-TRAF)	TGAATTGCG [G/A] GAATTGACCC	M	G	A	G	R
G617u1	WIAF-12256	HT1614	52	PPP1CA, protein phosphatase 1, catalytic subunit, alpha isoform	GAAGCTCAAC [C/T] TGGACTCGAT	S	C	T	L	L
G617u2	WIAF-12270	HT1614	792	PPP1CA, protein phosphatase 1, catalytic subunit, alpha isoform	AAGACGGCTA [C/T] GAGTCTTTTG	S	C	T	Y	Y
G618u1	WIAF-12238	HT27508	1598	protein phosphatase, 2A B56-alpha subunit	CATTGAACCA [A/C] CACAGTTCAA	M	A	C	T	P
G618u2	WIAF-12271	HT27508	1135	protein phosphatase, 2A B56-alpha subunit	ATCAGAAATT [C/T] GTACAAACAGC	S	C	T	F	F
G62u1	WIAF-10369	HT0855	214	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group 6	AGGAGTACCT [G/C] TCCTTTCTGT	S	G	C	L	L
G62u2	WIAF-10370	HT0855	926	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group 6	AAACTGTCT [T/C] TTGAAAGGAA	M	T	C	F	L
G62u3	WIAF-10428	HT0855	2904	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group 6	AGCACGGACA [C/T] GCAGGCCCGG	M	C	T	T	M

G62u4	WIAF-10430	HT0855	3368	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TGACCTCAC [A/G] TGAGTAGTAA	M	A	G	M	V
G62u5	WIAF-10451	HT0855	1376	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TTCTGGGAA [G/A] AAGCTGAAGC	M	G	A	E	K
G62u6	WIAF-10452	HT0855	3716	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TAAGCATTGC [A/G] GAGAGGCCAA	M	A	G	R	G
G62u7	WIAF-10453	HT0855	3967	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	CCCTGAAAGC [A/C] CTGAGGCTCT	S	A	C	A	A
G62u8	WIAF-10454	HT0855	4016	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TGTTGTTCCC [A/G] CCTGGAAGTGG	M	A	G	T	A
G62u9	WIAF-10455	HT0855	3979	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TGAGGCTCTC [T/C] CGTCAGCGGT	S	T	C	S	S
G62u10	WIAF-10456	HT0855	3729	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	GACGCCAAGT [T/G] TGAAGGAAGT	M	T	G	F	C
G62u11	WIAF-10476	HT0855	1275	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TCTGAGATG [G/A] TACTGACTAT	M	G	A	G	D
G62u12	WIAF-10477	HT0855	2017	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	TGATCTTGA [C/T] GAAGGACACA	S	C	T	D	D
G62u13	WIAF-10479	HT0855	3255	ERCC6, excision repair cross-complementing rodent repair deficiency, complementation group	CTACATATC [T/C] GTAAATGATG	S	T	C	S	S

G62u14	WIAF-10481	HT0855		BRCC6, excision repair cross-complementing rodent repair deficiency, complementation group 4317 6	GGCACCTGC [A/G]GGAAGCTTCT	M	A	G	Q	R
G620a1	WIAF-12116	HT1943		PPP2CB, protein phosphatase 2 (formerly 2A), catalytic subunit, 1256 beta isoform	TATCATGGAA [T/A]TAGATGACAC	M	T	A	L	I
G620a2	WIAF-12117	HT1943		PPP2CB, protein phosphatase 2 (formerly 2A), catalytic subunit, 1326 beta isoform	CCTCATGTTA [C/G]ACGGGGCAAC	M	C	G	T	R
G620u3	WIAF-12239	HT1943		PPP2CB, protein phosphatase 2 (formerly 2A), catalytic subunit, 819 beta isoform	TTTATGATG [A/G]ATGTCTGCGA	M	A	G	E	G
G623u1	WIAF-12260	HT3979		PPP1CB, protein phosphatase 1, catalytic subunit, beta isoform 459	TTCAATGGACA [A/G]TATACAGATT	S	A	G	Q	Q
G625u1	WIAF-12266	HT1961		PPP2R2A, protein phosphatase 2 (formerly 2A), regulatory subunit 227 B (PR 52), alpha isoform	CATTCTGGAG [A/G]ATTACTAGCA	M	A	G	E	G
G628a1	WIAF-12104	HT2780		PPP1CC, protein phosphatase 1, catalytic subunit, gamma isoform 1104	AGGGGTATGA [T/A]CACAAAGCAA	M	T	A	I	N
G628a2	WIAF-12105	HT2780		PPP1CC, protein phosphatase 1, catalytic subunit, gamma isoform 973	CCAATTATTG [C/T]GGAGAGTTTG	S	C	T	C	C
G628u3	WIAF-12311	HT2780		PPP1CC, protein phosphatase 1, catalytic subunit, gamma isoform 888	GATCTTATAT [G/T]TAGAGCCCAT	M	G	T	C	F
G630a1	WIAF-12103	HT5086		protein phosphatase 2A, 130 kDa regulatory subunit 704	AAAGATGCAG [A/G]TCTGAACCTCT	M	A	G	D	G
G630a2	WIAF-12106	HT5086		protein phosphatase 2A, 130 kDa regulatory subunit 1015	CGATGGGAAC [G/T]CCCCATCCTT	M	G	T	A	S
G630a3	WIAF-12107	HT5086		protein phosphatase 2A, 130 kDa regulatory subunit 1024	CGCCCCATCC [T/C]TTGGTTTACT	M	T	C	F	L
G630a4	WIAF-12108	HT5086		protein phosphatase 2A, 130 kDa regulatory subunit 837	ACTTAAAGGA [T/C]ATTGCAGGAG	S	T	C	D	D
G630u5	WIAF-12325	HT5086		protein phosphatase 2A, 130 kDa regulatory subunit 1200	TAAAGATGTG [C/T]TTGGACATCT	S	C	T	C	C

G630u6	WIAF-12326	HT5086	2810	protein phosphatase 2A, 130 kDa regulatory subunit	ATGTTACAGG[C/T]TGCAGGGGGA	M	C	T	A	V
G630u7	WIAF-12351	HT5086	512	protein phosphatase 2A, 130 kDa regulatory subunit	ATTATGGCAG[C/T]AACTTACAGA	M	C	T	A	V
G630u8	WIAF-12352	HT5086	703	protein phosphatase 2A, 130 kDa regulatory subunit	CAAGATGCA[G/A]ATCTGAATC	M	G	A	D	N
G630u9	WIAF-12353	HT5086	1069	protein phosphatase 2A, 130 kDa regulatory subunit	ACCTTTGTCT[C/T]ATAGAACTC	M	C	T	H	Y
G634u1	WIAF-11825	X04434	2283	IGF1R, insulin-like growth factor 1 receptor	TGCAAGTGGC[C/T]AACACCACCA	S	C	T	A	A
G634u2	WIAF-11826	X04434	2279	IGF1R, insulin-like growth factor 1 receptor	GTGATGCAAG[T/C]GGCCAAACACC	M	T	C	V	A
G634u3	WIAF-11781	X04434	1731	IGF1R, insulin-like growth factor 1 receptor	ACAAGGACGT[G/A]GAGCCGGGCA	S	G	A	V	V
G634a4	WIAF-13106	X04434	948	IGF1R, insulin-like growth factor 1 receptor	TCCAGCACGG[C/A]GAGTGCATGC	S	C	A	G	G
G634a5	WIAF-13107	X04434	1089	IGF1R, insulin-like growth factor 1 receptor	CTTCTGTCTCA[G/C]ATGCTCCAAG	M	G	C	Q	H
G634a6	WIAF-13108	X04434	2539	IGF1R, insulin-like growth factor 1 receptor	AGAGGAGCA[G/A]ATGACATTC	M	G	A	D	N
G634a7	WIAF-13109	X04434	2606	IGF1R, insulin-like growth factor 1 receptor	AAGTGGCCGG[A/C]ACCTGAGAAT	M	A	C	E	A
G634a8	WIAF-13111	X04434	1543	IGF1R, insulin-like growth factor 1 receptor	CTCCACCACC[A/T]CGTCGAAGAA	M	A	T	T	S
G634a9	WIAF-13112	X04434	1549	IGF1R, insulin-like growth factor 1 receptor	CACCACGTCTG[A/G]AGATCGCAT	M	A	G	K	E
G634a10	WIAF-13113	X04434	1596	IGF1R, insulin-like growth factor 1 receptor	CCCCTGACTA[C/T]AGGGAICTCA	S	C	T	Y	Y
G645u1	WIAF-12332	HT5191	1127	retinoic acid-binding protein II	TCTGCAGACT[C/T]TTCAGGAGAG	M	C	T	L	F
G645u2	WIAF-12333	HT5191	1048	retinoic acid-binding protein II	AAGCATTAGA[G/A]GCCTTACAGA	S	G	A	E	E
G646u1	WIAF-12303	X81479	1204	EMR1, efg-like module containing, mucin-like, hormone receptor-like sequence 1	CAAAATATCCA[T/C]GTGGACTAAA	M	T	C	M	T
G646u2	WIAF-12304	X81479	1919	EMR1, efg-like module containing, mucin-like, hormone receptor-like sequence 1	TTCTGCTGTG[T/G]CGTCCATCC	M	T	G	C	W

G646u3	WIAF-12316	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	590	CTTGCCAGCA [G/T] CATGCAACTT	M	G	T	E	D
G646u4	WIAF-12317	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	799	GCACCAAGCA [G/A] TGGACAGTTG	M	G	A	S	N
G646u5	WIAF-12318	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	558	TGAAGACGTG [A/G] ATGAATGTGC	M	A	G	N	D
G646u6	WIAF-12334	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	207	TTACTATTGC [A/G] CTTGCAACA	M	A	G	T	A
G646u7	WIAF-12335	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	458	TCACCAGCAG [G/C] GTCGCCCTG	M	G	C	R	S
G646u8	WIAF-12336	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	1308	CTCAGCAAAAT [G/A] TCACTCCGGC	M	G	A	V	I
G646u9	WIAF-12337	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	1285	ACACTGGCAT [C/T] TTTTGGAAA	M	C	T	S	F
G646u10	WIAF-12338	X81479	EMR1, egf-like module containing, mucin-like, hormone receptor-like sequence 1	2026	GACAAACAAG [C/T] GGGCTGGCC	M	C	T	T	M
G647u1	WIAF-12339	HT5190	RARA, retinoic acid receptor, alpha	174	TGCCTCCCTA [C/T] GCCTTCTCT	S	C	T	Y	Y
G648a1	WIAF-13332	HT0070	retinoic acid receptor, beta	469	AACGTGAGCC [A/G] GGAGCAGCGT	-	A	G	-	-
G648a2	WIAF-13333	HT0070	retinoic acid receptor, beta	532	ATTGTTTTTA [A/G] GGTGAGAAAT	-	A	G	-	-
G650u1	WIAF-12323	X52773	RXRA, retinoid X receptor, alpha	862	CTCGCGGAAC [G/A] ACCCTGTCAC	M	G	A	D	N
G650u2	WIAF-12341	X52773	RXRA, retinoid X receptor, alpha	102	TCCTGCCGCT [C/T] GATTCTCCA	S	C	T	L	L

002060 2445960

G650u3	WIAF-12348	X52773	673 RXRA, retinoid X receptor, alpha	GCCATGGGC [A/G] TGAAGCGGA	M	A	G	M	V
G650u4	WIAF-12349	X52773	902 RXRA, retinoid X receptor, alpha	GACAAACAGC [T/C] TTTCACCTG	M	T	C	L	P
G653a1	WIAF-13326	HT1458	RARB, retinoic acid receptor, beta	AGGAGAAAGC [T/C] CTCAAAGCAT	S	T	C	A	A
G655a1	WIAF-13327	J05252	PCSK2, proprotein convertase subtilisin/kexin type 2	CCTTCAGCAA [C/T] GGGAGGAAAA	S	C	T	N	N
G655a2	WIAF-13334	J05252	PCSK2, proprotein convertase subtilisin/kexin type 2	CCTATCCTTA [C/A] CCTCGGTACA	N	C	A	Y	*
G655a3	WIAF-13335	J05252	PCSK2, proprotein convertase subtilisin/kexin type 2	TTTCTGCTGC [C/T] GCCAACAAACA	S	C	T	A	A
G658u1	WIAF-11856	J02943	CBG, corticosteroid binding globulin	TCTATGACCT [T/C] GGAGATGTGC	S	T	C	L	L
G658u2	WIAF-13407	J02943	CBG, corticosteroid binding globulin	CCTTCATGAC [T/G] CAGAGCTCCC	M	T	G	S	A
G658u3	WIAF-13408	J02943	CBG, corticosteroid binding globulin	TTCATGACTC [A/G] GAGCTCCCCCT	S	A	G	S	S
G658u4	WIAF-13409	J02943	CBG, corticosteroid binding globulin	TCACCCAGGA [C/T] GCCCAGCTGA	S	C	T	D	D
G663u1	WIAF-13400	HT3157	1202 TPO, thyroid peroxidase	CGCCACGCGC [G/A] CCTGCGGCT	S	G	A	A	A
G663u2	WIAF-13401	HT3157	1282 TPO, thyroid peroxidase	GGCCGCGCCA [G/C] CGAGGTCCCC	M	G	C	S	T
G668a1	WIAF-13350	U53506	DIO2, deiodinase, iodothyronine, type II	TCGATGCCTA [C/A] AACAGGTGA	N	C	A	Y	*
G668a2	WIAF-13351	U53506	DIO2, deiodinase, iodothyronine, type II	TGCCTACAAA [C/A] AGGTGAAAT	M	C	A	Q	K
G668a3	WIAF-13352	U53506	DIO2, deiodinase, iodothyronine, type II	TGTCCTCACT [A/G] CAGAAGGAGG	M	A	G	T	A
G673a1	WIAF-13328	M57464	Human ret proto-oncogene mRNA for tyrosine kinase.	CGAGCCTGGG [G/A] AGCCCCGGGG	M	G	A	E	K
G673a2	WIAF-13336	M57464	Human ret proto-oncogene mRNA for tyrosine kinase.	GGCTCGCCGA [T/A] TTGCCCAGAT	M	T	A	F	I
G673a3	WIAF-13337	M57464	Human ret proto-oncogene mRNA for tyrosine kinase.	ACTGCCAGGC [G/A] TTCAGTGGCA	S	G	A	A	A
G673a4	WIAF-13338	M57464	Human ret proto-oncogene mRNA for tyrosine kinase.	TTGGAAAAAC [T/A] CTAGGAGGAG	S	T	A	T	T
G673a5	WIAF-13339	M57464	Human ret proto-oncogene mRNA for tyrosine kinase.	CGAGTGAGCT [T/G] CGAGACTGTC	S	T	G	L	L

G678a1	WIAP-13353	D49492	1439	GDF10, growth differentiation factor 10	TCGGCTGGAA [T/A] GAATGGATAA	M	T	A	N	K
G68u1	WIAP-10434	HT1115	1214	ERCC3, excision repair cross-complementing rodent repair deficiency, complementation group B 3 (xeroderma pigmentosum group B complementing)	CTGTGGAGCA [G/A] TGGAAGCCCC	S	G	A	Q	Q
G68u2	WIAP-10435	HT1115	1155	ERCC3, excision repair cross-complementing rodent repair deficiency, complementation group B 3 (xeroderma pigmentosum group B complementing)	TGTGACTGCT [G/C] CATGCACTGT	M	G	C	A	P
G68u3	WIAP-10436	HT1115	1327	ERCC3, excision repair cross-complementing rodent repair deficiency, complementation group B 3 (xeroderma pigmentosum group B complementing)	AGCACCTACT [C/T] CATGCTGGGC	M	C	T	S	F
G68u4	WIAP-10461	HT1115	926	ERCC3, excision repair cross-complementing rodent repair deficiency, complementation group B 3 (xeroderma pigmentosum group B complementing)	AGGAAATGAT [T/C] GAGGAATCCC	S	T	C	I	I
G68u5	WIAP-10464	HT1115	1430	ERCC3, excision repair cross-complementing rodent repair deficiency, complementation group B 3 (xeroderma pigmentosum group B complementing)	AAGTGCACAC [C/T] ATACAGCCA	S	C	T	T	T
G684a1	WIAP-13359	X51801	712	BMP7, bone morphogenetic protein 7 (osteogenic protein 1)	GTTTATCAGG [T/G] GCTCAGGAG	M	T	G	V	G
G684a2	WIAP-13360	X51801	719	BMP7, bone morphogenetic protein 7 (osteogenic protein 1)	AGGTGCTCCA [G/A] GAGCACTTGG	S	G	A	Q	Q
G684a3	WIAP-13361	X51801	796	BMP7, bone morphogenetic protein 7 (osteogenic protein 1)	GGCTGGCTGG [T/G] GTTTGACATC	M	T	G	V	G

G684a4	WIAF-13362	X51801		862	BMP7, bone morphogenetic protein 1) BMP7 (osteogenic protein 1)	GGCCTGCAGC [T/G] CTCGGTGGAG	M	T	G	L	R	
G684a5	WIAF-13363	X51801		658	BMP7, bone morphogenetic protein 1) BMP7 (osteogenic protein 1)	ATCTACAAGG [A/G] CTACATCCGG	M	A	G	D	G	
G684u6	WIAF-13834	X51801		1421	BMP7, bone morphogenetic protein 1) BMP7 (osteogenic protein 1)	GCCACTAGCT [C/T] CTCCGAGAAT	-	C	T	-	-	
G685a1	WIAF-13329	D89675		882	BMPL1B, bone morphogenetic protein receptor, type IB	GTTCCCTTTA [T/G] GATTATCTGA	N	T	G	Y	*	
G685a2	WIAF-13330	D89675		920	BMPL1B, bone morphogenetic protein receptor, type IB	GCTAAATCAA [T/C] GCTGAAGTTA	M	T	C	M	T	
G685a3	WIAF-13331	D89675		770	BMPL1B, bone morphogenetic protein receptor, type IB	TATCAGACAG [T/G] GTTGATGAGG	M	T	G	V	G	
G685a4	WIAF-13340	D89675		1303	BMPL1B, bone morphogenetic protein receptor, type IB	TCCTTATCAT [G/A] ACCTAGTGCC	M	G	A	D	N	
G685a5	WIAF-13341	D89675		1372	BMPL1B, bone morphogenetic protein receptor, type IB	GTTACGCCCC [T/G] CATTCCCAAA	M	T	G	S	A	
G685a6	WIAF-13342	D89675		1173	BMPL1B, bone morphogenetic protein receptor, type IB	TGTTGGACGA [G/A] AGCTTGAACA	S	G	A	E	E	
G686u1	WIAF-13816	Z48923		2705	BMPL2, bone morphogenetic protein receptor, type II (serine/threonine kinase)	AAATTTCGCA [G/A] CAACACAAA	M	G	A	S	N	
G686u2	WIAF-13817	Z48923		2749	BMPL2, bone morphogenetic protein receptor, type II (serine/threonine kinase)	TGGAGTTGCC [A/T] AGATGAATAC	N	A	T	K	*	
G687a1	WIAF-13343	HT1455		626	CALB1, calbindin 1, (28kD)	ATGATCAGGA [C/T] GGCATGGAT	S	C	T	D	D	
G696u1	WIAF-11839	HT27700		1075	calcium-sensing receptor	GGGACAAATT [G/C] CAGCTGATCA	M	G	C	A	P	
G696u2	WIAF-11840	HT27700		1551	calcium-sensing receptor	TACTGTGGA [C/T] ACCTTTCTGA	S	C	T	D	D	
G696u3	WIAF-11841	HT27700		1688	calcium-sensing receptor	TTACGGATAT [C/T] CTACAATGTG	M	C	T	S	F	
G696u4	WIAF-11842	HT27700		1698	calcium-sensing receptor	CCTACAATGT [G/T] TACTTAGCAG	S	G	T	V	V	
G696u5	WIAF-11858	HT27700		1767	calcium-sensing receptor	GGAGAGGGCT [C/T] TTCACCAATG	S	C	T	L	L	
G696u6	WIAF-11859	HT27700		1689	calcium-sensing receptor	TACGGATATC [C/T] TACAAATGCT	S	C	T	S	S	
G696u7	WIAF-11860	HT27700		2541	calcium-sensing receptor	TCGTGCTCTG [C/T] ATCTCATGCA	S	C	T	C	C	

G696u8	WIAP-11861	HT27700	2581	calcium-sensing receptor	TGTCCTCTG [G/A] TGTTGAGG	M	G	A	V	M
G696u9	WIAP-11863	HT27700	3159	calcium-sensing receptor	TCTCCGCA [G/C] CGTCCAGCA	M	G	C	K	N
G696u10	WIAP-11872	HT27700	562	calcium-sensing receptor	TCTATTCT [T/A] TTGAGTAGC	M	T	A	F	I
G696u11	WIAP-11878	HT27700	2941	calcium-sensing receptor	CATTCAGC [T/G] ATGCCAGC	M	T	G	Y	D
G696u12	WIAP-13386	HT27700	1145	calcium-sensing receptor	AGGATATCT [G/A] CATCGACTC	M	G	A	C	Y
G696u13	WIAP-13395	HT27700	670	calcium-sensing receptor	GATATTGCC [A/G] TAGAGAGAT	M	A	G	I	V
G696u14	WIAP-13396	HT27700	2243	calcium-sensing receptor	TTCTGTCCA [A/G] TGAGAACAC	M	A	G	N	S
G696u15	WIAP-13397	HT27700	2742	calcium-sensing receptor	AGCTGGAG [T/C] GAGATCATCT	S	T	C	D	D
G698u1	WIAP-13547	X61598	393	CBP1, collagen-binding protein 1	TCAGCAACTC [G/C] ACGCGCGCA	S	G	C	S	S
G698u2	WIAP-13549	X61598	628	CBP1, collagen-binding protein 1	CGGCGCCTG [C/T] TAGTCAAGC	S	C	T	L	L
G698u3	WIAP-13550	X61598	1230	CBP1, collagen-binding protein 1	GCGGCTCCT [G/A] CTATTCATTG	S	G	A	L	L
G701u1	WIAP-12382	HT27657	706	CGRP type I receptor	AACGATGTTG [C/A] AGCAGGAAT	M	C	A	A	E
G701u2	WIAP-12391	HT27657	841	CGRP type I receptor	TGACAAAT [A/T] TACCAGTGT	M	A	T	Y	F
G704u1	WIAP-14046	X60382	1396	COL10A1, collagen, type X, alpha 1 (Schmid metaphyseal chondrodysplasia)	AGGCAATCCA [G/A] GATTCCTGG	M	G	A	G	R
G704u2	WIAP-14070	X60382	1648	COL10A1, collagen, type X, alpha 1 (Schmid metaphyseal chondrodysplasia)	TGCCAACCAG [G/C] GGGTAACAGG	M	G	C	G	R
G704u3	WIAP-14071	X60382	1824	COL10A1, collagen, type X, alpha 1 (Schmid metaphyseal chondrodysplasia)	CATACCAGT [G/C] CATGTGAAG	S	G	C	V	V
G704u4	WIAP-14072	X60382	1582	COL10A1, collagen, type X, alpha 1 (Schmid metaphyseal chondrodysplasia)	AGTCATGCCT [G/C] AGGGTTTAT	M	G	C	E	Q
G705a1	WIAP-13228	J04177	686	COL11A1, collagen, type XI, alpha 1	AGAGAAAAAC [T/A] GTGACAATGA	S	T	A	T	T
G705a2	WIAP-13229	J04177	698	COL11A1, collagen, type XI, alpha 1	TGACAATGAT [T/A] GTTGATTGTA	S	T	A	I	I
G705a3	WIAP-13230	J04177	888	COL11A1, collagen, type XI, alpha 1	TAGTCCAGAC [T/A] GTGACTCTTC	M	T	A	C	S
G705a4	WIAP-13231	J04177	894	COL11A1, collagen, type XI, alpha 1	AGACTGTGAC [T/A] CTTACAGACC	M	T	A	S	T
G705a5	WIAP-13232	J04177	651	COL11A1, collagen, type XI, alpha 1	TGACGGGAAG [T/A] GGATCGGGT	M	T	A	W	R

G705a6	WIAP-13233	J04177	COL11A1, collagen, type XI, alpha 1	661	TGGCATCGGG [T/A] AGCAATCAGC	M	T	A	V	E
G705a7	WIAP-13234	J04177	COL11A1, collagen, type XI, alpha 1	1597	CGTCTGTGGCT [T/C] ACCAGGGGCT	M	T	C	L	S
G705a8	WIAP-13235	J04177	COL11A1, collagen, type XI, alpha 1	2745	TGGGTTTCCA [G/A] GTCCCAATGG	M	G	A	G	S
G705a9	WIAP-13236	J04177	COL11A1, collagen, type XI, alpha 1	4385	GTCCAGAAGG [T/A] CTTCCGGGCA	S	T	A	G	G
G705a10	WIAP-13237	J04177	COL11A1, collagen, type XI, alpha 1	4576	GAAAAGGTG [A/T] COGAGGGCTC	M	A	T	D	V
G705a11	WIAP-13238	J04177	COL11A1, collagen, type XI, alpha 1	4306	GCTAAGGGGG [A/C] AGCAGGTGCA	M	A	C	E	A
G705a12	WIAP-13239	J04177	COL11A1, collagen, type XI, alpha 1	4837	AGACATACTG [A/G] AGGCATGCAA	M	A	G	E	G
G705a13	WIAP-13240	J04177	COL11A1, collagen, type XI, alpha 1	4931	AACAAGACAT [C/T] GAGCATATGA	S	C	T	I	I
G705a14	WIAP-13346	J04177	COL11A1, collagen, type XI, alpha 1	299	AAGCACTAGA [T/G] TTTCACAAAT	M	T	G	D	E
G705a15	WIAP-13347	J04177	COL11A1, collagen, type XI, alpha 1	2225	GGGAGCCTGG [G/C] CCTCCAGGTC	S	G	C	G	G
G705u16	WIAP-13679	J04177	COL11A1, collagen, type XI, alpha 1	5493	AATTGATCAA [G/A] TACCTATTGT	M	G	A	V	I
G705u17	WIAP-13700	J04177	COL11A1, collagen, type XI, alpha 1	3484	GGAGTTCAAG [G/A] TCCTGTTGGT	M	G	A	G	D
G705u18	WIAP-13709	J04177	COL11A1, collagen, type XI, alpha 1	5392	GAGATGTCCT [A/T] TGACAATAAT	M	A	T	Y	F
G707u1	WIAP-12363	U32169	COL11A2, collagen, type XI, alpha 1	4996	TCCCTTGAGA [C/T] TCCGTGGGGC	M	C	T	L	F
G707u2	WIAP-12374	U32169	COL11A2, collagen, type XI, alpha 1	3580	CAATGGCGCT [G/A] ATGGCCCA	M	G	A	D	N
G707u3	WIAP-12385	U32169	COL11A2, collagen, type XI, alpha 1	2059	GCCTGGCTCA [G/A] ACGGACCCCC	M	G	A	D	N
G708a1	WIAP-13354	U73778	COL12A1, collagen, type XII, alpha 1	1885	GCCTCTCCTC [C/T] TGCAGAGACC	M	C	T	P	L
G708a2	WIAP-13355	U73778	COL12A1, collagen, type XII, alpha 1	3630	TGTTGGACAA [G/A] AAATGACAAC	M	G	A	E	K
G708a3	WIAP-13356	U73778	COL12A1, collagen, type XII, alpha 1	3905	GCTTGTTCGA [A/T] GCTGTGGCAA	M	A	T	Q	H
G708a4	WIAP-13357	U73778	COL12A1, collagen, type XII, alpha 1	7051	ATTCACCAG [C/A] CCGGATGTA	M	C	A	A	D
G708a5	WIAP-13358	U73778	COL12A1, collagen, type XII, alpha 1	8036	AAGAAGTAAA [G/A] ACATTATTTT	S	G	A	K	K

G708a6	WIAF-13364	U73778	COL12A1, collagen, type XII, alpha 1	1461	TGGCTCCTAT [A/T] GCATTGGGAT	M	A	T	S	C
G708a7	WIAF-13365	U73778	COL12A1, collagen, type XII, alpha 1	2344	ATTACTTGGA [C/T] TCAAGCTCCA	M	C	T	T	I
G708a8	WIAF-13366	U73778	COL12A1, collagen, type XII, alpha 1	5207	CAGATAAGAT [G/A] GAGACCATCT	M	G	A	M	I
G708a9	WIAF-13367	U73778	COL12A1, collagen, type XII, alpha 1	6592	GAGCCCATGG [A/T] AGCCTTTGTT	M	A	T	E	V
G708a10	WIAF-13368	U73778	COL12A1, collagen, type XII, alpha 1	7434	CCAGGATGAG [G/A] TCAAGAAGGC	M	G	A	V	I
G708a11	WIAF-13369	U73778	COL12A1, collagen, type XII, alpha 1	9108	ACCTCGGGG [C/G] TGCCTGGGCC	M	C	G	L	V
G708a12	WIAF-13370	U73778	COL12A1, collagen, type XII, alpha 1	9111	TCGGGGGCTG [C/T] CTGGGCCCCC	M	C	T	P	S
G708a13	WIAF-13371	U73778	COL12A1, collagen, type XII, alpha 1	9196	CCCCCTGGCC [G/A] TCCTGGAAC	M	G	A	R	H
G708a14	WIAF-13972	U73778	COL12A1, collagen, type XII, alpha 1	3044	CAGTATTGTC [C/A] ACTTACAGCA	S	C	A	A	A
G708a15	WIAF-13977	U73778	COL12A1, collagen, type XII, alpha 1	5853	TGTACTGTGA [G/C] TTCCCGTTTA	M	G	C	V	L
G710u1	WIAF-12371	D38163	COL19A1, collagen, type XIX, alpha 1	3082	AGGAACAAG [G/T] GCTCCATGGG	M	G	T	G	C
G710u2	WIAF-12388	D38163	COL19A1, collagen, type XIX, alpha 1	2089	TCCAGGGACT [C/T] CAGGGAATGA	M	C	T	P	S
G711u1	WIAF-12360	L25286	COL15A1, collagen, type XV, alpha 1	1449	TGTGGGTCCA [A/G] GCAGTGAAGA	M	A	G	S	G
G711u2	WIAF-12372	L25286	COL15A1, collagen, type XV, alpha 1	4001	ATATTCCAAT [A/G] TACTCCTTTG	M	A	G	I	M
G711u3	WIAF-12373	L25286	COL15A1, collagen, type XV, alpha 1	3867	CCATTGCAA [G/T] ATCTGTCCAC	M	G	T	D	Y
G711a4	WIAF-13372	L25286	COL15A1, collagen, type XV, alpha 1	395	CCAGCAGCAC [C/T] CGTGGTGGCG	S	C	T	T	T
G711a5	WIAF-13373	L25286	COL15A1, collagen, type XV, alpha 1	3101	AAGGCGACCA [G/A] GGAGCCACAG	S	G	A	Q	Q
G712u1	WIAF-13619	M92642	COL16A1, collagen, type XVI, alpha 1	3608	GGCACCAGG [G/A] ATTTCAAGGC	M	G	A	G	E
G712u2	WIAF-13620	M92642	COL16A1, collagen, type XVI, alpha 1	4944	CCATGAAGAC [C/T] ATGAAGGGGC	S	C	T	T	T
G712u3	WIAF-13621	M92642	COL16A1, collagen, type XVI, alpha 1	4707	CCAAAGGTGA [A/C] AAAGGGGACA	M	A	C	E	D
G712u4	WIAF-13654	M92642	COL16A1, collagen, type XVI, alpha 1	421	GCCACGCGA [C/A] GAGTATCCC	S	C	A	R	R

20250624 22:59:00

G712u5	WIAF-13655	M92642	COL16A1, collagen, type XVI, alpha 1	444	GGGGTCTCCC [G/A] GAGAGTTTG	S	G	A	P	P
G712u6	WIAF-13656	M92642	COL16A1, collagen, type XVI, alpha 1	338	CTCATGAAGA [A/C] GTCGTCCATC	M	A	C	K	T
G712u7	WIAF-13862	M92642	COL16A1, collagen, type XVI, alpha 1	3227	CCTGGTCCTC [C/T] GGGATTGCCA	M	C	T	P	L
G712u8	WIAF-13863	M92642	COL16A1, collagen, type XVI, alpha 1	3199	TCCTGGCTGT [G/T] TTGGGAGCCC	M	G	T	V	F
G712u9	WIAF-13878	M92642	COL16A1, collagen, type XVI, alpha 1	318	ACCTCATCCA [C/T] CGACTCAGCC	S	C	T	H	H
G712u10	WIAF-13882	M92642	COL16A1, collagen, type XVI, alpha 1	1346	ACAGGCGAGA [A/G] GGGCCAGAAA	M	A	G	K	R
G712u11	WIAF-13883	M92642	COL16A1, collagen, type XVI, alpha 1	1309	GTCAGGAGCT [C/T] TGGACCCCTC	S	C	T	L	L
G715a1	WIAF-13344	Z74615	COL1A1, collagen, type I, alpha 1	3504	TCCTGGTGAA [C/G] AAGTGCCCTC	M	C	G	Q	E
G717u1	WIAF-12639	Z74616	COL1A2, collagen, type I, alpha 2	3988	ATGAGGAGAC [T/C] GGCAACCTGA	S	T	C	T	T
G720u1	WIAF-12367	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	3494	GGTGCAATCG [G/A] CAGTCCAGGA	M	G	A	G	D
G720u2	WIAF-12383	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	3035	GGTCTCAAGG [G/A] TGAAGTGGG	M	G	A	G	D
G720a3	WIAF-13374	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	214	TCCTGGTCAG [T/C] CCTATGCGGA	M	T	C	S	P
G720a4	WIAF-13375	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	1953	CTGGACCTCA [A/G] GGACCCCCAG	S	A	G	Q	Q
G720a5	WIAF-13376	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	2194	TAGAGGTGGA [G/A] CTGGTCCCC	M	G	A	A	T
G720a6	WIAF-13377	X14420	COL3A1, collagen, type III, alpha 1 (Ehlers-Danlos syndrome type IV, autosomal dominant)	3731	GGGATTGGAG [G/A] TGAAGAAGCT	M	G	A	G	D

G722u1	WIAF-14132	HT3162	140	COL4A2, collagen, type IV, alpha 2	GAGATTGGCG [C/T] GACTGGTGAT	M	C	T	A	V
G724a1	WIAF-12120	X81053	3892	COL4A4, collagen, type IV, alpha 4	CTCGTGGAAA [G/A] AAAGTCCCC	S	G	A	K	K
G724a2	WIAF-12121	X81053	4187	COL4A4, collagen, type IV, alpha 4	GAAAGGACCA [A/G] TGGATTCCC	M	A	G	M	V
G724a3	WIAF-12122	X81053	3802	COL4A4, collagen, type IV, alpha 4	ATGATGTGGG [G/A] CCACCTGGTC	S	G	A	G	G
G724a4	WIAF-12123	X81053	1838	COL4A4, collagen, type IV, alpha 4	ACCAGGAAAG [C/A] ATGTGCTCTC	M	C	A	H	N
G724u5	WIAF-12364	X81053	376	COL4A4, collagen, type IV, alpha 4	CTGTTTGCCA [C/T] TGTGTTCTCTG	S	C	T	H	H
G724u6	WIAF-12365	X81053	2018	COL4A4, collagen, type IV, alpha 4	TCCAGGGGAT [C/G] ATGAAGATGC	M	C	G	H	D
G724u7	WIAF-12366	X81053	4756	COL4A4, collagen, type IV, alpha 4	GCCTTCCGGT [A/G] TTTAGCACGC	S	A	G	V	V
G724u8	WIAF-12377	X81053	3595	COL4A4, collagen, type IV, alpha 4	CTGACCACCC [A/G] GGGTGCCACG	S	A	G	P	P
G724u9	WIAF-12378	X81053	3516	COL4A4, collagen, type IV, alpha 4	GGAGCATCCG [G/C] AGAGCAGGCG	M	G	C	G	A
G724u10	WIAF-12379	X81053	4288	COL4A4, collagen, type IV, alpha 4	CTGCTCTTCC [A/G] GGTCCACAG	S	A	G	P	P
G724u11	WIAF-12380	X81053	5140	COL4A4, collagen, type IV, alpha 4	GCCACTTTT [C/A] GCAATAAGT	M	C	A	F	L
G724u12	WIAF-12387	X81053	207	COL4A4, collagen, type IV, alpha 4	GACTTGCTTG [C/T] GATGTGTTCT	-	C	T	-	-
G727u1	WIAF-12362	D90279	5135	COL5A1, collagen, type V, alpha 1	TTCAGGTTT [A/T] CTGCACTTC	M	A	T	Y	F
G727u2	WIAF-12369	D90279	4686	COL5A1, collagen, type V, alpha 1	AACAGGGTAT [C/T] ACTGTCCTT	S	C	T	I	I
G727u3	WIAF-12370	D90279	4608	COL5A1, collagen, type V, alpha 1	TCGGTCTCTCC [G/C] GGTGACACGG	S	G	C	P	P
G727a4	WIAF-13300	D90279	2034	COL5A1, collagen, type V, alpha 1	ACGGCTGGC [T/A] GGGTTGCCAG	S	T	A	A	A
G727a5	WIAF-13301	D90279	2073	COL5A1, collagen, type V, alpha 1	GTGACCTTGG [T/C] CCTTCGGCC	S	T	C	G	G
G727a6	WIAF-13302	D90279	3763	COL5A1, collagen, type V, alpha 1	CGGGCAGAAA [G/A] GTGATGAAGG	M	G	A	G	S

G729u1	WIAF-11844	L02870	2345	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	ATGGACTGGA [G/A] CCAGATACTG	S	G	A	E	E
G729u2	WIAF-11845	L02870	3083	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	TATCCTGGCG [G/A] CCACTCAGAG	S	G	A	R	R
G729u3	WIAF-11846	L02870	3031	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GACTCGGTGA [C/T] TTTGGCCTGG	M	C	T	T	I
G729u4	WIAF-11851	L02870	1289	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	CGGACTATGA [G/T] GTGACCCGTGA	M	G	T	E	D
G729u5	WIAF-11852	L02870	1032	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	CCAAGTGA [G/T] TGATTGCCCT	M	G	T	V	L
G729u6	WIAF-11853	L02870	1897	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	CGCCGGGAGC [C/T] GGAACCTCCA	M	C	T	P	L
G729u7	WIAF-11854	L02870	1827	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GCTTAGCTAC [A/T] CTGTGCGGGT	M	A	T	T	S
G729u8	WIAF-11855	L02870	1893	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	TGTCCGCCGG [G/A] AGCCGGAAC	M	G	A	E	K

G729u9	WIAF-11864	L02870	2142	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GGGCCCTGCT [G/A] CAGTCATCGT	M	G	A	A	T
G729u10	WIAF-11865	L02870	2353	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GAGCCAGATA [C/T] TGAGTATAAG	M	C	T	T	I
G729u11	WIAF-11866	L02870	2221	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	TCATCTGTCA [C/T] CATTACCTGG	M	C	T	T	I
G729u12	WIAF-11869	L02870	6585	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	ACCAGGAGAG [C/T] GTGGTATGGC	M	C	T	R	C
G729u13	WIAF-11870	L02870	8169	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GGGTGACCGA [G/T] GCTTTGACGG	M	G	T	G	C
G729u14	WIAF-11877	L02870	438	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	CGCATCCGT [G/A] AGCTTAGCTA	M	G	A	E	K
G729u15	WIAF-11882	L02870	3481	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	AGGATCCGTG [A/T] CATGCCCTAC	M	A	T	D	V
G729u16	WIAF-11883	L02870	5654	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	ACGAGAGAACC [T/C] GGGGACCCTG	S	T	C	P	P

G729u17	WIAF-11884	L02870	7124	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	TGCCAGGGCC [G/C] CGAGGCGAGA	S	G	C	P	P
G729u18	WIAF-11885	L02870	7757	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	GCTTGGATGG [T/C] GACAAAGGAC	S	T	C	G	G
G729u19	WIAF-13389	L02870	1615	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	ACCGTGGTTC [C/T] CACTGGACCA	M	C	T	P	L
G729u20	WIAF-13390	L02870	2930	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	TCTTAGGGCC [G/A] GCTGGAGAAG	S	G	A	P	P
G729u21	WIAF-13399	L02870	5145	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	CCAGGGAGAT [C/T] CTGGAGAGGA	M	C	T	P	S
G729u22	WIAF-13411	L02870	3472	COL7A1, collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)	ATCTTGCAAA [G/A] GATCCGTGAC	M	G	A	R	K
G730a1	WIAF-13303	X57527	305	COL8A1, collagen, type VIII, alpha 1	ATGGGCAAGG [A/G] AGCGGTTCCC	M	A	G	E	G
G732u1	WIAF-12616	M95610	936	COL9A2, collagen, type IX, alpha 2	CAGGCGGAC [A/G] GCCCGGAAGT	S	A	G	T	T
G732u2	WIAF-12617	M95610	696	COL9A2, collagen, type IX, alpha 2	AAGGAGAGA [C/T] GGGCCCTCAT	S	C	T	D	D
G732u3	WIAF-12619	M95610	1288	COL9A2, collagen, type IX, alpha 2	AAGTGGGTGA [C/T] CCAGGGGTGG	M	C	T	P	S
G732u4	WIAF-12620	M95610	962	COL9A2, collagen, type IX, alpha 2	CCACCAGGGC [C/G] TAGCGGGTGT	M	C	G	P	R

G752u1	WIAF-11843	HT1782	1405	CHGA, chromogranin A (parathyroid secretory protein 1)	CGGCATTGA [A/G] GCAGAGCTGG	S	A	G	E	E
G752u2	WIAF-11873	HT1782	1187	CHGA, chromogranin A (parathyroid secretory protein 1)	GGACAACGGG [G/A] ACAGTTCCAT	M	G	A	D	N
G754a1	WIAF-13382	K02043	663	NPPA, natriuretic peptide precursor A	GTACAATGCC [G/A] TGTCACGCG	M	G	A	V	M
G756u1	WIAF-12395	HT3508	2086	SCNN1A, sodium channel, nonvoltage-gated 1 alpha	CAGTTCCTCC [A/G] CCTGTCCTCT	M	A	G	T	A
G757u1	WIAF-12420	HT28563	797	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	CCTGCAGGCC [A/C] CCAACATCTT	M	A	C	T	P
G757u2	WIAF-12421	HT28563	1006	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	GAACCTGAATT [C/T] GGCCTGAAGT	S	C	T	F	F
G757u3	WIAF-12430	HT28563	1768	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	TCATCGACTT [T/C] GTGTGGATCA	S	T	C	F	F
G757u4	WIAF-12494	HT28563	662	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	AAGCAGCTCA [G/C] CATCAGAAAA	M	G	C	A	P
G757u5	WIAF-12506	HT28563	1091	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	GATGCTTCAC [G/C] AGCAGAGGTC	M	G	C	E	Q
G757u6	WIAF-12507	HT28563	1452	SCNN1B, sodium channel, nonvoltage-gated 1, beta (Liddle syndrome)	ACCTGCATTG [G/T] CATGTCCAAG	M	G	T	G	V
G758u1	WIAF-12621	HT27856	415	SCNN1D, sodium channel, nonvoltage-gated 1, delta	CGGGAACCCA [C/T] GTCGGCCGAG	M	C	T	R	C
G758u2	WIAF-12632	HT27856	325	SCNN1D, sodium channel, nonvoltage-gated 1, delta	CCTCTTTGAG [C/T] GTCACCTGGCA	M	C	T	R	C
G758u3	WIAF-12634	HT27856	879	SCNN1D, sodium channel, nonvoltage-gated 1, delta	ATGCGTCTG [G/A] ACAGTCTCAGC	N	G	A	W	*
G758u4	WIAF-12635	HT27856	1138	SCNN1D, sodium channel, nonvoltage-gated 1, delta	CGTGGAGGTG [G/C] AGCTGCTACA	M	G	C	E	Q
G762u1	WIAF-12622	HT27531	1850	NPR3, natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor C)	TAGGAGCTGG [C/T] TTGCTAATGG	S	C	T	G	G

G762u2	WIAF-12623	HT27531	1926 C)	NPR3, natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor	AGAAGAAAGT [A/G] ACCTTGAAA	M	A	G	N	D
G762u3	WIAF-12624	HT27531	1791 C)	NPR3, natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor	CAAATCATCA [G/T] GTGGCTTAGA	M	G	T	G	C
G762u4	WIAF-12636	HT27531	1963 C)	NPR3, natriuretic peptide receptor C/guanylate cyclase C (atrionatriuretic peptide receptor	GAAGATTCCA [T/C] CAGATCCCAT	M	T	C	I	T
G763u1	WIAF-12659	HT3183	1633 B)	NPR2, natriuretic peptide receptor B/guanylate cyclase B (atrionatriuretic peptide receptor	CTGGGCCCTT [C/T] CCTGATGAAC	M	C	T	S	F
G763u2	WIAF-12678	HT3183	668 B)	NPR2, natriuretic peptide receptor B/guanylate cyclase B (atrionatriuretic peptide receptor	TGCCATCACT [T/C] CTGCTGTGG	S	T	C	L	L
G763u3	WIAF-12684	HT3183	2354 B)	NPR2, natriuretic peptide receptor B/guanylate cyclase B (atrionatriuretic peptide receptor	TGTTTGAAGT [C/T] AAACATATGA	S	C	T	L	L
G764u1	WIAF-12698	HT1221	3021 A)	NPR1, natriuretic peptide receptor A/guanylate cyclase A (atrionatriuretic peptide receptor	CCCCGTTACT [G/T] TCTCTTTGGG	M	G	T	C	F
G764u2	WIAF-12708	HT1221	588 A)	NPR1, natriuretic peptide receptor A/guanylate cyclase A (atrionatriuretic peptide receptor	GAGCGCCAAG [C/T] GCTCATGCTC	M	C	T	A	V

G764u3	WIAF-12709	HT1221	1897 A	NPRL, natriuretic peptide receptor A/guanylate cyclase A (atrionatriuretic peptide receptor	GTCCCCGTGG [G/A] AGCCTGCAGG	S	G	A	G	G	
G765u1	WIAF-10012	HT2456	604	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	GCTGGCACAA [A/G] GCTGCGGGCA	S	A	G	N	N	
G765u2	WIAF-10014	HT2456	2350	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	TGATGGCCAC [A/G] TCCCGGAAAT	S	A	G	T	T	
G765u3	WIAF-10025	HT2456	1688	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	CCCACCTGCAC [C/A] AGTGTGACAT	M	C	A	Q	K	
G765u4	WIAF-10027	HT2456	3220	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	TCCCCTTCAG [C/T] TACCTGTCG	S	C	T	S	S	
G765u5	WIAF-10028	HT2456	3409	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	TCAGGTACTT [T/C] GTCAGCTTCA	S	T	C	F	F	
G765u6	WIAF-10040	HT2456	775	DCP1, dipeptidyl carboxypeptidase 1 (angiotensin I converting enzyme)	AGCCCCCTCTA [C/T] CTGAACCTCC	S	C	T	Y	Y	
G772u1	WIAF-12626	HT2121	1064	AVPR2, arginine vasopressin receptor 2 (nephrogenic diabetes insipidus)	TCAGCAGCAG [C/T] GTGTCCTCAG	S	C	T	S	S	
G772u2	WIAF-12627	HT2121	998	AVPR2, arginine vasopressin receptor 2 (nephrogenic diabetes insipidus)	CCTTTGTGCT [A/G] CTCATGTTGC	S	A	G	L	L	
G773u1	WIAF-12644	HT2141	163	SLC6A6, solute carrier family 6 (neurotransmitter transporter, taurine), member 6	CTAGCAAGAT [C/T] GACTTTGTGC	S	C	T	I	I	

G773u2	WIAP-12645	HT2141	SLC6A6, solute carrier family 6 (neurotransmitter transporter, taurine), member 6	445	TCGTCATCCT [G/C] GCCTGGGCCA	S G C L L
G773u3	WIAP-12665	HT2141	SLC6A6, solute carrier family 6 (neurotransmitter transporter, taurine), member 6	289	TGTTTGGGAG [C/T] GGCTGCTG	S C T S S
G773u4	WIAP-12666	HT2141	SLC6A6, solute carrier family 6 (neurotransmitter transporter, taurine), member 6	382	CCTGTGTTCTC [T/C] GGTATGGCT	S T C S S
G776u1	WIAP-11857	U66088	SLC5A5, solute carrier family 5 (sodium iodide symporter), member 1457	5	TAGAGACCT [C/T] ATCAACCTC	S C T L L
G776u2	WIAP-11871	U66088	SLC5A5, solute carrier family 5 (sodium iodide symporter), member 2039	5	GATTGTTGTG [G/C] TGGACCTCG	M G C W C
G776u3	WIAP-13398	U66088	SLC5A5, solute carrier family 5 (sodium iodide symporter), member 1379	5	GGCTTTTCCT [G/A] GCCTGTGCTT	S G A L L
G777u1	WIAP-12646	HT27843	SMRT	4348	ATACAATATC [A/G] GCCAGCCTGG	M A G S G
G777u2	WIAP-12654	HT27843	SMRT	2031	CTGAGCTGGG [T/C] AAGCCGCGC	S T C G G
G777u3	WIAP-12655	HT27843	SMRT	2052	AGAGCCCTC [G/A] ACCTATGAGG	S G A L L
G777u4	WIAP-12675	HT27843	SMRT	2205	CTCGTGAGAT [C/T] GCCAAGTCCC	S C T I I
G778u1	WIAP-14093	HT1449	TG, thyroglobulin	8212	ATCTGCTCTC [T/C] GAGACATCT	M T C L P
G778u2	WIAP-14111	HT1449	TG, thyroglobulin	6033	ATGTGAACGA [C/T] GGTGCGATGC	M C T R W
G778u3	WIAP-14112	HT1449	TG, thyroglobulin	6894	GTATCTCAAT [G/T] GTTCATCCC	M G T V L
G778u4	WIAP-14125	HT1449	TG, thyroglobulin	2375	ATGGCCCTCC [T/C] GAGCAGGTCT	S T C P P
G778u5	WIAP-14136	HT1449	TG, thyroglobulin	1931	AGGATGTCCA [A/G] TGCCTTTCCG	S A G Q Q
G783u1	WIAP-12649	X97674	H.sapiens mRNA for transcriptional intermediary factor 2.	4008	CTAGTGGTAT [G/C] CCAGCACTA	M G C M I
G783u2	WIAP-12658	X97674	H.sapiens mRNA for transcriptional intermediary factor 2.	2566	GCCTGGCAGT [G/A] AGCTGGACAA	M G A E K
G783u3	WIAP-12671	X97674	H.sapiens mRNA for transcriptional intermediary factor 2.	3828	CTCTGAGGCC [T/C] GGAGTACCAA	S T C P P
G785u1	WIAP-13385	HT1291	TTR, transthyretin (prealbumin, amyloidosis type I)	386	CCAACGACTC [C/T] GGCCCCCGCC	S C T S S

000000-2445900

G787u1	WIAP-12652	HT27477	TRIP15: thyroid receptor 468 interacting protein 15	GAAATTTATA [T/C] TTAGAACGAG	S	T	C	Y	Y
G792u1	WIAP-12661	HT27476	thyroid receptor interactor 14	CAGCTGGAAC [G/A] TGAAGAGGCG	M	G	A	V	M
G793u1	WIAP-12643	HT5152	thyroid receptor interactor 8	GGAAGCTTTT [C/G] AAAGAATGTT	N	C	G	S	*
G794u1	WIAP-12664	HT5136	PSMC5, proteasome (prosome, 1110 macropain) 26S subunit, ATPase, 5	GCCTGTGCAC [G/A] GAAGCTGGCA	S	G	A	T	T
G797u1	WIAP-11847	HT3919	140 glutamate receptor 3, flip isoform	CTCAGGAGG [A/G] TTCCCAACA	S	A	G	G	G
G797u2	WIAP-11848	HT3919	759 glutamate receptor 3, flip isoform	GGTTGTGATC [C/T] TAGGGAACA	S	C	T	L	L
G797u3	WIAP-11849	HT3919	1253 glutamate receptor 3, flip isoform	GCTACTGGAA [C/T] GAGTATGAAA	S	C	T	N	N
G797u4	WIAP-11850	HT3919	1770 glutamate receptor 3, flip isoform	TCCTTTCCCTA [G/A] TCAGCAGGTT	M	G	A	V	I
G797u5	WIAP-13404	HT3919	2711 glutamate receptor 3, flip isoform	GCTACAACGT [G/A] TATGGAACAG	S	G	A	V	V
G797u6	WIAP-13405	HT3919	2376 glutamate receptor 3, flip isoform	CTCAGCATTA [G/A] GAACGCCCTGT	M	G	A	G	R
G798u1	WIAP-11868	X77748	GRM3, glutamate receptor, 2655 metabotropic 3	TGCAGACGAC [A/G] ACCATGTGCA	S	A	G	T	T
G798u2	WIAP-11879	X77748	GRM3, glutamate receptor, 2771 metabotropic 3	CACAGACTGC [A/G] CCTCAACAGG	M	A	G	H	R
G798a3	WIAP-12085	X77748	GRM3, glutamate receptor, 2699 metabotropic 3	GTGGTCTTGG [G/C] CTGTTTGTIT	M	G	C	G	A
G798a4	WIAP-12086	X77748	GRM3, glutamate receptor, 2738 metabotropic 3	ATCCTGTTC [A/G] ACCCCAGAAG	M	A	G	Q	R
G798a5	WIAP-12087	X77748	GRM3, glutamate receptor, 2072 metabotropic 3	ACACCCCTGG [T/C] CAAAGCATCG	M	T	C	V	A
G798a6	WIAP-12088	X77748	GRM3, glutamate receptor, 2235 metabotropic 3	CCCTGTCTAC [C/T] AAGACAAACT	S	C	T	T	T
G798u7	WIAP-13391	X77748	GRM3, glutamate receptor, 1131 metabotropic 3	GGCCCAATGC [C/T] TCCTTCACCT	S	C	T	A	A
G799u1	WIAP-11880	M81883	GAD1, glutamate decarboxylase 1 (brain, 67kD)	CAACAATGC [C/T] TGGAACTGGC	S	C	T	L	L
G799u2	WIAP-11881	M81883	GAD1, glutamate decarboxylase 1 (brain, 67kD)	AGGTATACT [C/T] CAAGGATGCA	S	C	T	L	L

000000-2445960

G799u3	WIAF-13392	M81883		GAD1, glutamate decarboxylase 1 661 (brain, 67kD)	GGTGGCCCA [T/C] GGATGCACCA	S	T	C	H	H
G799u4	WIAF-13393	M81883		GAD1, glutamate decarboxylase 1 556 (brain, 67kD)	AGCTGATGGC [G/A] TCCTCGACCC	S	G	A	A	A
G799u5	WIAF-13410	M81883		GAD1, glutamate decarboxylase 1 1229 (brain, 67kD)	CCTCATGGAA [C/T] AAATAACACT	N	C	T	Q	*
G801u1	WIAF-13403	D49394		HTR3, 5-hydroxytryptamine 1596 (serotonin) receptor 3	TTTACCTGCT [A/G] GCGGTGCTGG	S	A	G	L	L
G803a1	WIAF-13118	U66406		EFNB3, ephrin-B3 1446	CTGGGCTGG [G/A] GGGTGGAGGT	M	G	A	G	E
G804u1	WIAF-11887	Z26653		LAMA2, laminin, alpha 2 (merosin, 7237 congenital muscular dystrophy)	TCACTGATGG [G/T] CACATAAAAG	S	G	T	G	G
G804u2	WIAF-11901	Z26653		LAMA2, laminin, alpha 2 (merosin, 9351 congenital muscular dystrophy)	GCAAGCCACT [G/C] GAGGTTAATT	M	G	C	W	S
G804u3	WIAF-11924	Z26653		LAMA2, laminin, alpha 2 (merosin, 8740 congenital muscular dystrophy)	ACACTACCCG [A/G] AGAATTGCTC	S	A	G	R	R
G804u4	WIAF-11943	Z26653		LAMA2, laminin, alpha 2 (merosin, 8577 congenital muscular dystrophy)	ACCAAAATCA [A/G] TGATGGCCAG	M	A	G	N	S
G804a5	WIAF-12089	Z26653		LAMA2, laminin, alpha 2 (merosin, 3372 congenital muscular dystrophy)	CTCTGTGACT [G/A] CTTCTCCCT	M	G	A	C	Y
G804a6	WIAF-13227	Z26653		LAMA2, laminin, alpha 2 (merosin, 7047 congenital muscular dystrophy)	GTCAGTCCTC [A/G] GGTGGAGAGAT	M	A	G	Q	R
G804u7	WIAF-13437	Z26653		LAMA2, laminin, alpha 2 (merosin, 6791 congenital muscular dystrophy)	TGTCAGAGCC [C/T] TGGATGGACC	S	C	T	L	L
G805u1	WIAF-13416	U14755		LHX1, LIM homeobox protein 1 799	AAGTAACAGC [A/G] GTGTTGCCAA	M	A	G	S	G
G805u2	WIAF-13417	U14755		LHX1, LIM homeobox protein 1 743	GGCAGGAAC [T/C] CTACATCATC	M	T	C	L	P
G805u3	WIAF-13428	U14755		LHX1, LIM homeobox protein 1 639	GCCGTCAGGG [C/A] ATCTCCCTTA	S	C	A	G	G
G806u1	WIAF-11886	AF026547		CSPG3, chondroitin sulfate proteoglycan 3 (neurocan) 2656	TTGAGTTCC [A/G] GCCATGTCTA	S	A	G	P	P

000000-2246560

G806u2	WIAF-11895	AF026547	529	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	TGACCTTCGC [T/C] GAGGCCCAGG	S	T	C	A	A
G806u3	WIAF-11896	AF026547	477	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	GAGGTGACAG [G/A] TGTGTGTTTC	M	G	A	G	D
G806u4	WIAF-11917	AF026547	89	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	ACAGGATATC [A/G] CCGATGCCAG	M	A	G	T	A
G806u5	WIAF-11918	AF026547	213	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	AGCGCAGCCC [G/C] AGATGCCCT	M	G	C	R	P
G806u6	WIAF-11929	AF026547	769	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	GCTTGCCCG [G/A] GAGCTGGGG	S	G	A	R	R
G806u7	WIAF-11931	AF026547	3148	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	ACATTGATGA [C/T] TGCCTCTGCA	S	C	T	D	D
G806u8	WIAF-11949	AF026547	209	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	GCCAAGCGCA [G/A] CCCGAGATGC	M	G	A	A	T
G806a9	WIAF-13114	AF026547	3430	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	ATGAAAAACAC [G/A] TGGATCGGCC	S	G	A	T	T
G806u10	WIAF-13420	AF026547	2113	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	CCAGGGCAGA [C/G] TTCAGAGAAA	M	C	G	D	E
G806u11	WIAF-13431	AF026547	94	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	ATATCACCGA [T/G] GCCAGCGAAA	M	T	G	D	E
G806u12	WIAF-13432	AF026547	275	CSPG3, chondroitin sulfate proteoglycan 3 (neurocan)	ACAGGACTTG [C/T] CCATCCTGGT	M	C	T	P	S
G808a1	WIAF-13117	Y13276	177	TLX, tailless homolog (Drosophila)	GCATGAGCAA [G/a] CCAGCCGGAT	S	G	a	K	K
G810u1	WIAF-11890	X98248	990	SORT1, sortilin 1	ATAAGGATAC [C/A] ACAAGAAGGA	S	C	A	T	T
G810u2	WIAF-11891	X98248	1093	SORT1, sortilin 1	GGCAGCAAT [G/T] ATGACATGGT	M	G	T	D	Y
G810u3	WIAF-11907	X98248	1683	SORT1, sortilin 1	GACAGGAAG [T/G] CAATGCTGCC	S	T	G	G	G
G810u4	WIAF-11908	X98248	1433	SORT1, sortilin 1	ATCTCCGAGA [A/C] ACTGAATGTT	M	A	C	K	T
G810u5	WIAF-11909	X98248	1354	SORT1, sortilin 1	GAAGCCTGAA [A/G] ACAGTGAATG	M	A	G	N	D
G810u6	WIAF-11910	X98248	2180	SORT1, sortilin 1	TACCGGAAA [T/A] TCCAGGGGAC	M	T	A	I	N
G810u7	WIAF-11911	X98248	2264	SORT1, sortilin 1	AACTTTTGA [G/A] TCCGAAAAA	M	G	A	S	N

004060" 24465960

G810u8	WIAF-11925	X98248	1993	SORT1, sortilin 1	TCGAGACTAT [G/A] TTGTGACAA	M	G	A	V	I
G810u9	WIAF-11939	X98248	1351	SORT1, sortilin 1	GAGGAGCCT [G/C] AAAACAGTGA	M	G	C	E	Q
G810u10	WIAF-11940	X98248	2232	SORT1, sortilin 1	AAGTAAAGA [C/T] TTGAAAAAGA	S	C	T	D	D
G810a11	WIAF-13115	X98248	1769	SORT1, sortilin 1	TCCATGAATA [T/A] CAGCATTGG	M	T	A	I	N
G810a12	WIAF-13116	X98248	1757	SORT1, sortilin 1	CCTGGAGCTA [G/A] GTCCATGAAT	M	G	A	R	K
G811u1	WIAF-11893	HT3676	900	synapsin I, alt. transcript 1	TGACCAAGAC [G/A] TATGCCACTG	S	G	A	T	T
G811u2	WIAF-11894	HT3676	758	synapsin I, alt. transcript 1	ACCTTCTACC [C/T] CAATCACAAA	M	C	T	P	L
G811u3	WIAF-11927	HT3676	996	synapsin I, alt. transcript 1	CGTCAGTGTC [A/T] GGGAACTGGA	S	A	T	S	S
G811u4	WIAF-11928	HT3676	1054	synapsin I, alt. transcript 1	CATGTCIGAC [A/G] GATACAAAGCT	M	A	G	R	G
G811u5	WIAF-13418	HT3676	249	synapsin I, alt. transcript 1	TGTCCAACGC [G/A] GTCAAGCAGA	S	G	A	A	A
G811u6	WIAF-13419	HT3676	432	synapsin I, alt. transcript 1	TTAAAGTAGA [G/A] CAGGCCGGAAT	S	G	A	E	E
G812u1	WIAF-11898	HT4564	163	STX1A, syntaxin 1A (brain)	CCAAACCCGA [T/C] GAGAAGACGA	S	T	C	D	D
G812u2	WIAF-11942	HT4564	604	STX1A, syntaxin 1A (brain)	TACACGACAT [G/T] TTCATGGACA	M	G	T	M	I
G813u1	WIAF-11934	U72508	939	Human B7 mRNA, complete cds.	TATGACAGAG [G/A] ACAGAGGATG	M	G	A	G	E
G813u2	WIAF-11948	U72508	619	Human B7 mRNA, complete cds.	GCATCCACAT [G/C] GTGACAGGTC	M	G	C	M	I
G816u1	WIAF-11897	HT4230	151	HTR2B, 5-hydroxytryptamine (serotonin) receptor 2B	CTAACTGGTC [T/G] GGATTACAGA	S	T	G	S	S
G816u2	WIAF-11930	HT4230	189	HTR2B, 5-hydroxytryptamine (serotonin) receptor 2B	GAAATGAAC [A/G] GATTGTTGAG	M	A	G	Q	R
G818u1	WIAF-11902	HT2694	753	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	GAGTTTTC [C/T] TGCACTCAAT	S	C	T	H	H
G818u2	WIAF-11903	HT2694	775	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	TGTGAGACAC [A/G] GTTCAGATCC	M	A	G	S	G
G818u3	WIAF-11904	HT2694	1211	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	TATAATCCAT [A/C] TACACGGAGT	M	A	C	Y	S

000000 2445960

G818u4	WIAF-11905	HT2694	1081	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	GATTACCTGC [A/C] AACAGGAATG	M	A	C	K	Q
G818u5	WIAF-11933	HT2694	795	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	CCTTCTATAC [C/T] CCAGAGCCAG	S	C	T	T	T
G818u6	WIAF-11935	HT2694	1239	TPH, tryptophan hydroxylase (tryptophan 5-monooxygenase)	TCCTGAAAGA [C/T] ACCAAGAGCA	S	C	T	D	D
G822u1	WIAF-11906	HT0207	936	ASMT, acetylserotonin N- methyltransferase	CAGACGGAAA [G/T] TGCTCACACC	M	G	T	K	N
G822u2	WIAF-11919	HT0207	637	ASMT, acetylserotonin N- methyltransferase	TGCTGGGACA [C/T] GGATAAAGCT	M	C	T	R	W
G822u3	WIAF-11936	HT0207	318	ASMT, acetylserotonin N- methyltransferase	GAAAAGCTTT [C/T] TATCGAACA	S	C	T	F	F
G822u4	WIAF-11937	HT0207	116	ASMT, acetylserotonin N- methyltransferase	AATGACTACG [C/T] CAACGGCTTC	M	C	T	A	V
G822u5	WIAF-11938	HT0207	930	ASMT, acetylserotonin N- methyltransferase	ACTGGGCAGA [C/T] GGAAAGTGCT	S	C	T	D	D
G822u6	WIAF-13427	HT0207	120	ASMT, acetylserotonin N- methyltransferase	ACTAGGCCAA [C/A] GGCTTCATGG	M	C	A	N	K
G825u1	WIAF-11888	HT4974	236	ADAR, adenosine deaminase, RNA- specific	GCTCAGATAC [C/T] AGCAGCCTGG	N	C	T	Q	*
G825u2	WIAF-11900	HT4974	3076	ADAR, adenosine deaminase, RNA- specific	TCTTTGACAA [A/G] TCCTGCAGGG	S	A	G	K	K
G825u3	WIAF-11912	HT4974	2537	ADAR, adenosine deaminase, RNA- specific	CTTCATTGGG [G/C] AGAACGAGAA	M	G	C	E	Q
G825u4	WIAF-11941	HT4974	3558	ADAR, adenosine deaminase, RNA- specific	GATGGCTATG [A/G] CCTGGAGATC	M	A	G	D	G
G825a5	WIAF-12090	HT4974	1305	ADAR, adenosine deaminase, RNA- specific	CCTGAGACCA [A/G] AAGAAACGCA	M	A	G	K	R
G825u6	WIAF-13426	HT4974	3683	ADAR, adenosine deaminase, RNA- specific	CCGCAGGGAT [C/T] TACTGAGACT	S	C	T	L	L
G826u1	WIAF-12554	X99383	2109	ADARB1, adenosine deaminase, RNA- specific, B1 (homolog of rat RED1)	AGATTACCAA [A/G] CCCAACGTGT	S	A	G	K	K
G826u2	WIAF-12566	X99383	1698	ADARB1, adenosine deaminase, RNA- specific, B1 (homolog of rat RED1)	TGTCTCGCAG [T/G] GACAAGATTG	M	T	G	S	R
G829u1	WIAF-13735	U49262	1404	DVL3, dishevelled 3 (homologous to Drosophila dsh)	GGTTGGAGG [T/C] CCGTGACTGC	M	T	C	V	A

G83u1	WIAF-10449	HT1576	1338	DNMT1, DNA (cytosine-5-)-methyltransferase 1	ATGATGACCC [G/A] TCTCTTGAAG	S	G	A	P	P
G83u2	WIAF-10450	HT1576	1871	DNMT1, DNA (cytosine-5-)-methyltransferase 1	AAGCTGGTCT [A/G] CCAGATCTTC	M	A	G	Y	C
G83u3	WIAF-10468	HT1576	928	DNMT1, DNA (cytosine-5-)-methyltransferase 1	AAATCCACAG [A/G] TTTCTGATGA	M	A	G	I	V
G83u4	WIAF-10469	HT1576	1562	DNMT1, DNA (cytosine-5-)-methyltransferase 1	AATTCGACT [C/T] GACCTATGAG	M	C	T	S	L
G83u5	WIAF-10471	HT1576	2424	DNMT1, DNA (cytosine-5-)-methyltransferase 1	GGGCCACGTC [G/A] GACCCTCTGG	S	G	A	S	S
G83u6	WIAF-10473	HT1576	3790	DNMT1, DNA (cytosine-5-)-methyltransferase 1	GTTCTTCCTC [C/T] TGGAGAAATGT	S	C	T	L	L
G83u7	WIAF-10486	HT1576	1581	DNMT1, DNA (cytosine-5-)-methyltransferase 1	AGGACCTGAT [C/A] AACAAAGATCG	S	C	A	I	I
G832u1	WIAF-12577	L13387	1129	PAFAH1B1, platelet-activating factor acetylhydrolase, isoform 1b, alpha subunit (45kD)	AGACATTTCAC [A/T] GGACACACAGAG	S	A	T	T	T
G835u1	WIAF-12555	U38276	1311	SEMA3F, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3F	CCTCTGGCTC [C/A] GTGTTCCGAG	S	C	A	S	S
G835u2	WIAF-12556	U38276	1229	SEMA3F, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3F	ACTCACTTTG [A/T] TGAGCTCCAG	M	A	T	D	V
G835u3	WIAF-12557	U38276	1473	SEMA3F, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3F	GAACCTTCAC [G/A] CCATCTATGA	S	G	A	T	T
G835a4	WIAF-13138	U38276	1726	SEMA3F, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3F	TGACCAGGAG [A/T] TGGAGGAGCT	M	A	T	M	L
G836u1	WIAF-12592	U28369	1056	SEMA3B, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3B	AACGACGTGG [G/A] CGGCCAGCGC	M	G	A	G	D
G836u2	WIAF-12609	U28369	1479	SEMA3B, sema domain, immunoglobulin domain (Ig), short basic domain, secreted, 3B	GTCCGTGCCA [C/T] TGGGGGCGC	M	C	T	T	I

G838u1	WIAF-12590	U72671	ICAM5, intercellular adhesion molecule 5, telencephalin	1107	CGCAGCTGGG [A/G] CCAAGCTCT	M	A	G	T	A
G838u2	WIAF-12591	U72671	ICAM5, intercellular adhesion molecule 5, telencephalin	966	CAGGCAGCTG [A/G] TCTGCAAGT	M	A	G	I	V
G840a1	WIAF-12109	HT961	SOS1, son of sevenless (Drosophila) homolog 1	2232	CTCAGGCAAA [T/C] GGAGTAAGCC	S	T	C	N	N
G840a2	WIAF-12110	HT961	SOS1, son of sevenless (Drosophila) homolog 1	2404	ACCGTCTGAA [C/G] TTGTAGGAG	M	C	G	L	V
G840u3	WIAF-12213	HT961	SOS1, son of sevenless (Drosophila) homolog 1	3813	CAAGGTACC [G/A] CGTCGATGCT	S	G	A	P	P
G841u1	WIAF-12153	HT97420	SMOH, smoothened (Drosophila) homolog	1372	TTTGGCTTC [C/G] TGGCCTTGG	M	C	G	L	V
G841u2	WIAF-12179	HT97420	SMOH, smoothened (Drosophila) homolog	858	CCCAGTTCAT [G/T] GATGCTGCC	M	G	T	M	I
G841u3	WIAF-12185	HT97420	SMOH, smoothened (Drosophila) homolog	1164	CTGTGAGTGG [C/G] ATTGTTTGT	S	C	G	G	G
G847u1	WIAF-12588	L41939	EPHB2, EphB2	2019	GGTCTGCAGT [G/T] GCCACCTGAA	M	G	T	G	C
G847u2	WIAF-12596	L41939	EPHB2, EphB2	1806	GTGTAACAGA [A/C] GACGGGGTT	S	A	C	R	R
G847u3	WIAF-12613	L41939	EPHB2, EphB2	2885	AGCCATCAA [G/C] ATGGGGCAGT	M	G	C	K	N
G848u1	WIAF-12685	L40636	EPHB1, EphB1	2484	GTCAACAGTA [A/G] CCTGGTGTGC	M	A	G	N	S
G848u2	WIAF-12690	L40636	EPHB1, EphB1	2020	CCTTCACTTA [T/C] GAGGATCCA	S	T	C	Y	Y
G849u1	WIAF-11920	D83492	EPHB6, EphB6	1544	ACCTGTGTGG [C/T] TCATGCAGAG	M	C	T	A	V
G849u2	WIAF-11921	D83492	EPHB6, EphB6	3301	CTTTGGGATA [C/T] TCATGTGGGA	M	C	T	L	F
G849u3	WIAF-13412	D83492	EPHB6, EphB6	1139	GAGACCTTCA [C/T] CCTTACTAC	M	C	T	T	I
G849u4	WIAF-13413	D83492	EPHB6, EphB6	1895	TTTGAGGTGC [A/C] AGGCTCAGCA	M	A	C	Q	P
G849u5	WIAF-13414	D83492	EPHB6, EphB6	2338	CTATGACCAG [G/A] CAGAAGACA	M	G	A	A	T
G849u6	WIAF-13415	D83492	EPHB6, EphB6	2567	GGGCTTTGG [C/G] CTTCTCTCTG	M	C	G	A	G
G849u7	WIAF-13422	D83492	EPHB6, EphB6	2860	GGCCATCCAG [G/A] CCCTGTGGGC	M	G	A	A	T
G849u8	WIAF-13423	D83492	EPHB6, EphB6	2782	GGAGTCAAT [G/C] GGACAGGCTC	M	G	C	G	R
G849u9	WIAF-13424	D83492	EPHB6, EphB6	3038	TTCTCAGGC [A/G] CGGGAGGGC	M	A	G	Q	R
G849u10	WIAF-13425	D83492	EPHB6, EphB6	3637	AGCCATTGGA [C/T] TGGAGTGCTA	S	C	T	L	L
G856u1	WIAF-12625	D45906	LIMK2, LIM domain kinase 2	1323	AGCTGAACCT [G/C] CTGACAGAT	S	G	C	L	L
G858u1	WIAF-12630	U65019	MADH2, MAD (mothers against decapentaplegic, Drosophila) homolog 2	864	TTTGGTGTTC [G/A] ATAGCATATT	S	G	A	S	S
G86u1	WIAF-10437	HT1701	RAD51, RAD51 (S. cerevisiae) homolog (E. coli RecA homolog)	263	TGAAGCAAAT [G/C] CAGATACTTC	M	G	C	A	P

000000 26465960

G86u2	WIAF-10465	HT1701	RAD51, RAD51 (<i>S. cerevisiae</i>) 861 homolog (E coli RecA homolog)	GCATCAGCCA [T/C] GATGGTAGAA	M	T	C	M	T
G86u3	WIAF-10466	HT1701	RAD51, RAD51 (<i>S. cerevisiae</i>) 924 homolog (E coli RecA homolog)	TACAGACACAG [A/G] CTACTCGGT	M	A	G	D	G
G864a1	WIAF-13139	X82324	POU3F4, POU domain, class 3, 183 transcription factor 4	CAGCAATGGG [C/T] ATCCCTCGG	M	C	t	H	Y
G866u1	WIAF-12637	HT0101	2576 glutamate receptor (GB:M64752)	AAATCCCGTA [G/A] TGAATCCAAG	M	G	A	S	N
G866u2	WIAF-12638	HT0101	1131 glutamate receptor (GB:M64752)	TAAACAGAAA [C/T] GTGCAGTTTA	S	C	T	N	N
G869u1	WIAF-13406	HT33620	GRIN2C, glutamate receptor, ionotropic, N-methyl D-aspartate 3627 2C	AGATCAGCAG [G/T] GTAGCCCGTG	M	G	T	R	S
G870u1	WIAF-11889	HT4468	SLC1A1, solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system 714 Xag), member 1	CAGAAGAGTC [C/G] TTTCACAGCTG	S	C	G	S	S
G870u2	WIAF-11913	HT4468	SLC1A1, solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system 314 Xag), member 1	CTACAGAAAT [T/A] CTACTTTGCT	M	T	A	F	Y
G870u3	WIAF-11914	HT4468	SLC1A1, solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system 579 Xag), member 1	AAGTCAGTAC [G/A] GTGGATGCCA	S	G	A	T	T
G870u4	WIAF-11922	HT4468	SLC1A1, solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system 706 Xag), member 1	GAATGACACA [G/A] AAGAGTCCTT	M	G	A	E	K

G870u5	WIAF-11923	HT4468			SLC1A1, solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1	978		GGAGATCAT [A/G] GAAGTTGAAG	M	A	G	I	M
G871u1	WIAF-11892	HT3187			SLC1A3, solute carrier family 1 (glial high affinity glutamate transporter), member 3	1004		TTCTCTTAAC [G/C] AAGCCATCAT	M	G	C	E	Q
G871u2	WIAF-11915	HT3187			SLC1A3, solute carrier family 1 (glial high affinity glutamate transporter), member 3	1154		TGTTGCTTA [C/T] TCATTCAAGC	M	C	T	L	F
G871u3	WIAF-11926	HT3187			SLC1A3, solute carrier family 1 (glial high affinity glutamate transporter), member 3	1412		GGCTGCCATT [T/G] TCATTGCTCA	M	T	G	F	V
G871u4	WIAF-11944	HT3187			SLC1A3, solute carrier family 1 (glial high affinity glutamate transporter), member 3	1217		AAACCCCTTGG [G/A] TTTTATTGG	M	G	A	V	I
G872u1	WIAF-13433	HT4077			SLC1A2, solute carrier family 1 (glial high affinity glutamate transporter), member 2	1271		CTGTTGGAGC [A/C] ACCATTAAAC	S	A	C	A	A
G879u1	WIAF-11899	HT28317			GRM2, glutamate receptor, metabotropic 2	1273		GACTTTTGTG [T/C] CAAGTCAAG	M	T	C	L	P
G879u2	WIAF-11932	HT28317			GRM2, glutamate receptor, metabotropic 2	2349		CTTCTATGTC [A/G] CCTCCAGTGA	M	A	G	T	A
G879u3	WIAF-13421	HT28317			GRM2, glutamate receptor, metabotropic 2	2186		ATGCAAGTAT [G/T] TTGGGCTGCG	M	G	T	M	I
G879u4	WIAF-13429	HT28317			GRM2, glutamate receptor, metabotropic 2	2567		CCCAGTTTGT [C/T] CCCACTGTGT	S	C	T	V	V
G879u5	WIAF-13436	HT28317			GRM2, glutamate receptor, metabotropic 2	2046		ACAGTGGCC [A/G] TCTGCCTGGC	M	A	G	I	V
G879u6	WIAF-13438	HT28317			GRM2, glutamate receptor, metabotropic 2	2425		GTGCTTGGCT [G/T] CCTCTTTGGG	M	G	T	C	F
G879u7	WIAF-13439	HT28317			GRM2, glutamate receptor, metabotropic 2	2463		CCTCTTCCAG [C/T] CGCAGAAGAA	M	C	T	P	S
G880u1	WIAF-12164	HT33719			GRM4, glutamate receptor, metabotropic 4	2117		AGCCCGACCT [T/G] GGCACCTGCT	S	T	G	L	L

G880u2	WIAF-12176	HT33719	GRM4, glutamate receptor, 2427 metabotropic 4	GGACCTGTGC [C/T] TCATCTGCCT	M	C	T	L	F
G880u3	WIAF-12192	HT33719	GRM4, glutamate receptor, 2372 metabotropic 4	ACCAGCGGAC [A/G] CTCGACCCCC	S	A	G	T	T
G883a1	WIAF-13140	HT48863	GRM7, glutamate receptor, 1408 metabotropic 7	ATCGCAAAATG [C/a] ACAGGACAGG	N	C	a	C	*
G883a2	WIAF-13141	HT48863	GRM7, glutamate receptor, 2027 metabotropic 7	TCCTGTCTTC [C/t] TGGCAATGTT	S	C	t	L	L
G883a3	WIAF-13147	HT48863	GRM7, glutamate receptor, 1813 metabotropic 7	TGTGCACACT [A/g] CCATGTAAGC	S	A	g	L	L
G883a4	WIAF-13148	HT48863	GRM7, glutamate receptor, 1536 metabotropic 7	TGTGCTGACT [A/t] CCGGGGTGTC	M	A	t	Y	F
G883a5	WIAF-13149	HT48863	GRM7, glutamate receptor, 2473 metabotropic 7	AAGCCAGAGG [G/a] GTTCTCAAGT	S	G	a	G	G
G883a6	WIAF-13150	HT48863	GRM7, glutamate receptor, 2434 metabotropic 7	TCATAGACTA [C/t] GATGAACACA	S	C	t	Y	Y
G884u1	WIAF-11916	U95025	GRM8, glutamate receptor, 1052 metabotropic 8	CGAACTCTTG [C/a] CAATAATCGA	M	C	A	A	D
G884u2	WIAF-11945	U95025	GRM8, glutamate receptor, 2016 metabotropic 8	AAACAAACCG [T/C] ATCCACCAGAA	S	T	C	R	R
G884u3	WIAF-11946	U95025	GRM8, glutamate receptor, 1852 metabotropic 8	GAGGGCTTCA [G/a] GACGCGAACT	M	G	A	G	R
G884u4	WIAF-11947	U95025	GRM8, glutamate receptor, 2078 metabotropic 8	ATTAGTCCAG [C/G] ATCTCAGCTG	M	C	G	A	G
G884u5	WIAF-13430	U95025	GRM8, glutamate receptor, 1897 metabotropic 8	TTTTTCTCTGT [T/G] ATTCAATCAC	M	T	G	Y	D
G884u6	WIAF-13435	U95025	GRM8, glutamate receptor, 2364 metabotropic 8	TTACCAATGTA [T/C] ACCACCTGCA	S	T	C	Y	Y
G885u1	WIAF-13434	AF002700	GFR2, GDNF family receptor alpha 1363 2	AACTCAGGCC [C/A] CAGCAGAGCC	M	C	A	P	H
G886a1	WIAF-13142	U95847	GFR1, GDNF family receptor alpha 497 1	GAAGTCGCTC [T/a] ACAACTGCCG	M	T	a	Y	N
G886a2	WIAF-13143	U95847	GFR1, GDNF family receptor alpha 1385 1	GTCTGAGAAT [G/a] AAATTTCCAC	M	G	a	E	K
G886a3	WIAF-13151	U95847	GFR1, GDNF family receptor alpha 781 1	GCGTGTCCAA [T/c] GATGTCTGCA	S	T	C	N	N
G892u1	WIAF-11956	U12140	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TGGCAATCC [A/G] TTTACATGCT	S	A	G	P	P
G892u2	WIAF-11957	U12140	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GGATCAAGAC [T/A] CTCCAAGAGG	S	T	A	T	T

G892u3	WIAF-11958	U12140	956	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GCAAAATCTGG [C/T] CGCACCCTAAC	M	C	T	A	V
G892u4	WIAF-11960	U12140	1738	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CTCCAAAGTTT [G/A] GCATGAAGG	M	G	A	G	S
G892u5	WIAF-11962	U12140	2486	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GTCGGTGGCC [A/G] CACAATGCTG	M	A	G	H	R
G892u6	WIAF-11965	U12140	1106	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TCCTTAAGGA [T/C] AACTAACATT	M	T	C	I	T
G892u7	WIAF-11966	U12140	2085	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	AGGATGCCAG [T/C] GACAAATGCAC	S	T	C	S	S
G892u8	WIAF-11967	U12140	2230	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GGACCTCAAC [A/C] AGTTCTCTCAG	M	A	C	K	Q
G892u9	WIAF-11968	U12140	2223	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	AGCATGGGGA [C/T] CTCACAAAGT	S	C	T	D	D
G892u10	WIAF-11992	U12140	1602	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GTAATGAAAT [C/T] CCTTCCACAG	S	C	T	I	I
G892u11	WIAF-11998	U12140	1354	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TACTAAATA [C/T] ATGTTACCAA	M	C	T	H	Y
G892u12	WIAF-11999	U12140	1944	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CAITTTGTTCA [G/C] CACATCAAGC	M	G	C	Q	H
G892u13	WIAF-12000	U12140	2103	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CACCAAGGA [C/T] TTCCACCGTG	S	C	T	D	D
G892u14	WIAF-12001	U12140	1860	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CTGTCAATTAT [T/C] GGAATGACCA	S	T	C	I	I
G892a15	WIAF-13144	U12140	1868	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	ATTGGAATCA [C/G] CAAGATCCCT	M	C	G	T	S
G892a16	WIAF-13145	U12140	1903	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CCAGTACTTT [G/T] GCATCACCAA	M	G	T	G	C

G892a17	WIAF-13146	U12140	1965	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GACATAACAT [T/G] GTTCTGAAAA	M	T	G	I	M
G892u18	WIAF-13442	U12140	958	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	AAATCTGGCC [G/T] CACCTAACCT	M	G	T	A	S
G892u19	WIAF-13446	U12140	2502	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TGCTGCCCAT [T/C] CGCTGGATGC	S	T	C	I	I
G892u20	WIAF-13447	U12140	2317	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GATGCTGCAT [A/T] TAGCCGACGA	M	A	T	I	L
G892u21	WIAF-13448	U12140	2364	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CGTCCAGCA [C/A] TTCGTGCACC	M	C	A	H	Q
G892u22	WIAF-13449	U12140	2507	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	CCCATTCGCT [G/A] GATGCCTCCA	N	G	A	W	*
G892u23	WIAF-13471	U12140	2389	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TTTGGCCACC [A/C] GGAACGCTCT	S	A	C	R	R
G892u24	WIAF-13472	U12140	2416	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GGAGAACTTG [C/T] TGGTGAAAT	S	C	T	L	L
G892u25	WIAF-13474	U12140	359	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	GGGATGTCGT [C/T] CTGGATAAGG	M	C	T	S	F
G892u26	WIAF-13479	U12140	1044	NTRK2, neurotrophic tyrosine kinase, receptor, type 2	TGTATTGGGA [T/C] GTTGGTAACC	S	T	C	D	D
G9u1	WIAF-10222	J03826	1130	FDXR, ferredoxin reductase	GGTATAAGAG [C/T] CGCCCTGTCTG	S	C	T	S	S
G9u2	WIAF-10258	J03826	388	FDXR, ferredoxin reductase	CCGGAGCTGC [A/G] GGAGGCCTAC	M	A	G	Q	R
G900u1	WIAF-11970	HT3470	497	STX4A, syntaxin 4A (placental)	TGCAATTCAA [T/C] GCAGTCGGAA	M	T	C	M	T
G901u1	WIAF-11969	HT27792	758	STX3A, syntaxin 3A	TGCACACAGT [G/A] GACCACGTGG	S	G	A	V	V
G901u2	WIAF-11971	HT27792	317	STX3A, syntaxin 3A	ACGTCCCGAA [C/A] AAATGAAGA	M	C	A	N	K
G901u3	WIAF-12002	HT27792	611	STX3A, syntaxin 3A	AGCAAGCCCT [C/T] AGTGAGATTG	S	C	T	L	L
G901u4	WIAF-12003	HT27792	909	STX3A, syntaxin 3A	GCTGAATTAA [G/A] AGTGGCCTAA	-	G	A	-	-
G901u5	WIAF-12004	HT27792	163	STX3A, syntaxin 3A	ATTGAGGAAA [C/T] TCGGCTTAAC	M	C	T	T	I
G901u6	WIAF-13152	HT27792	82	STX3A, syntaxin 3A	CAGCTGACAC [A/G] GGATGATCAT	M	A	G	Q	R

G901u7	WIAF-13453	HT27792	828	STX3A, syntaxin 3A	CCGGAAGAAA [T/C] TCATAATTAT	S	T	C	L	L
G901u8	WIAF-13455	HT27792	226	STX3A, syntaxin 3A	TACAGTATCA [T/C] TCCTCTGCA	M	T	C	I	T
G902u1	WIAF-13454	HT27744	848	STX5A, syntaxin 5A	ACTTCCAGTC [T/A] GTCACCTCCA	S	T	A	S	S
G902u2	WIAF-13456	HT27744	338	STX5A, syntaxin 5A	ATTTCGTGAG [A/G] GCCAAGGGCA	S	A	G	R	R
G905u1	WIAF-12202	HT27789	487	CREBL1, cAMP responsive element binding protein-like 1	TCCAGATCAA [C/T] GTTATCCCA	S	C	T	N	N
G905u2	WIAF-12219	HT27789	151	CREBL1, cAMP responsive element binding protein-like 1	ATTCTGGCCT [A/T] GATGAAGTGG	S	A	T	L	L
G905u3	WIAF-12230	HT27789	649	CREBL1, cAMP responsive element binding protein-like 1	AGTCCCTGTC [C/G] CCTTCAGGAT	S	C	G	S	S
G906u1	WIAF-12214	HT4372	2127	N-ethylmaleimide-sensitive factor	AAGGGAAGAA [G/A] GTCTGGATAG	S	G	A	K	K
G906u2	WIAF-12221	HT4372	514	N-ethylmaleimide-sensitive factor	GGGAGAGCCT [G/A] CGACAGGGAA	M	G	A	A	T
G908u1	WIAF-12201	HT3665	98	RAB5A, member RAS oncogene family	GCCAAAATAC [T/G] GGAAATAAAA	S	T	G	T	T
G91u1	WIAF-10438	HT1848	496	ERCC1, excision repair cross-complementing rodent repair deficiency, complementation group 1 (includes overlapping antisense 1 sequence)	TCGTGCGCAA [C/T] GTGCCTCTGGG	S	C	T	N	N
G91u2	WIAF-10439	HT1848	367	ERCC1, excision repair cross-complementing rodent repair deficiency, complementation group 1 (includes overlapping antisense 1 sequence)	CTGGGGCCAC [G/A] TGCCCCACAG	S	G	A	T	T
G914a1	WIAF-13210	HT3672	252	synaptobrevin 1	GCAGTGCTGC [C/A] AAGCTAAGA	S	C	A	A	A
G915a1	WIAF-12115	D63506	1390	Homo sapiens mRNA for unc-18homologue, complete cds.	TTACCTTGGT [G/A] TTCCCATTTG	M	G	A	V	I
G915u2	WIAF-12293	D63506	685	Homo sapiens mRNA for unc-18homologue, complete cds.	ACAGCTTGT [G/A] AAAAAAAGCT	M	G	A	E	K
G916a1	WIAF-13209	HT28523	308	Huntingtin associated protein 1-like protein	GAGCAGTTTT [C/T] GGAGGCCAGC	M	C	T	S	L
G916a2	WIAF-13211	HT28523	762	Huntingtin associated protein 1-like protein	CGGAGGAGTT [G/C] GTGCCCCAGG	M	G	C	L	F
G916a3	WIAF-13212	HT28523	560	Huntingtin associated protein 1-like protein	GAGCTCAGAA [C/T] GTCTCTAAGG	M	C	T	T	M

000000 2445960

G917u1	WIAF-11972	U79734	1075	HIP1, huntingtin interacting protein 1	AGAGCCAGCG [G/A] GTTGTGCTGC	S	G	A	R	R
G917u2	WIAF-11973	U79734	1005	HIP1, huntingtin interacting protein 1	GACCACTTAA [T/C] TGAGCGACTA	M	T	C	I	T
G917u3	WIAF-11977	U79734	1539	HIP1, huntingtin interacting protein 1	CTGCAAGGCA [G/A] CCTGGAAACT	M	G	A	S	N
G917u4	WIAF-12005	U79734	817	HIP1, huntingtin interacting protein 1	TGGTGTGTAT [C/T] CCTGCAGAGG	S	C	T	I	I
G917u5	WIAF-12006	U79734	1906	HIP1, huntingtin interacting protein 1	GCTGGAGCCA [G/C] TAATCTGGCCT	M	G	C	Q	H
G917a6	WIAF-13157	U79734	993	HIP1, huntingtin interacting protein 1	AAGGATGAGA [A/G] GGACCACTTA	M	A	G	K	R
G919u1	WIAF-11974	D30742	707	CAMK4, calcium/calmodulin-dependent protein kinase IV	ACTGGGCACC [T/C] GAAATTCCTTA	S	T	C	P	P
G919u2	WIAF-11991	D30742	1139	CAMK4, calcium/calmodulin-dependent protein kinase IV	AGAGCCACAA [G/A] GCTAGCCGAG	S	G	A	K	K
G919u3	WIAF-12007	D30742	834	CAMK4, calcium/calmodulin-dependent protein kinase IV	CATGTTCCAGG [A/T] GAAATCTGAA	N	A	T	R	*
G919u4	WIAF-13443	D30742	1088	CAMK4, calcium/calmodulin-dependent protein kinase IV	TGGCCTCTTC [C/G] CGCCTGGGAA	S	C	G	S	S
G920u1	WIAF-11979	X78520	1952	CLCN3, chloride channel 3	ATGACATTC [T/C] GATCGTCCAG	S	T	C	P	P
G920u2	WIAF-11980	X78520	1819	CLCN3, chloride channel 3	ATAGCCTTCC [C/T] TAATCCATAC	M	C	T	P	L
G920u3	WIAF-11981	X78520	2094	CLCN3, chloride channel 3	CATTGGAGCG [A/G] TCGCAGGAAG	M	A	G	I	V
G920u4	WIAF-11983	X78520	2822	CLCN3, chloride channel 3	ATATTTTCCG [A/G] AAGCTGGGAC	S	A	G	R	R
G920u5	WIAF-11984	X78520	2745	CLCN3, chloride channel 3	GCCATTGAAG [C/T] TTCGAAGCAT	M	C	T	L	F
G920u6	WIAF-11987	X78520	2499	CLCN3, chloride channel 3	TCCCTTAGCT [G/T] TCCTGACACA	M	G	T	V	F
G920u7	WIAF-12008	X78520	1251	CLCN3, chloride channel 3	CATCATCAGA [G/A] GTTACTTGGG	M	G	A	G	S
G920u8	WIAF-12011	X78520	888	CLCN3, chloride channel 3	AGTAGTACA [C/T] TAACAGGATT	S	C	T	L	L
G920u9	WIAF-13459	X78520	2804	CLCN3, chloride channel 3	CAATGGAGAT [T/C] GTGGTGGATA	S	T	C	I	I
G921u1	WIAF-11954	J02908	931	CLU, clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)	GAGAGGTTGA [C/T] CAGGAATAC	M	C	T	T	I

G921u2	WIAF-11955	J02908		880	CLU, clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)	CCCTCCACGG [C/T] TAAGCTCCGG	M	C	T	A	V
G921u3	WIAF-11990	J02908		1051	CLU, clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)	CTCACGCAAG [G/C] CGAAGACCAG	M	G	C	G	A
G921u4	WIAF-13469	J02908		986	CLU, clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)	TCAACACCTC [C/T] TCCTTGCTGG	S	C	T	S	S
G923u1	WIAF-11993	M19650		1059	Human 2',3'-cyclic nucleotide 3'-phosphodiesterase mRNA, complete cds.	GAGCTAAGCC [G/A] GGGCAAGCTC	M	G	A	R	Q
G923u2	WIAF-11994	M19650		1062	Human 2',3'-cyclic nucleotide 3'-phosphodiesterase mRNA, complete cds.	CTAAGCCGGG [G/T] CAAGCTCTAT	M	G	T	G	V
G923u3	WIAF-13445	M19650		1141	Human 2',3'-cyclic nucleotide 3'-phosphodiesterase mRNA, complete cds.	TCCTCACGGG [G/A] TACTACGGGA	S	G	A	G	G
G925u1	WIAF-11953	L11315		666	CAK, cell adhesion kinase	GGGTCAATGAG [T/C] GTCTGTCTGC	S	T	C	S	S
G925u2	WIAF-11959	L11315		2562	CAK, cell adhesion kinase	TGCTGCCAT [C/T] CGCTGGATGG	S	C	T	I	I
G925u3	WIAF-11996	L11315		2049	CAK, cell adhesion kinase	AAGATCTGGT [T/C] AGTCTTGATT	S	T	C	V	V
G925u4	WIAF-13440	L11315		1601	CAK, cell adhesion kinase	TACAGAGAGC [C/T] CCGGCTCGT	M	C	T	P	L
G925u5	WIAF-13441	L11315		1629	CAK, cell adhesion kinase	CGCCCCACTC [C/T] GCTCCCTGTG	S	C	T	S	S
G925u6	WIAF-13451	L11315		2262	CAK, cell adhesion kinase	TGGAGAACGG [C/T] GACCTCAACC	S	C	T	G	G
G926u1	WIAF-11961	AF018956		577	NRP1, neuropilin 1	TGAAAGCTTT [G/T] ACCTGGAGCC	M	G	T	D	Y
G926u2	WIAF-11963	AF018956		1683	NRP1, neuropilin 1	CCAGCGGATT [C/G] ATCAGGATCT	M	C	G	F	L
G926u3	WIAF-11975	AF018956		2176	NRP1, neuropilin 1	GACCTTCTGG [T/C] ATCATCATGTC	M	T	C	Y	H

G926u4	WIAF-11976	AF018956	2092	NRP1, neuropilin 1	TTCCCAAGCT [G/T] ACGAAATCA	M	G	T	D	Y
G926a5	WIAF-13158	AF018956	747	NRP1, neuropilin 1	TTTTTTACAC [C/T] GACAGCGGA	S	C	T	T	T
G926a6	WIAF-13159	AF018956	996	NRP1, neuropilin 1	ACTTGGGCTT [T/C] CTGCGTTTG	S	T	C	L	L
G926u7	WIAF-13444	AF018956	644	NRP1, neuropilin 1	GAAATCTGGG [A/C] TGGATTCCCT	M	A	C	D	A
G926u8	WIAF-13450	AF018956	1738	NRP1, neuropilin 1	CAGAATGGAG [C/G] TCGTGGGCTG	M	C	G	L	V
G926u9	WIAF-13452	AF018956	537	NRP1, neuropilin 1	TTGCTTTTGC [G/A] CCAAGATGT	S	G	A	A	A
G926u10	WIAF-13457	AF018956	2197	NRP1, neuropilin 1	TGGTCCCAC [G/A] TCGGCACACT	M	G	A	V	I
G927u1	WIAF-11978	AF022860	870	NRP2, neuropilin 2	GGATTGCTAA [T/C] GAACAGATCA	S	T	C	N	N
G927u2	WIAF-11982	AF022860	1674	NRP2, neuropilin 2	ATGACACCCC [T/G] GACATCGAA	S	T	G	P	P
G927u3	WIAF-11985	AF022860	1250	NRP2, neuropilin 2	TGGCACTCAG [G/A] TATCGCCCTC	M	G	A	G	D
G927u4	WIAF-11986	AF022860	1071	NRP2, neuropilin 2	ATGGCTACTA [C/T] GTCAAATCTT	S	C	T	Y	Y
G927u5	WIAF-12009	AF022860	726	NRP2, neuropilin 2	GTTTCATCGAC [G/A] GGGATCCTCT	S	G	A	T	T
G927u6	WIAF-12010	AF022860	2522	NRP2, neuropilin 2	GCAACCTCAG [G/T] GTCTGGGCCC	M	G	T	G	V
G927u7	WIAF-12012	AF022860	123	NRP2, neuropilin 2	GCTATATCAC [C/T] TCTCCCGGTT	S	C	T	T	T
G927a8	WIAF-13160	AF022860	2427	NRP2, neuropilin 2	CTTTTCAGT [G/T] GACATCCGAG	S	G	T	V	V
G927a9	WIAF-13161	AF022860	2430	NRP2, neuropilin 2	TTGCAGTGGG [C/G] ATCCCGAGAA	M	C	G	D	E
G927a10	WIAF-13162	AF022860	2463	NRP2, neuropilin 2	AAGGATATGA [A/G] GATGAATTTG	S	A	G	E	E
G927a11	WIAF-13163	AF022860	2473	NRP2, neuropilin 2	AGATGAATTT [G/T] ATGATGAATA	M	G	T	D	Y
G927u12	WIAF-13480	AF022860	724	NRP2, neuropilin 2	TCGTTTCATCG [A/T] CGGGGATCCT	M	A	T	T	S
G927u13	WIAF-13481	AF022860	767	NRP2, neuropilin 2	ATGGCGGTGG [C/T] CAAGGATGGC	M	C	T	A	V
G930a1	WIAF-13164	HT2608	609	GABRA2, gamma-aminobutyric acid (GABA) A receptor, alpha 2	ACAATGGGAA [G/a] AAATCAGTAG	S	G	a	K	K
G931a1	WIAF-13153	HT2609	1111	GABRA3, gamma-aminobutyric acid (GABA) A receptor, alpha 3	ACTGGTTTCAT [A/g] GCCGTCCTGTT	M	A	g	I	M
G931a2	WIAF-13165	HT2609	1448	GABRA3, gamma-aminobutyric acid (GABA) A receptor, alpha 3	TGTCAGCAAG [G/A] TTGACAAAAT	M	G	A	V	I
G932a1	WIAF-13154	HT27773	1077	GABRA4, gamma-aminobutyric acid (GABA) A receptor, alpha 4	CAAAAGAAAAG [A/G] CATCAAAGCC	M	A	G	T	A
G932a2	WIAF-13155	HT27773	1189	GABRA4, gamma-aminobutyric acid (GABA) A receptor, alpha 4	AGAACAAAATG [C/A] TTTGGTTCAC	M	C	A	A	D
G936u1	WIAF-12308	HT3432	1027	GABRB2, gamma-aminobutyric acid (GABA) A receptor, beta 2	AATTACGATG [C/T] TTCAGCTGCA	M	C	T	A	V
G936u2	WIAF-12327	HT3432	362	GABRB2, gamma-aminobutyric acid (GABA) A receptor, beta 2	AAGGCTATGA [C/T] ATTCTGTCTGA	S	C	T	D	D

G936u3	WIAF-12328	HT3432	571	GABR2, gamma-aminobutyric acid (GABA) A receptor, beta 2	CTCTGGGTGC [C/T] TGATACCTAT	M	C	T	P	L
G939u1	WIAF-12330	HT2236	1219	GABR2, gamma-aminobutyric acid (GABA) receptor, rho 2	CTGGATGGAA [G/C] CTACAGTGAG	M	G	C	S	T
G939u2	WIAF-12355	HT2236	1003	GABR2, gamma-aminobutyric acid (GABA) receptor, rho 2	ACCACCATCA [T/C] CACGGGCGTG	M	T	C	I	T
G939u3	WIAF-12356	HT2236	1041	GABR2, gamma-aminobutyric acid (GABA) receptor, rho 2	CGTCTCCTAC [G/A] TCAAGGCGGT	M	G	A	V	I
G950u1	WIAF-13622	U64871	785	Human putative G protein-coupled receptor (GPR19) gene, complete cds.	GTCTGTCTCC [A/C] GTTCACCACT	M	A	C	Q	P
G950u2	WIAF-13624	U64871	443	Human putative G protein-coupled receptor (GPR19) gene, complete cds.	GATAACAGCA [A/C] GCCACATTGT	M	A	C	K	T
G950u3	WIAF-13625	U64871	818	Human putative G protein-coupled receptor (GPR19) gene, complete cds.	CTGGGTAGTG [C/T] AACGTGCAAG	M	C	T	A	V
G955a1	WIAF-13166	HT3860	5110	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	CTGGCCTCTT [T/C] ACCGTGGAGA	S	T	c	F	F
G955a2	WIAF-13167	HT3860	3842	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	CTACCCCAAC [C/a] CAGAAACTAC	M	C	a	P	T
G955a3	WIAF-13168	HT3860	5624	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	GTGTGCCCA [G/a] AGTCGAGCC	M	G	a	E	K
G955a4	WIAF-13169	HT3860	5703	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	ATCAGCTTCT [A/g] CATGCTCTGT	M	A	g	Y	C
G955a5	WIAF-13170	HT3860	5809	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	ACCACCTGGA [T/c] GAGTTAAAA	S	T	c	D	D
G955a6	WIAF-13171	HT3860	6616	calcium channel, voltage-gated, alpha 1 subunit, L type, alt. transcript 1	CCGGCTCCAA [C/t] GCCAACATCA	S	C	t	N	N
G956u1	WIAF-14187	HT2199	1334	calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	CTTCAATAG [C/T] CCTTTGGTA	M	C	T	A	V

002060" 24425960

G956u2	WIAF-14188	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	1452	AAGAGACCC [A/T] GCTCCATGTG	S A T P P
G956u3	WIAF-14189	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	1614	GCTGGACAGA [C/T] GTCCTCTACT	S C T D D
G956u4	WIAF-14190	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	2540	GGCAAGTTTA [A/T] TTTTGATGAA	M A T N I
G956u5	WIAF-14191	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	3210	TGCTGAGCAG [T/C] GCTGCCTGTG	S T C S S
G956u6	WIAF-14192	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	3326	TTGAAGATGA [C/T] AACTTTTGGGA	M C T T I
G956u7	WIAF-14193	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	3274	ACTGGGTTAC [T/C] TTGACTATGC	M T C F L
G956u8	WIAF-14194	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	5127	TGCCTCTCAA [C/T] AGTGACGGGA	S C T N N
G956u9	WIAF-14195	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	5173	TGCTTTGGTT [C/T] GAACGGCTCT	N C T R *
G956u10	WIAF-14200	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	1437	CAGATATCGT [A/G] GCTCAAGAGG	S A G V V
G956u11	WIAF-14201	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	2567	ACCAAGCGGA [G/T] CACCTTTGAC	M G T S I
G956u12	WIAF-14202	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	4464	TCACCTTTTT [C/T] CGTCTTTTCC	S C T F F
G956u13	WIAF-14215	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	6927	GCTACAGCGA [C/T] CAAGAGCCAG	S C T D D
G956u14	WIAF-14216	HT2199		calcium channel, voltage-gated, alpha 1D subunit, DHP-sensitive	6858	CCCGAGCCAA [C/T] GGGGATGTGG	S C T N N
G957u1	WIAF-12306	HT4229		calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	915 2	TACATCGAGC [G/A] TGCCTTCATGA	M G A ? R

G957u2	WIAF-12309	HT4229	3555 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	GCCTACTACAT [C/T] GTGAACCTGC	S	C	T	I	I
G957u3	WIAF-12310	HT4229	4116 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	ATCTAGATCA [C/T] GAGAAAAACA	S	C	T	H	H
G957u4	WIAF-12313	HT4229	5181 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	AGAACGAGAA [T/C] GAAGCTGCG	S	T	C	N	N
G957u5	WIAF-12314	HT4229	5971 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	TATGGACCCC [G/A] CCGATGACGG	S	G	A	T	T
G957u6	WIAF-12315	HT4229	5985 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	ATGACGGACA [G/T] TTCCAAGAAC	M	G	T	Q	H
G957u7	WIAF-12329	HT4229	3100 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	GCTGGCAGGA [G/A] GCCTTGATGA	M	G	A	G	S
G957u8	WIAF-12331	HT4229	6492 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	CCCTCCTTTC [C/T] TACAGCTCCC	M	C	T	?	R
G957u9	WIAF-12354	HT4229	3839 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	AACGCTTTGG [G/C] AACCAACAAA	M	G	C	G	A
G957u10	WIAF-12357	HT4229	4753 2	calcium channel, voltage-gated, alpha 1E subunit, alt. transcript	TGACTTTCATC [A/G] CCGTGATTGG	M	A	G	T	A
G960u1	WIAF-12305	HT3336	1246	CACNB3, calcium channel, voltage-dependent, beta 3 subunit	TTGATGCCCT [C/T] TGATGAGGCC	M	C	T	S	F
G960u2	WIAF-12340	HT3336	1288	CACNB3, calcium channel, voltage-dependent, beta 3 subunit	TGGACAGGAT [C/T] TTCACAGCCT	M	C	T	S	F
G960u3	WIAF-12345	HT3336	641	CACNB3, calcium channel, voltage-dependent, beta 3 subunit	AGGCTCTCTT [C/T] GACTTCCTCA	S	C	T	F	F
G960u4	WIAF-12346	HT3336	576	CACNB3, calcium channel, voltage-dependent, beta 3 subunit	CATGCGGCTT [G/A] TGGTGTGCTGT	M	G	A	V	M
G961u1	WIAF-12322	U95019	2037	CACNB2, calcium channel, voltage-dependent, beta 2 subunit	ACTCTGCCTA [C/T] GTAGAGCCAA	S	C	T	Y	Y

G961u2	WIAF-12347	U95019		2007	CACNB2, calcium channel, voltage-dependent, beta 2 subunit	CATTTGACTC [G/A] GAAACCCAGG	S	G	A	S	S
G962u1	WIAF-12324	U95020		1423	CACNB4, calcium channel, voltage-dependent, beta 4 subunit	CCAATTGAAA [G/A] ACGAAGTCTA	M	G	A	R	K
G962u2	WIAF-12342	U95020		167	CACNB4, calcium channel, voltage-dependent, beta 4 subunit	GGAGCAGGTT [G/T] AAAAGATCCG	M	G	T	L	F
G962u3	WIAF-12350	U95020		1571	CACNB4, calcium channel, voltage-dependent, beta 4 subunit	ACACTTACAA [A/G] CCCCATAGGA	S	A	G	K	K
G965u1	WIAF-12312	U40583		1276	CHRNA7, cholinergic receptor, nicotinic, alpha polypeptide 7	TCCTGCACGG [T/C] GGGCAACCCC	S	T	C	G	G
G968a1	WIAF-12119	HT27592		1008	CHRNA1, cholinergic receptor, nicotinic, alpha polypeptide 1 (muscle)	ACACACACCA [C/T] CGCTCACCCA	S	C	T	H	H
G968u2	WIAF-12368	HT27592		1136	CHRNA1, cholinergic receptor, nicotinic, alpha polypeptide 1 (muscle)	AAGATTTTTA [C/T] AGAAGACATT	M	C	T	T	I
G973a1	WIAF-13172	HT48774		800	CHRNA2, cholinergic receptor, nicotinic, alpha polypeptide 2 (neuronal)	ACACTTCAGA [C/t] GTGGTGATTG	S	C	t	D	D
G973a2	WIAF-13173	HT48774		927	CHRNA2, cholinergic receptor, nicotinic, alpha polypeptide 2 (neuronal)	CTGGAAACCCC [G/a] CTGATTTTGG	M	G	a	A	T
G977u1	WIAF-13949	Y08419		366	CHRNA5, cholinergic receptor, nicotinic, alpha polypeptide 5	AAGTTATACG [T/C] GTTCCTTCAG	S	T	C	R	R
G978a1	WIAF-13179	Y08417		1331	CHRNA3, cholinergic receptor, nicotinic, beta polypeptide 3	CCATTAGATA [C/a] ATTTCGAGAC	N	C	a	Y	*
G983a1	WIAF-13214	HT0374		236	NPY, neuropeptide Y	GATACTACTC [G/A] CGGTGCGAC	S	G	A	S	S
G983a2	WIAF-13215	HT0374		290	NPY, neuropeptide Y	GAAAACGATC [C/T] AGCCACAGAG	S	C	T	S	S
G983a3	WIAF-13216	HT0374		111	NPY, neuropeptide Y	GCGACTGGGG [C/T] TGTCGGACT	S	C	T	L	L
G987a1	WIAF-13174	HT27830		159	PPYR1, pancreatic polypeptide receptor 1	TGGTCTTCAT [C/T] GTCACCTCCT	S	C	T	I	I
G987a2	WIAF-13175	HT27830		222	PPYR1, pancreatic polypeptide receptor 1	TGATGTGTGT [G/A] ACTGTGAGGC	S	G	A	V	V
G987a3	WIAF-13176	HT27830		322	PPYR1, pancreatic polypeptide receptor 1	GCCGCTGACC [G/T] CCGTCTACAC	M	G	T	A	S

20250624 14:59:50

G987a4	WIAF-13177	HT27830	1074	PPYR1, pancreatic polypeptide receptor 1	TGGAGGAGTC [G/A] GAGCATCTGC	S	G	A	S	S
G987a5	WIAF-13178	HT27830	975	PPYR1, pancreatic polypeptide receptor 1	CCTCCACCTG [C/T] GTCACCCAT	S	C	T	C	C
G987a6	WIAF-13180	HT27830	615	PPYR1, pancreatic polypeptide receptor 1	AGTTCTCTGGC [A/G] GATAAGTGG	S	A	G	A	A
G987a7	WIAF-13181	HT27830	718	PPYR1, pancreatic polypeptide receptor 1	GGGCTTCATC [C/T] TGCTCTGTTA	S	C	T	L	L
G987a8	WIAF-13182	HT27830	745	PPYR1, pancreatic polypeptide receptor 1	CATCTACCGG [C/T] GCCTGGACAG	M	C	t	R	C
G987a9	WIAF-13183	HT27830	842	PPYR1, pancreatic polypeptide receptor 1	GTGATGGTGG [T/A] GGCCTTTGCC	M	T	A	V	E
G987a10	WIAF-13184	HT27830	852	PPYR1, pancreatic polypeptide receptor 1	TGGCCTTTGC [C/T] GTGCTCTGGC	S	C	T	A	A
G987a11	WIAF-13185	HT27830	889	PPYR1, pancreatic polypeptide receptor 1	CAACAGCCTG [G/a] AAGACTGGCA	M	G	a	E	K
G987a12	WIAF-13186	HT27830	924	PPYR1, pancreatic polypeptide receptor 1	CCATCTGCCA [C/T] GGGAACTCA	S	C	T	H	H
G989u1	WIAF-13573	D86519	891	NPY6R, neuropeptide Y receptor Y6	TGACTCATGC [C/T] TACTGGGCA	S	C	T	A	A
G989u2	WIAF-13588	D86519	465	NPY6R, neuropeptide Y receptor Y6	ACCACCCAGC [A/G] TCTAATACAA	S	A	G	A	A
G989u3	WIAF-13591	D86519	980	NPY6R, neuropeptide Y receptor Y6	GAGCCCTTCC [G/A] CAACCTCTCT	M	G	A	R	H
G991u1	WIAF-12390	HT97376	336	Notch2	AAGTACTTG [C/T] GTTCAGAAA	S	C	T	C	C
G993u1	WIAF-12359	U95299	1343	NOTCH4, Notch (Drosophila) homolog 4	TCCACACTCT [G/T] CCTGTGTCTAG	M	G	T	C	F
G993u2	WIAF-12361	U95299	2020	NOTCH4, Notch (Drosophila) homolog 4	TAAGGACCAG [A/G] AAGACAAGGC	M	A	G	K	E
G993u3	WIAF-12384	U95299	5775	NOTCH4, Notch (Drosophila) homolog 4	GGGCCTATTG [G/T] CATTCGCCGA	S	G	T	S	S
G996a1	WIAF-13213	HT3329	356	OPRM1, opioid receptor, mu 1	CTTAGATGGC [A/G] ACCTGTCCGA	M	A	G	N	D
LPLa4	WIAF-13314	HT1320	443	LPL, lipoprotein lipase	ATGTATGAGA [G/T] TTGGGTGCCA	M	G	T	S	I
LPLa5	WIAF-13315	HT1320	579	LPL, lipoprotein lipase	GACAGGATGT [G/A] GCCCGGTTTA	S	G	A	V	V
LPLa6	WIAF-13316	HT1320	609	LPL, lipoprotein lipase	TGGAGGAGGA [G/A] TTTAACTACC	S	G	A	E	E
LPLa7	WIAF-13317	HT1320	1338	LPL, lipoprotein lipase	CAATAAGAC [C/A] TACTCTCTCC	S	C	A	T	T
LPLa8	WIAF-13318	HT1320	1117	LPL, lipoprotein lipase	CAATCTGGGC [T/G] ATGAGATCAA	M	T	G	Y	D
LPLa9	WIAF-13319	HT1320	715	LPL, lipoprotein lipase	CAGAATTACT [G/A] GCCTCGATCC	M	G	A	G	S
LPLa10	WIAF-13320	HT1320	834	LPL, lipoprotein lipase	CTGCTCGAAG [C/A] ATTGGAATCC	M	C	A	S	R
LPLa11	WIAF-13321	HT1320	951	LPL, lipoprotein lipase	GACTTGGAGA [T/A] GTGGACCAGC	M	T	A	D	E
LPLa12	WIAF-13322	HT1320	1595	LPL, lipoprotein lipase	AATAAGAACT [C/G] AGGCTGAAC	N	C	G	S	*

LPLa13	WIAF-13323	HT1320	1597 LPL, lipoprotein lipase	TAAGAAGTCA [G/A] GCTGAAACTG	M	G	A	G	S
LPLa14	WIAF-13324	HT1320	1606 LPL, lipoprotein lipase	AGGCTGAAAC [T/C] GGGCGAATCT	-	T	C	-	-
LPLa15	WIAF-13325	HT1320	1611 LPL, lipoprotein lipase	GAAACTGGGC [G/A] AATCTACAGA	-	G	A	-	-

002060-2445500

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

[illegible]